

Sildenafil 25 mg film-coated tablets
Sildenafil 50 mg film-coated tablets
Sildenafil 100 mg film-coated tablets
 Sildenafil citrate

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Sildenafil is and what it is used for
2. What you need to know before you take Sildenafil
3. How to take Sildenafil
4. Possible side effects
5. How to store Sildenafil
6. Contents of the pack and other information

1. What Sildenafil is and what it is used for

Sildenafil belongs to a group of medicines called **phosphodiesterase type 5 (PDE 5) inhibitors**. It works by helping to relax the blood vessels in your penis, allowing blood to flow into your penis when you get sexually excited. Sildenafil will only help you to get an erection if you are sexually stimulated.

Sildenafil is a treatment for men with **erectile dysfunction**, sometimes known as impotence. This is when a man cannot get, or keep a hard, erect penis suitable for sexual activity.

2. What you need to know before you take Sildenafil

Do not take Sildenafil:

- If you are **allergic** to sildenafil or any of the other ingredients of this medicine (listed in section 6).
- If you are taking medicines called **nitrates**, as the combination may cause a potentially dangerous decrease in your blood pressure. Tell your doctor if you are taking any of these medicines which are often given for relief of angina pectoris (or "chest pain"). **If you are not certain, ask your doctor or pharmacist.**
- If you are using any of the drugs known as **nitric oxide donors** such as amyl nitrite ("poppers") as the combination may also lead to a potentially dangerous decrease in your blood pressure.
- If you are taking **riociguat**. This drug is used to treat pulmonary arterial hypertension (i.e., high blood pressure in the lungs) and chronic thromboembolic pulmonary hypertension (i.e., high blood pressure in the lungs secondary to blood clots). PDE5 inhibitors, such as sildenafil have been shown to increase the hypotensive effects of this medicine. If you are taking riociguat or are unsure tell your doctor.
- If you have a **severe heart or liver** problem.
- If you have recently had a **stroke** or a **heart attack**, or if you have **low blood pressure**.
- If you have certain rare inherited eye diseases (such as **retinitis pigmentosa**).
- If you have ever had loss of vision because of **non-arteritic anterior ischaemic optic neuropathy** (NAION).

Warnings and precautions

Tell your doctor before taking Sildenafil:

- If you have **sickle cell anaemia** (an abnormality of red blood cells), leukaemia (cancer of blood cells), multiple myeloma (cancer of bone marrow)
- If you have a deformity of your penis or **Peyronie's Disease**.
- If you have problems with your **heart**. Your doctor should in that case carefully check whether your heart can take the additional strain of having sex.
- If you currently have a **stomach ulcer**, or a bleeding problem (such as **haemophilia**).
- If you experience sudden decrease or loss of **vision**, stop taking Sildenafil and contact your doctor immediately.

You should not use Sildenafil with any other oral or local treatments for erectile dysfunction.

You should not use Sildenafil with treatments for pulmonary arterial hypertension (PAH) containing sildenafil or any other PDE5 inhibitors.

You should not take Sildenafil if you do not have erectile dysfunction.

You should not take Sildenafil if you are a woman.

Children and adolescents

Sildenafil should not be given to individuals under the age of 18.

Patients with kidney or liver problems

You should tell your doctor if you have **kidney** or **liver** problems. Your doctor may decide on a lower dose for you.

Other medicines and Sildenafil

Tell your doctor or pharmacist if you are taking, have recently taken other medicines or might take any other medicines, including medicines obtained without prescription.

Sildenafil tablets may interfere with some medicines, especially those used to treat **chest pain**. In the event of a medical emergency, you should tell any healthcare professional treating your condition that you have taken Sildenafil and when you did. Do not take Sildenafil with other medicines unless your doctor tells you that you can.

You should not take Sildenafil if you are taking medicines called **nitrates** as the combination of these products may cause a potentially dangerous decrease in your blood pressure. Always tell your doctor or pharmacist if you are taking any of these medicines that are often used for the relief of angina pectoris (or "chest pain").

You should not take Sildenafil if you are using any of the medicines known as **nitric oxide donors** such as amyl nitrite ("poppers") as the combination may also lead to a potentially dangerous decrease in your blood pressure.

If you are taking medicines known as protease inhibitors, such as for the **treatment of HIV**, your doctor may start you on the lowest dose (25 mg) of Sildenafil.

Some patients who take **alpha-blocker** therapy for the treatment of high blood pressure or prostate enlargement may experience dizziness or light-headedness which may be caused by low blood pressure upon sitting or standing up quickly. Certain patients have experienced these symptoms when taking Sildenafil with alpha-blockers. This is most likely to occur within 4 hours after taking Sildenafil. In order to reduce the likelihood that these symptoms occur, you should be on a regular daily dose of your alpha-blocker before you start Sildenafil. Your doctor may start you on a lower dose (25 mg) of Sildenafil.

Sildenafil with food and drink

Sildenafil can be taken with or without food. However, you may find that Sildenafil takes longer to start working if you take it with a heavy meal.

Drinking alcohol can temporarily impair your ability to get an erection. To get the maximum benefit from your medicine, you are advised not to drink excessive amounts of alcohol before taking Sildenafil.

Pregnancy and Breast-feeding

Sildenafil is not indicated for use by women.

Driving and using machines

Sildenafil can cause **dizziness** and can affect **vision**. You should be aware of how you react to Sildenafil before you drive or use machinery.

3. How to take Sildenafil

Always take Sildenafil exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure. The recommended starting dose is 50 mg.

You should not take Sildenafil more than once a day.

You should take Sildenafil about **one hour** before you plan to have sex. Swallow the tablet whole with a glass of water. The 50mg and 100mg tablets can be divided into equal doses.

If you have the impression that the effect of Sildenafil is too strong or too weak, talk to your doctor or pharmacist.

Sildenafil will only help you to get an erection if you are **sexually stimulated**. The amount of time Sildenafil takes to work varies from person to person, but it normally takes between half an hour and one hour. You may find that Sildenafil takes longer to work if you take it with a heavy meal.

If Sildenafil does not help you to get an erection, or if your erection does not last long enough for you to complete sexual intercourse you should tell your doctor.

If you take more Sildenafil than you should

You may experience an increase in side effects and their severity. Doses above 100 mg do not increase the efficacy.

You should not take more tablets than your doctor tells you to.

Contact your doctor if you take more tablets than you should.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects although not everybody gets them. The side effects reported in association with the use of Sildenafil are usually mild to moderate and of a short duration.

If you experience any of the following serious side effects stop taking Sildenafil and seek medical help immediately:

- An allergic reaction - this occurs **uncommonly** (may affect up to 1 in 100 people). Symptoms include sudden wheeziness, difficulty in breathing or dizziness, swelling of the eyelids, face, lips or throat.
- Chest pains - this occurs **uncommonly**
If this occurs during or after intercourse:
 - Get in a semi-sitting position and try to relax.
 - **Do not use nitrates** to treat your chest pain.
- Prolonged and sometimes painful erections – this occurs **rarely** (may affect up to 1 in 1,000 people). If you have an erection which lasts for more than 4 hours, you should contact a doctor immediately.
- A sudden decrease or loss of vision - this occurs **rarely**
- Serious skin reactions - this occurs **rarely**. Symptoms may include severe peeling and swelling of the skin, blistering of the mouth, genitals and around the eyes, fever.
- Seizures or fits - this occurs **rarely**

Other side effects:

Very common (may affect more than 1 in 10 people): headache.

Common (may affect up to 1 in 10 people): nausea, facial flushing, hot flush (symptoms include a sudden feeling of heat in your upper body), indigestion, colour tinge to vision, blurred vision, visual disturbance, stuffy nose and dizziness.

Uncommon (may affect up to 1 in 100 people): vomiting, skin rash, eye irritation, bloodshot eyes/red eyes, eye pain, seeing flashes of light, visual brightness, light sensitivity, watery eyes, pounding heartbeat, rapid heartbeat, high blood pressure, low blood pressure, muscle pain, feeling sleepy, reduced sense of touch, vertigo, ringing in the ears, dry mouth, blocked or stuffy sinuses, inflammation of the lining of the nose (symptoms include runny nose, sneezing and stuffy nose), upper

abdominal pain, gastro-oesophageal reflux disease (symptoms include heartburn), presence of blood in urine, pain in the arms or legs, nosebleed, feeling hot and feeling tired.

Rare (may affect up to 1 in 1,000 people): fainting, stroke, heart attack, irregular heartbeat, temporary decreased blood flow to parts of the brain, feeling of tightening of the throat, numb mouth, bleeding at the back of the eye, double vision, reduced sharpness of vision, abnormal sensation in the eye, swelling of the eye or eyelid, small particles or spots in your vision, seeing halos around lights, dilation of the pupil of the eye, discolouration of the white of the eye, penile bleeding, presence of blood in semen, dry nose, swelling of the inside of the nose, feeling irritable and sudden decrease or loss of hearing.

From post-marketing experience cases of unstable angina (a heart condition) and sudden death have been reported rarely. Of note most but not all, of the men who experienced these side effects had heart problems before taking this medicine. It is not possible to determine whether these events were directly related to Sildenafil.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse: This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the **Yellow Card Scheme** at www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Sildenafil

Keep this medicine out of the sight and reach of children. Store in the outer carton in order to protect from light. Do not use Sildenafil after the expiry date which is stated on the carton. The expiry date refers to the last day of the month. Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Sildenafil contains

- **Sildenafil 25 mg:** Each tablet contains 35.12 mg of sildenafil citrate quantitatively equivalent to 25 mg sildenafil.
- **Sildenafil 50 mg:** Each tablet contains 70.24 mg of sildenafil citrate quantitatively equivalent to 50 mg of sildenafil.
- **Sildenafil 100 mg:** Each tablet contains 140.48 mg of sildenafil citrate quantitatively equivalent to 100 mg of sildenafil.

The other ingredients are:

- Tablet core: microcrystalline cellulose, calcium hydrogen phosphate (anhydrous), povidone 25, croscarmellose sodium, magnesium stearate.
- Film-coat: Hypromellose, microcrystalline cellulose, titanium dioxide (E171), brilliant blue FCF aluminium lake (E133), macrogol stearate, macrogol 6000.

What Sildenafil looks like and contents of the pack

- **Sildenafil 25 mg** film-coated tablets are blue, oval and biconvex.
- **Sildenafil 50 mg** film-coated tablets are blue, oval and biconvex with break-marks on both sides to facilitate ease of dosing. The tablet can be divided into equal doses.
- **Sildenafil 100 mg** film-coated tablets are blue, oval and biconvex with break-marks on both sides to facilitate ease of dosing. The tablet can be divided into equal doses.

The tablets are provided in blister packs containing 4, 10, 12 or 20 film-coated tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

hameln pharmaceuticals ltd
Nexus, Gloucester Business Park
Gloucester, GL3 4AG, United Kingdom

Manufacturer

hameln rds a.s.
Horná 36, 900 01 Modra, Slovak Republic

This leaflet was last revised in February 2016.