



Package leaflet: Information for the patient

Movicol® Paediatric Plain 6.9g powder for oral solution

Macrogol 3350, Sodium Chloride, Sodium Hydrogen Carbonate, Potassium Chloride

Read all of this leaflet carefully before you start to give this medicine to your child because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for your child. Do not pass it on to others. It may harm them, even if their signs of illness are the same as your child's.
- If your child gets any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Movicol Paediatric Plain is and what it is used for
2. What you need to know before you give Movicol Paediatric Plain
3. How to give Movicol Paediatric Plain
4. Possible side effects
5. How to store Movicol Paediatric Plain
6. Contents of the pack and other information

1. What Movicol Paediatric Plain is and what it is used for

The name of this medicine is Movicol Paediatric Plain 6.9g powder for oral solution. It is a laxative for the treatment of chronic constipation in children aged 1 to 11 years and for the treatment of very bad constipation (called faecal impaction) in children aged 5 to 11 years.

Movicol Paediatric Plain helps your child to have a comfortable bowel movement even if they have been constipated for a long time.

2. What you need to know before you give Movicol Paediatric Plain

Do not give Movicol Paediatric Plain if your doctor has told you that your child has:

- a blockage in their intestine (gut obstruction, ileus)
- a perforated gut wall
- severe inflammatory bowel disease like ulcerative colitis, Crohn's disease or toxic megacolon
- an allergy to the active substances of Movicol Paediatric Plain (listed in section 6).

If your child has a heart or kidney problem, an impaired gag reflex or reflux oesophagitis (heart-burn caused by back-flow of stomach contents into the oesophagus), you should talk to your doctor before giving Movicol Paediatric Plain.

**Warnings and precautions**

When taking Movicol Paediatric Plain you should continue to take plenty of fluids. The fluid content of Movicol should not replace your regular liquid intake.

Other medicines and Movicol Paediatric Plain

Tell your doctor if your child is taking or has recently taken or might take any other medicines.

Some medicines, e.g. anti-epileptics, may not work as effectively during use with Movicol Paediatric Plain. When taking large volumes of Movicol Paediatric Plain (e.g. for faecal impaction), other medicines may be flushed through your digestive system. Your child should not take other medicines by mouth together with Movicol Paediatric Plain and for one hour before and for one hour after taking Movicol Paediatric Plain.

If you need to thicken fluids in order to swallow them safely, Movicol Paediatric Plain may counteract the effect of the thickener.

Pregnancy and breast feeding

Movicol Paediatric Plain can be given during pregnancy and whilst breastfeeding. Ask your pharmacist or doctor for advice before giving Movicol Paediatric Plain.

Movicol Paediatric Plain contains sodium

Movicol Paediatric Plain contains 93.4 mg sodium (main component of cooking/table salt) per sachet. This is equivalent to 4.6% of the recommended maximum daily dietary intake of sodium for an adult.

3. How to give Movicol Paediatric Plain

This medicine can be given at any time with or without food.

Dissolve the content of each sachet in 62.5 ml (1/4 of a glass) of water and drink.

Chronic Constipation:

The dose of Movicol Paediatric Plain depends on the age of the child and their response to treatment.

- As a starting dose, children aged 1 to 6 years should be given 1 sachet (=1 dose) each day.
- Children aged 7 to 11 should be given 2 sachets each day.

Your doctor may tell you to increase the number of sachets taken until your child has a soft bowel movement. If the dose does need increasing, this should be done every other day. For children below 2 years of age, no more than 2 sachets should be taken in one day. For children 2 to 11 years, normally, no more than 4 sachets are required to be taken in one day.

It is not necessary to take all of the drink at one time, if the child prefers it, half the drink can be taken in the morning and half in the evening.

Faecal impaction:



Before your child takes Movicol Paediatric Plain for faecal impaction, it should be confirmed medically that your child has this condition.

A course of treatment with Movicol Paediatric Plain is as follows:

	Number of Movicol Paediatric Plain sachets						
Age (years)	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
5-11	4	6	8	10	12	12	12

The daily number of sachets should be taken in divided doses, all taken within 12 hours (e.g. 8am – 8pm). Each sachet dissolved in 62.5 ml (1/4 of a glass) of water. Treatment can be stopped when the medicine has worked. This is shown by your child passing large volumes of stool and/or watery diarrhoea.

How to mix

- Open the sachet and pour the contents into a glass. Add about 62.5 ml or ¼ glass of water to the glass. Stir well until all the powder has dissolved and the Movicol Paediatric Plain solution is clear or slightly hazy, then give it to your child to drink.
- If you like you can add a flavour such as orange squash to the drink

Duration of treatment

Chronic constipation:

Treatment needs to be for a prolonged period of time, at least 6 – 12 months.

Faecal impaction:

Treatment with Movicol Paediatric Plain can be for up to 7 days.

If your child takes more Movicol Paediatric Plain than they should:

Your child may develop excessive diarrhoea. If this occurs, stop taking Movicol Paediatric Plain and drink plenty of fluids. If you are worried contact your doctor or pharmacist.

If you forget to give Movicol Paediatric Plain:

Give the dose as soon as you remember.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Tell your doctor immediately and stop giving Movicol Paediatric Plain if your child:

- gets a serious allergic reaction which causes difficulty in breathing or swelling of the face, lips, tongue or throat.

Very common side effects (more than 1 person in 10)

- stomach pains
- stomach rumbles



Common side effects (more than 1 in 100 but less than 1 in 10 people):

- diarrhoea
- vomiting (being sick)
- nausea (feeling sick)
- anal discomfort (uncomfortable sensation around the bottom).

If your child is being treated for **chronic constipation**, diarrhoea will usually improve when the dose is reduced.

Uncommon side effects (more than 1 in 1000 but less than 1 in 100 people)

- swollen stomach
- wind

Rare side effects (less than 1 person in 1000)

- allergic reactions, which may include, breathlessness or difficulty in breathing.

Other side effects reported include:

Skin rash (hives), itching, reddening of the skin or a nettle rash, swollen hands, feet or ankles, headaches, indigestion, high or low levels of potassium in the blood

Reporting of side effects

If your child gets any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Movicol Paediatric Plain

Keep this medicine out of the sight and reach of children.

- Do not use Movicol Paediatric Plain after the expiry date which is stated on the sachet and carton.
- The expiry date refers to the last day of that month.
- This medicinal product does not require any special storage conditions.
- Once you have made up Movicol Paediatric Plain in water, if your child cannot drink it straight away keep it covered and in the fridge (2°C – 8°C). Throw away any solution not used within a 6-hour period.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.



6. Contents of the pack and other information

What Movicol Paediatric Plain contains

Each sachet of Movicol Paediatric Plain contains the following:

Macrogol 3350	6.563 g
Sodium Chloride	0.1754 g
Sodium Hydrogen Carbonate	0.0893 g
Potassium Chloride	0.0251 g

Movicol Paediatric Plain contains no colours, flavours or sweeteners.

When it is made into a drink with $\frac{1}{4}$ glass (62.5ml) of water, each sachet gives the equivalent of:

Sodium	65 millimoles/litre
Chloride	53 millimoles/litre
Potassium	5.4 millimoles/litre
Bicarbonate	17 millimoles/litre

What Movicol Paediatric Plain looks like and contents of the pack

Movicol Paediatric Plain is available in rectangular sachets and tubular (stick-pack) sachets.

Movicol Paediatric Plain is a white powder.

Movicol Paediatric Plain is available in boxes of 6, 8, 10, 20, 30, 40, 50, 60 or 100 sachets. Not all sachets and pack sizes may be marketed.

Marketing authorisation holder:

Norgine Pharmaceuticals Limited,
ARC Uxbridge, Building 01,
Sanderson Road, Uxbridge, UB8 1DH, UK

Manufacturer:

Norgine Limited, New Road, Hengoed, Mid Glamorgan, CF82 8SJ, U.K.

Or

Norgine B.V., Antonio Vivaldistraat 150, 1083 HP Amsterdam, The Netherlands.

Or

Meribel Pharma Höganäs AB, Sporthallsvägen 6, 263 35 Höganäs, Sweden.

This leaflet was last revised in 12/2025.

Other sources of information

If you need the information on this leaflet in an alternative format, such as large print, or Braille please ring 0800 198 5000.