



## Package leaflet: Information for the user

### **Irbesartan and Hydrochlorothiazide 150 mg/ 12.5 mg film-coated tablets** **Irbesartan and Hydrochlorothiazide 300 mg/ 12.5 mg film-coated tablets** **Irbesartan and Hydrochlorothiazide 300 mg/ 25 mg film-coated tablets** irbesartan/ hydrochlorothiazide

#### **Read all of this leaflet carefully before you start taking this medicine, because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Irbesartan and Hydrochlorothiazide is and what it is used for
2. What you need to know before you take Irbesartan and Hydrochlorothiazide
3. How to take Irbesartan and Hydrochlorothiazide
4. Possible side effects
5. How to store Irbesartan and Hydrochlorothiazide
6. Contents of the pack and other information

#### **1. What Irbesartan/ Hydrochlorothiazide is and what it is used for**

Irbesartan and Hydrochlorothiazide is a combination of two active substances, irbesartan and hydrochlorothiazide.

Irbesartan belongs to a group of medicines known as angiotensin-II receptor antagonists. Angiotensin-II is a substance produced in the body that binds to receptors in blood vessels causing them to tighten. This results in an increase in blood pressure. Irbesartan prevents the binding of angiotensin-II to these receptors, causing the blood vessels to relax and the blood pressure to lower. Hydrochlorothiazide is one of a group of medicines (called thiazide diuretics) that causes increased urine output and so causes a lowering of blood pressure.

The two active ingredients in Irbesartan and Hydrochlorothiazide work together to lower blood pressure further than if either was given alone.

**Irbesartan and Hydrochlorothiazide is used to treat high blood pressure**, when treatment with irbesartan or hydrochlorothiazide alone did not provide adequate control of your blood pressure.

#### **2. What you need to know before you take Irbesartan/ Hydrochlorothiazide**

##### **Do not take Irbesartan and Hydrochlorothiazide**

- if you are **allergic** to irbesartan or any of the other ingredients of this medicine (listed in section 6)
- if you are **allergic** to hydrochlorothiazide or any other sulfonamide-derived medicines
- if you are **more than 3 months pregnant**. (It is also better to avoid Irbesartan and Hydrochlorothiazide in early pregnancy – see pregnancy section)
- if you have **severe liver or kidney problems**
- if you have **difficulty in producing urine**
- if your doctor determines that you have **persistently high calcium or low potassium levels in your blood**
- if you have diabetes or **impaired kidney function** and you are treated with a blood pressure lowering medicine containing aliskiren

##### **Warnings and precautions**

Talk to your doctor before taking Irbesartan/ Hydrochlorothiazide and if any of the following apply to you:

- if you have had skin cancer or if you develop an unexpected skin lesion during the treatment. Treatment with hydrochlorothiazide, particularly long term use with high doses, may increase the risk of some types of skin and lip cancer (non-melanoma skin cancer). Protect your skin from sun exposure and UV rays while taking Irbesartan/ Hydrochlorothiazide
- if you get **excessive vomiting or diarrhoea**
- if you suffer from **kidney problems** or have a **kidney transplant**
- if you suffer from **heart problems**
- if you suffer from **liver problems**
- if you suffer from **diabetes**
- if you develop low blood sugar levels (symptoms may include sweating, weakness, hunger, dizziness, trembling, headache, flushing or paleness, numbness, having a fast, pounding heart beat), particularly if you are being treated for diabetes.
- if you suffer from **lupus erythematosus** (also known as lupus or SLE)
- if you suffer from **primary aldosteronism** (a condition related to high production of the hormone aldosterone, which causes sodium retention and, in turn, an increase in blood pressure).
- if you are taking any of the following medicines used to treat high blood pressure:
  - an ACE-inhibitor (for example enalapril, lisinopril, ramipril), in particular if you have diabetes-related kidney problems.
  - aliskiren
- If you have decrease in your vision or pain in one of the eyes while taking Irbesartan/ Hydrochlorothiazide. These could be symptoms of fluid accumulation in the vascular layer of the eye (choroidal effusion) or an increase of pressure in your eye (glaucoma) and can happen within hours to weeks of taking Irbesartan/ Hydrochlorothiazide. This can lead to permanent vision loss, if not treated. If you earlier have had a penicillin or sulfonamide allergy, you can be at higher risk of developing this. You should discontinue Irbesartan/ Hydrochlorothiazide treatment and seek prompt medical attention.

Your doctor may check your kidney function, blood pressure, and the amount of electrolytes (e.g. potassium) in your blood at regular intervals.

See also information under the heading “Do not take Irbesartan/ Hydrochlorothiazide”

You must tell your doctor if you think you are (or might become) pregnant. Irbesartan and

Hydrochlorothiazide is not recommended in early pregnancy, and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see pregnancy section).

##### **You should also tell your doctor:**

- if you are on a **low-salt diet**
- if you have signs such as **abnormal thirst, dry mouth, general weakness, drowsiness, muscle pain or cramps, nausea, vomiting**, or an **abnormally fast heart beat** which may indicate an excessive effect of hydrochlorothiazide (contained in Irbesartan and Hydrochlorothiazide)
- if you experience an increased **sensitivity of the skin to the sun** with symptoms of sunburn (such as redness, itching, swelling, blistering) occurring more quickly than normal
- if you are **going to have an operation** (surgery) or **be given anaesthetics**

The hydrochlorothiazide contained in this medicine could produce a positive result in an anti-doping test.

##### **Children and adolescents**

Irbesartan/ Hydrochlorothiazide should not be given to children and adolescents (under 18 years).

##### **Other medicines and Irbesartan/ Hydrochlorothiazide**

Tell your doctor or pharmacist if you are taking or have recently or might take any other medicines.

Diuretic agents such as the hydrochlorothiazide contained in Irbesartan and Hydrochlorothiazide may have an effect on other medicines. Preparations containing lithium should not be taken with Irbesartan and Hydrochlorothiazide without close supervision by your doctor.

Your doctor may need to change your dose and/or to take other precautions:

If you are taking an ACE-inhibitor or aliskiren (see also information under the headings “Do not take Irbesartan/ Hydrochlorothiazide” and “Warnings and precautions”)

##### **You may need to have blood checks if you take:**

- potassium supplements
- salt substitutes containing potassium
- potassium-sparing medicines or other diuretics (water tablets)
- some laxatives
- medicines for the treatment of gout
- therapeutic vitamin D supplements
- medicines to control heart rhythm
- medicines for diabetes (oral agents as repaglinide or insulins)
- carbamazepine (a medicine for the treatment of epilepsy)

It is also important to tell your doctor if you are taking other medicines to reduce your blood pressure, steroids, medicines to treat cancer, pain killers, arthritis medicines, or colestyramine and colestipol resins for lowering blood cholesterol.

##### **Irbesartan and Hydrochlorothiazide with food and drink and alcohol**

Irbesartan and Hydrochlorothiazide can be taken with or without food.

Due to the hydrochlorothiazide contained in Irbesartan and Hydrochlorothiazide, if you drink alcohol while on treatment with this medicine, you may have an increased feeling of dizziness on standing up, specially when getting up from a sitting position.

##### **Pregnancy, breast-feeding and fertility**

###### **Pregnancy**

You must tell your doctor if you think you are (or might become) pregnant. Your doctor will normally advise you to stop taking Irbesartan and Hydrochlorothiazide before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Irbesartan and Hydrochlorothiazide. Irbesartan and Hydrochlorothiazide is not recommended during pregnancy, and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

###### **Breast-feeding**

Tell your doctor if you are breast-feeding or about to start breast-feeding. Irbesartan and Hydrochlorothiazide is not recommended for mothers who are breast-feeding, and your doctor may choose another treatment for you if you wish to breast-feed, especially if your baby is newborn, or was born prematurely.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

###### **Driving and using machines**

Irbesartan and Hydrochlorothiazide is unlikely to affect your ability to drive or use machines. However, occasionally dizziness or weariness may occur during treatment of high blood pressure. If you experience these, talk to your doctor before attempting to drive or use machines.

###### **Irbesartan and Hydrochlorothiazide contains lactose**

Irbesartan/ Hydrochlorothiazide contains lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

###### **Irbesartan contain Sodium:**

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially ‘sodium-free’.

#### **3. How to take Irbesartan/ Hydrochlorothiazide**

Always take this medicine exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

##### **Dosage**

The recommended dose of Irbesartan and Hydrochlorothiazide is one or two tablets a day.

■ Black

Version: 11

Date & Time: 29.04.2021 & 5.40 pm

Submission Code: N07323

P15XXXX



Irbesartan and Hydrochlorothiazide will usually be prescribed by your doctor when your previous treatment did not reduce your blood pressure enough. Your doctor will instruct you how to switch from the previous treatment to Irbesartan and Hydrochlorothiazide.

#### Method of administration

Irbesartan and Hydrochlorothiazide is for **oral use**. Swallow the tablets with a sufficient amount of fluid (e.g. one glass of water). You can take Irbesartan and Hydrochlorothiazide with or without food. Try to take your daily dose at about the same time each day. It is important that you continue to take Irbesartan and Hydrochlorothiazide until your doctor tells you otherwise.

The maximal blood pressure lowering effect should be reached 6-8 weeks after beginning treatment.

#### If you take more Irbesartan and Hydrochlorothiazide than you should

If you accidentally take too many tablets, contact your doctor immediately.

#### Children should not take Irbesartan and Hydrochlorothiazide

Irbesartan and Hydrochlorothiazide should not be given to children under 18 years of age. If a child swallows some tablets, contact your doctor immediately.

#### If you forget to take Irbesartan and Hydrochlorothiazide

If you accidentally miss a daily dose, just take the next dose as normal. Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Some of these effects may be serious and may require medical attention.

Rare cases of allergic skin reactions (rash, urticaria), as well as localised swelling of the face, lips and/or tongue have been reported in patients taking irbesartan.

**If you get any of the above symptoms or get short of breath**, stop taking Irbesartan and Hydrochlorothiazide and contact your doctor immediately.

The frequency of the side effects listed below is defined using the following convention:

Common: may affect up to 1 in 10 people

Uncommon: may affect up to 1 in 100 people

Side effects reported in clinical studies for patients treated with irbesartan/ hydrochlorothiazide were:

**Common side effects** (may affect up to 1 in 10 people):

- nausea/vomiting
- abnormal urination
- fatigue
- dizziness (including when getting up from a lying or sitting position)
- blood tests may show raised levels of an enzyme that measures the muscle and heart function (creatinine kinase) or raised levels of substances that measure kidney function (blood urea nitrogen, creatinine).

**If any of these side effects causes you problems**, talk to your doctor.

**Uncommon side effects** (may affect up to 1 in 100 people):

- diarrhoea
- low blood pressure
- fainting
- heart rate increased
- flushing
- swelling
- sexual dysfunction (problems with sexual performance)
- blood tests may show lowered levels of potassium and sodium in your blood.

**Not known** (frequency cannot be estimated from the available data):

- Skin and lip cancer (Non-melanoma skin cancer)

**If any of these side effects causes you problems**, talk to your doctor.

#### Side effects reported since the launch of Irbesartan/ Hydrochlorothiazide

Some undesirable effects have been reported since marketing of Irbesartan/ Hydrochlorothiazide. Undesirable effects where the frequency is not known are: headache, ringing in the ears, cough, taste disturbance, indigestion, pain in joints and muscles, liver function abnormal and impaired kidney function, increased level of potassium in your blood and allergic reactions such as rash, hives, swelling of the face, lips, mouth, tongue or throat. Uncommon cases of jaundice (yellowing of the skin and/or whites of the eyes) have also been reported.

As for any combination of two active substances, side effects associated with each individual component cannot be excluded.

#### Side effects associated with irbesartan alone

In addition to the side effects listed above, chest pain, severe allergic reactions (anaphylactic shock), decreased number of red blood cells (anaemia – symptoms may include tiredness, headaches, being short of breath when exercising, dizziness and looking pale) and decrease in the number of platelets (a blood cell essential for the clotting of the blood) and low blood sugar levels have also been reported.

#### Side effects associated with hydrochlorothiazide alone

Loss of appetite; stomach irritation; stomach cramps; constipation; jaundice (yellowing of the skin and/or whites of the eyes); inflammation of the pancreas characterised by severe upper stomach pain, often with nausea and vomiting; sleep disorders; depression; blurred vision; decrease in vision or pain in your eyes due to high pressure (possible signs of fluid accumulation in the vascular layer of the eye (choroidal effusion) or acute angle-closure glaucoma); lack of white blood cells, which can result in frequent infections, fever; decrease in the number

of platelets (a blood cell essential for the clotting of the blood), decreased number of red blood cells (anaemia) characterised by tiredness, headaches, being short of breath when exercising, dizziness and looking pale; kidney disease; lung problems including pneumonia or build-up of fluid in the lungs; increased sensitivity of the skin to the sun; inflammation of blood vessels; a skin disease characterized by the peeling of the skin all over the body; cutaneous lupus erythematosus, which is identified by a rash that may appear on the face, neck, and scalp; allergic reactions; weakness and muscle spasm; altered heart rate; reduced blood pressure after a change in body position; swelling of the salivary glands; high sugar levels in the blood; sugar in the urine; increases in some kinds of blood fat; high uric acid levels in the blood, which may cause gout.

It is known that side effects associated with hydrochlorothiazide may increase with higher doses of hydrochlorothiazide.

#### Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via Yellow Card Scheme Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard), or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

### 5. How to store Irbesartan/ Hydrochlorothiazide

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and on the blister/label of the bottle after EXP. The expiry date refers to the last day of that month.

Store below 30°C.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

### 6. Contents of the pack and other information

#### What Irbesartan and Hydrochlorothiazide contains

- The active substance is irbesartan & hydrochlorothiazide.  
Each film-coated tablet contains 150 mg of irbesartan and 12.5 mg of hydrochlorothiazide.  
Each film-coated tablet contains 300 mg of irbesartan and 12.5 mg of hydrochlorothiazide.  
Each film-coated tablet contains 300 mg of irbesartan and 25 mg of hydrochlorothiazide.
- The other ingredients are  
*Tablet core:* Lactose monohydrate, Sodium Starch Glycolate (Type A), Povidone K30, Silica Colloidal Anhydrous, Talc, Sodium stearyl fumarate.

#### Tablet coat:

150mg/12.5mg & 300mg/12.5mg film-coated tablets

Lactose monohydrate  
Hypromellose (E464)  
Titanium dioxide (E171)  
Macrogol 4000  
Iron oxide yellow (E172)  
Iron oxide red (E172)

#### 300 mg/25 mg film-coated tablets

Hypromellose (E464)  
Titanium dioxide (E171)  
Macrogol 4000  
Iron oxide red (E172)  
Iron oxide black (E172)

#### What Irbesartan and Hydrochlorothiazide looks like and contents of the pack

Film-coated Tablet.

#### 150 mg/ 12.5 mg

Peach coloured film-coated biconvex oval shaped tablets, debossed with "H 35" on one side and plain on other side.

#### 300 mg/ 12.5 mg

Peach coloured film-coated biconvex oval shaped tablets, debossed with "H 36" on one side and plain on other side.

#### 300 mg/ 25 mg

Pink coloured film-coated biconvex oval shaped tablets, debossed with "H 37" on one side and plain on other side.

Tablets are available in Polyamide/Aluminium/PVC/ Aluminium blister pack and white opaque HDPE bottle with white opaque polypropylene closure pack.

#### Pack size:

Blister pack: 1, 14, 28, 30, 56, 90, 98, 100 & 500 tablets

Bottle pack: 30 & 90 tablets

Not all pack sizes may be marketed.

#### Marketing Authorisation Holder

Milpharm Limited  
Ares Block  
Odyssey Business Park  
West End Road  
Ruislip HA4 6QD  
United Kingdom

#### Manufacturer

APL Swift Services (Malta) Limited  
HF26, Hal Far Industrial Estate, Hal Far  
Birzebbugia, BBG 3000  
Malta

or

Milpharm Limited  
Ares Block  
Odyssey Business Park  
West End Road  
Ruislip HA4 6QD  
United Kingdom

**This leaflet was last revised in 04/2021.**