

 <p>Trihexyphenidyl Tablets 5mg</p> <p>Each tablet contains 5mg trihexyphenidyl hydrochloride</p> <p>84 tablets</p> <p>MA holder: Genus Pharmaceuticals, Linthwaite, Huddersfield, HD7 5QH, UK. PM61315 92793621807</p> <p>POM PL 17225/0002 *Trademark</p>	<p>PACKAGE LEAFLET: INFORMATION FOR THE USER</p> <p>Trihexyphenidyl Tablets 5mg</p> <p>Read all of this leaflet carefully before you start taking this medicine</p> <ul style="list-style-type: none"> - Keep this leaflet. You may need to read it again. - If you have any further questions, ask your doctor or pharmacist. - This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours. - If any of the side-effects get serious, or if you notice any side-effects not listed in this leaflet, please tell your doctor or pharmacist. <p>In this leaflet</p> <ol style="list-style-type: none"> 1. What Trihexyphenidyl Tablets are and what they are used for 2. Before you take Trihexyphenidyl Tablets 3. How to take Trihexyphenidyl Tablets 4. Possible side-effects 5. How to store Trihexyphenidyl Tablets 6. Further information 	<p>1. WHAT TRIHEXYPHENIDYL TABLETS ARE AND WHAT THEY ARE USED FOR</p> <p>The name of your medicine is Trihexyphenidyl Tablets. Trihexyphenidyl is an antispasmodic (muscle relaxing) drug. Trihexyphenidyl is used to treat Parkinson's disease. Trihexyphenidyl helps patients with Parkinson's disease by improving muscle control and reducing stiffness so that they can move more naturally. Trihexyphenidyl can also be used to treat movement disorders (extreme restlessness, involuntary movements or muscle spasms) which may be caused by certain drugs.</p> <p>2. BEFORE YOU TAKE TRIHEXYPHENIDYL TABLETS</p> <p>Do not take Trihexyphenidyl Tablets if you:</p> <ul style="list-style-type: none"> • are allergic (hypersensitive) to trihexyphenidyl or any other ingredient in the tablets (see section 6 – Further information for details) <p>Take special care with Trihexyphenidyl Tablets if you:</p> <ul style="list-style-type: none"> • have glaucoma • have heart, liver or kidney disease or suffer from high blood pressure • have prostate problems
--	---	--

<ul style="list-style-type: none"> • have trouble passing urine • have a stomach ulcer or suffer from severe heartburn or acid indigestion or any other digestive disorders • suffer from a condition called myasthenia gravis • suffer from tardive dyskinesia (unusual movements of the face and tongue) • are elderly or under the age of 21. <p>Taking other medicines</p> <p>Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription (like aspirin).</p> <p>Take special care if you are taking any of the following:</p> <ul style="list-style-type: none"> • monoamine oxidase inhibitors (e.g. moclobemide, tranylcypromine) or tricyclic antidepressants (e.g. amitriptyline) for the treatment of depression • phenothiazines (e.g. chlorpromazine, thioridazine) or clozapine for the treatment of mental illness • antihistamines for the treatment of allergies 	<ul style="list-style-type: none"> • disopyramide for the treatment of an irregular heartbeat • levodopa for the treatment of Parkinson's disease • amantadine for viral infections • nefopam, a painkiller • metoclopramide or domperidone • parasympathomimetics (e.g. neostigmine, bethanechol) for the treatment of urinary retention. <p>Pregnancy and breast-feeding</p> <p>Do not take this medicine if you are pregnant, planning to become pregnant, or breast-feeding without consulting your doctor first.</p> <p>Driving and using machines</p> <p>Trihexyphenidyl can cause blurred vision or dizziness. Do not drive or operate machinery if this happens to you.</p>	<p>Important information about the ingredients of Trihexyphenidyl Tablets</p> <p>This medicine contains starch. This is suitable for people with coeliac disease. Patients with wheat allergy (different from coeliac disease) should not take this medicine.</p> <p>3. HOW TO TAKE TRIHEXYPHENIDYL TABLETS</p> <p>Always take Trihexyphenidyl Tablets exactly as your doctor has told you to. The label on your medicine should also tell you. You should check with your doctor or pharmacist if you are not sure.</p> <p>The dose of trihexyphenidyl varies from person to person depending on the symptoms. You will normally start on a low dose and this may be increased gradually by your doctor. The usual dose is between 5-15mg per day (each tablet contains 5mg). Some people may need less, especially if they are also taking other medicines. Also, if you are over 65 years of age, you may need less than the usual dose.</p> <p>Trihexyphenidyl Tablets should be taken at meal times (just before or just after a meal). If you are not sure how many tablets to take or when to take them, ask your pharmacist. Trihexyphenidyl Tablets should be swallowed whole with a glass of water.</p>
--	---	---

<p>If you take more Trihexyphenidyl Tablets than you should</p> <p>If you have taken an overdose of Trihexyphenidyl Tablets (that is more than the doctor has told you to) get medical help immediately, either by calling your doctor or going to the nearest hospital casualty department. Remember to take the labelled medicine bottle with you, whether there are any Trihexyphenidyl Tablets left or not.</p> <p>If you forget to take Trihexyphenidyl Tablets</p> <p>If you forget to take a tablet you should take it as soon as you remember. However, if this is within 2 hours of your next dose you should skip the missed tablet and carry on taking the rest of your tablets as usual.</p> <p>Do not take a double dose to make up for the forgotten tablet.</p> <p>If you stop taking Trihexyphenidyl Tablets</p> <p>Do not stop taking this medicine suddenly as your symptoms may get worse. If your dose needs to be lowered, your doctor will reduce the amount you should take gradually.</p>	<p>4. POSSIBLE SIDE-EFFECTS</p> <p>Like all medicines, Trihexyphenidyl Tablets can cause side-effects, although not everybody gets them. If you experience any of the following very rare side-effects you should tell your doctor or go to your nearest casualty department immediately:</p> <ul style="list-style-type: none"> • allergic reaction such as swelling of the eyelids, face or lips or skin rash • difficulty breathing or tightness in the chest • difficulty swallowing <p>If you experience any of the following rare side-effects you should tell your doctor as soon as possible:</p> <ul style="list-style-type: none"> • restlessness or agitation • feelings of confusion or delusion (loss of the sense of reality) • hallucinations • difficulty in sleeping • false sense of well-being • immediate or short-term memory loss • worsening of myasthenia gravis (very tired or weak muscles) 	<ul style="list-style-type: none"> • large pupils and lack of near vision • eye discomfort and increased sensitivity to light • painful increased pressure in the eye • faster heartbeat • being sick • reddening and dryness of skin or skin rashes • inability to pass urine or difficulty passing urine • thirst • fever (raised body temperature) <p>However you should tell your doctor if any of the following common side-effects are severe or become troublesome:</p> <ul style="list-style-type: none"> • dryness of mouth (this can be improved by sucking pastilles) • constipation • blurring of vision or dizziness • feeling sick or nervous <p>If any of the side-effects get serious, or if you notice any side-effects not listed in this leaflet, please tell your doctor or pharmacist.</p>
---	--	--

<p>Reporting of side effects</p> <p>If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for 'MHRA Yellow Card' in the Google Play or Apple App Store.</p> <p>By reporting side effects you can help provide more information on the safety of this medicine.</p> <p>5. HOW TO STORE TRIHEXYPHENIDYL TABLETS</p> <p>Keep out of the sight and reach of children.</p> <p>Do not take Trihexyphenidyl Tablets after the expiry date which is printed on the pack. The expiry date refers to the last day of the month.</p> <p>Store below 25°C. Keep your tablets in the original bottle until you need to take one. This protects the tablets from moisture which may spoil them. If your doctor decides to stop your medicine you should return any unused tablets to your pharmacist.</p>	<p>Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help protect the environment.</p> <p>6. FURTHER INFORMATION</p> <p>What Trihexyphenidyl Tablets contain</p> <p>The active substance in Trihexyphenidyl Tablets is trihexyphenidyl hydrochloride. Other ingredients are: calcium hydrogen phosphate, starch and starch pregelatinised, magnesium stearate and purified water.</p> <p>What Trihexyphenidyl Tablets look like and the contents of the pack</p> <p>Trihexyphenidyl Tablets 5mg are white tablets with GP4436 on the side. Each bottle contains 84 tablets.</p>	<p>POM PL 17225/0002 Trihexyphenidyl Tablets 5mg</p> <p>Marketing Authorisation holder: Genus Pharmaceuticals, Linthwaite, Huddersfield, HD7 5QH, UK.</p> <p>Manufacturer: Thornton & Ross Limited, Linthwaite, Huddersfield, HD7 5QH, UK.</p> <p>This leaflet was last revised in July 2018</p>
---	---	---