

CONTENTS: The solution contains Hexetidine 0.1g/100 ml. Other ingredients: Polysorbate 80, citric acid, saccharin sodium, levomenthol, eucalyptus oil, ethanol 96%, azorubin (E122) sodium calcium edetate, sodium hydroxide (E524), and purified water.

Oraldene has an antibacterial and antifungal action which relieves mouth and throat infections.

It can also be used before and after dental surgery.

Read and retain carton for full instructions.

Oraldene is a registered trade mark.

PL holder: McNeil Products Ltd,
High Wycombe, HP12 4EG.

McNeil
Products Ltd.

PL 15513/0067

Oraldene®

Hexetidine

Medicated Relief for
Mouth & Throat
Infections

Antibacterial
Mouthwash to relieve
gingivitis mouth ulcers
sore gums sore throat
200 ml e

Do not use if bottle
seal is broken
when purchased.

Keep out of the
sight and reach of children.

DIRECTIONS: Mouthwash
for topical use only. **DO NOT
SWALLOW. SHAKE WELL
BEFORE USE.**

**ADULTS AND CHILDREN 12
YEARS AND OLDER**

Rinse the mouth or gargle for
30 seconds with at least 15 ml
(approx. 1 tablespoonful) of
undiluted Oraldene 2 to 3 times a
day. **CHILDREN UNDER 12 YEARS -
not recommended.**

If symptoms persist discontinue use
and talk to your doctor or dentist.
Store below 30°C.

Use within 6 months of opening.
Keep the bottle in the outer carton
in order to protect from light.



XXXXXX

EXP

LOT

Oraldene®

Hexetidine

Oraldene®

Hexetidine

How to use this medicine

Check the table below to see how much medicine to use.

i Children under 12 years old

Oraldene is not recommended for children under 12 years of age.

i Adults and Children aged 12 years and over

Age	Dose
Adults and children aged 12 years and over	Rinse the mouth or gargle for 30 seconds with at least 15 ml (approx. 1 tablespoonful) of undiluted Oraldene two to three times a day or as otherwise directed.

- Do not swallow.
- Shake well before use.
- If symptoms persist discontinue use and talk to your doctor or dentist.

! If anyone accidentally swallows this product

Contact a doctor or your nearest Accident and Emergency department (Casualty) taking this pack with you.

! If you forget to use the medicine

Use the next dose as soon as you remember unless it is time to use it again. Do not use two quantities in one go.

Medicated Relief for Mouth & Throat Infections

Antibacterial Mouthwash to relieve *gingivitis* *mouth ulcers* *sore gums* *sore throat*

200 ml e

Possible side-effects

Like all medicines, Oraldene can cause side-effects although not everybody gets them. If you experience the following, stop using the medicine and talk to your doctor:

- Swelling of the face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing
- Allergic reactions including itchy rash Irritation, tingling or numbness of the tongue or inside the mouth
- Shortness of breath Cough Dry mouth Feeling sick (nausea) Sickness (vomiting) Salivary gland enlargement
- Taste disturbance or loss of taste Pain, swelling, redness or heat Tongue or tooth discolouration
- Blistering or ulceration

Reporting of side-effects

If you get any side-effects, talk to your doctor, pharmacist or nurse. This includes any possible side-effects not listed in this carton. You can also report side-effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side-effects you can help provide more information on the safety of this medicine.

Storing this medicine

Do not use if bottle seal is broken when purchased. Store below 30°C.

Use within 6 months of opening.

Keep the bottle in the outer carton in order to protect from light.

Keep this medicine out of the sight and reach of children. Do not use this medicine after the expiry date which is stated on the label. The expiry date refers to the last day of that month.

Retain carton for full instructions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

Further information

What's in this medicine?

The active ingredient is: Hexetidine 0.1 g/100 ml

Other ingredients are: Polysorbate 80, citric acid, saccharin sodium, levomenthol, eucalyptus oil, ethanol 96%, azorubin (E122) sodium calcium edetate, sodium hydroxide (E524), and purified water.

What the medicine looks like

Oraldene is a clear red solution available in 200 ml bottles. It contains 0.1 g of the active ingredient in every 100 ml of medicine.

What the medicine is for

The mouthwash contains hexetidine, which is an antibacterial and antifungal and is used to help treat throat and mouth infections, including mouth ulcers, thrush, sore or bleeding gums (called gingivitis), bad breath (halitosis) or sore throat. It can also be used before and after dental surgery to help prevent infections.

Before using this medicine

This medicine is suitable for most people, but a few people should not use it. If you are in any doubt, talk to your doctor or pharmacist.

X Do not use this medicine...

- If you have ever had a bad reaction to Oraldene or any of its ingredients. If this applies to you, get advice from your doctor or pharmacist without using Oraldene.
- If you are pregnant or breast-feeding
 - Ask your doctor or pharmacist for advice before using Oraldene if you are pregnant or breast-feeding.

! Excipients with known effect

- This medicine contains 607.8 mg of alcohol (ethanol) in each 15 ml. The amount in 15 ml of this medicine is equivalent to less than 15 ml beer or 6 ml wine. The small amount of alcohol in this medicine will not have any noticeable effects.
- Azorubin (E122) may cause allergic reactions.
- This medicine contains less than 1 mmol sodium (23 mg) per 15 ml, that is to say essentially 'sodium-free'.

This carton was revised September 2022. Oraldene is a registered trade mark.

© McNeil Products Limited 2022

Product Licence holder: McNeil Products Ltd, High Wycombe, HP12 4EG.

Manufacturer: Delpharm Orléans, 5 avenue de Concy, 45071 Orléans, Cedex 2, France.

PL 15513/0067