

Package Leaflet: Information for the user
STREPSILS Extra Triple Action Blackcurrant Lozenges
Hexylresorcinol

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

This medicine is available without prescription. However, you still need to take Strepils Extra Triple Action Blackcurrant Lozenges carefully to get the best results from it. Strepils Extra Triple Action Blackcurrant Lozenges will be referred to as this medicine throughout this leaflet.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your pharmacist. This includes any possible side effects not listed in the leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse after 3 days.

In this leaflet:

1. What this medicine is and what it is used for
2. What you need to know before you take this medicine
3. How to take this medicine
4. Possible side effects
5. How to store this medicine
6. Contents of the pack and other information

1. What this medicine is and what it is used for

This medicine contains an active ingredient for use in throat medications. It is used for the symptomatic relief of sore throat and its associated pain. The active ingredient has a mild antiseptic effect to kill the bacteria associated with mouth and throat infections and also acts as a local anaesthetic to numb the pain associated with a sore throat. Sucking the lozenge slowly in the mouth helps to lubricate and soothe the discomfort of the painful area in the throat, and also releases the active ingredient to help relieve the soreness and discomfort of mouth and throat infections.

2. What you need to know before you take this medicine

This medicine is recommended for adults, children (over 6 years old) and the elderly. Children under 6 years old should not use this medicine.

Do not take this medicine if you:

- are allergic (hypersensitive) to Hexylresorcinol or any of the other ingredients shown in Section 6.
- are under 6 years old.

Warnings and precautions

Tell your doctor or pharmacist if you:

- are taking or have recently taken any other medicines, including medicines obtained without a prescription.
- suffer or are suffering from any other throat problems

Pregnancy and breast-feeding

It is not recommended to use this medicine if you think you are pregnant, are pregnant or are breastfeeding. Ask your doctor or pharmacist for advice before taking any medicine.

Important information about some of the ingredients in this medicine

- This medicine contains only very low levels of gluten (from wheat starch). It is regarded as 'gluten-free' and is very unlikely to cause problems if you have coeliac disease. One lozenge contains no more than 22.42 micrograms of gluten. If you have wheat allergy (different from coeliac disease) you should not take this medicine.
- This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.
- This medicine contains fragrance with d-Limonene and Linalool. d-Limonene and Linalool may cause allergic reactions.
- This medicine contains carmoisine (E1220) which may cause allergic reactions.
- This medicine contains sucrose (1.408 g per lozenge) and glucose (1.121g per lozenge). This is to be taken into account in patients with diabetes mellitus. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.
- This medicine contains Sulphites – Sulphur Dioxide (E220) present in liquid glucose which may rarely cause severe hypersensitivity reactions and bronchospasm.
- This medicine contains 10.8 mg propylene glycol in each lozenge.



Pharmacode reads top
to bottom on the
front of the leaflet.

Pharmacode reads
bottom to top on the
back of the leaflet.



3. How to take this medicine

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Adults, Children (over 6 years old) and the elderly – One lozenge to be dissolved slowly in the mouth every 2 – 3 hours up to a maximum of 12 lozenges in 24 hours

Remove one lozenge from the foil blister packaging.

Replace the foil blister back into the cardboard carton.

Place lozenge into the mouth allowing it to dissolve slowly.

Dissolve one lozenge slowly in the mouth every 2 – 3 hours.

It is not recommended that children under 6 years use this medicine.

Remember that young children can choke on lozenges.

If symptoms persist for more than 3 days or if anything unusual happens please consult your doctor.

If you take more of this medicine than you should

You may experience stomach discomfort

Do not take any more of this medicine and consult your doctor or pharmacist.

If you forget to take this medicine

Do not take a double dose to make up for a forgotten lozenge. Continue to use this medicine normally.

If you have any further questions on the use of this medicine ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If you experience any hypersensitivity to this medicine i.e. itching, difficulty in breathing, swelling of the face, neck, tongue or throat and rash (severe allergic reactions) stop taking this medicine and consult your doctor or pharmacist.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store this medicine

Keep out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month and the year.

Do not store above 25°C. Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Contents of the pack and other information

What this medicine contains

The active substance is: Hexylresorcinol 2.4mg

The other ingredients are: Liquid sucrose, Liquid glucose (containing wheat starch (containing gluten) and sulphites – sulphur dioxide (E220)), Blackcurrant flavour (containing Propylene glycol, d-Limonene and Linalool), Levomenthol, Carmoisine Edicol (E122), Patent Blue V (E131).

What this medicine looks like and contents of the pack

A round purple lozenge. In pack sizes of 2, 4, 6, 8, 12, 16, 24, 32, 36 and 48 lozenges.

Not all pack sizes may be marketed.

Marketing Authorisation Holder:

Reckitt Benckiser Healthcare (UK) Limited, Slough, SL1 3UH

Manufacturers:

Reckitt Benckiser Healthcare International Limited, Thane road, Nottingham, NG90 2DB, UK.

RB NL Brands B.V., WTC Schiphol Airport, Schiphol Boulevard 207, 1118 BH Schiphol, NL.

Licence Number: PL 00063/0392

Date of preparation: December 2021