

# bonjela cool mint gel

choline salicylate, cetalkonium chloride

## INFORMATION FOR THE USER

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist have told you.

- Keep this leaflet. You may need to read it again.  
- Ask your pharmacist if you need more information or advice.

- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

- You must talk to a doctor if you do not feel better or if you feel worse after 7 days.

What is in this leaflet:

### 1. What bonjela Cool Mint Gel is and what it is used for

### 2. What you need to know before you use bonjela Cool Mint Gel

### 3. How to use bonjela Cool Mint Gel

### 4. Possible side effects

### 5. How to store bonjela Cool Mint Gel

### 6. Contents of the pack and further information

## 1. WHAT BONJELA COOL MINT GEL IS AND WHAT IT IS USED FOR

### bonjela Cool Mint Gel is used to

- relieve the pain and discomfort of common mouth ulcers and cold sores.
- aid healing of sore spots and ulcers due to dentures and braces.

bonjela Cool Mint Gel is a clear, almost colourless gel, which is applied to the inside of the mouth or gums. It contains choline salicylate (a pain killer and anti-inflammatory) which soothes sore areas in the mouth and an antiseptic called cetalkonium chloride, which kills bacteria and helps prevent infection.

### bonjela Cool Mint Gel is intended for adults and children over the age of 16.

## 2. WHAT YOU NEED TO KNOW BEFORE YOU USE BONJELA COOL MINT GEL

**Do not give to children or adolescents under 16 years of age. This is because there is a possible link between salicylates and Reye's syndrome when given to children. Reye's syndrome is a very rare disease which affects the brain and liver and can be fatal.**

### Do not use if you are

- suffering from a stomach ulcer
- allergic to salicylates, Aspirin or other NSAIDs (e.g. ibuprofen), or to any other ingredients in this product (see section 6)

### Warnings and precautions

This product contains salicylates. Talk to your doctor before taking this product with Aspirin or other salicylates.

### Other medicines and bonjela

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. Salicylates may increase the effect of blood thinning medicines (anticoagulants) and can decrease the effect of medicines to treat gout (uricosurics) such as probenecid, benzbromarone.

### Pregnancy and breast-feeding

If you are pregnant, planning to become pregnant, or breast-feeding, speak to your doctor or pharmacist.



0000000

**Important information about some of the ingredients in bonjela Cool Mint Gel**  
bonjela Cool Mint Gel contains a small amount of ethanol (alcohol), less than 100mg per dose.

## 3. HOW TO USE BONJELA COOL MINT GEL

Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist have told you. Check with your doctor or pharmacist if you are not sure.

**For adults and children over the age of 16 years only.**

### - Mouth ulcers and sores

Massage approximately one centimetre (half an inch) of bonjela Cool Mint Gel onto the sore area. This can be repeated after 3 hours.

### - Denture sores

Massage approximately one centimetre of bonjela Cool Mint Gel onto the sore area, and leave at least 30 minutes before reinserting the dentures. Do not apply bonjela Cool Mint Gel directly onto the dentures.

**Do not take more medicine than the label or leaflet tells you as salicylate toxicity may occur as a result.**

If your symptoms do not go away after 7 days of treatment, you should see your doctor or dentist.

**In the case of a missed dose**, simply apply a new dose and follow the instructions described above. **In the event of an overdose**, seek medical advice.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

## 4. POSSIBLE SIDE EFFECTS

Like all medicines, bonjela Cool Mint Gel can cause side effects, although not everybody gets them.

**STOP taking this product and seek immediate medical help if you experience:**

- unexplained wheezing or coughing, asthma, shortness of breath, difficulty breathing, itching, face, tongue or throat swelling, which can be signs of serious allergic reactions.

### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard)  
By reporting side effects you can help provide more information on the safety of this medicine.

## 5. HOW TO STORE BONJELA COOL MINT GEL

Keep out of the sight and reach of children.

Do not use bonjela Cool Mint Gel after the expiry date which is stated on the carton and on the tube. The expiry date refers to the last day of that month. Do not store above 25°C.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment

## 6. CONTENTS OF THE PACK AND FURTHER INFORMATION

### What bonjela Cool Mint Gel contains

The active ingredients (which make the medicine work) are choline salicylate 8.714% w/w, cetalkonium chloride 0.01 % w/w. The other ingredients are ethanol, glycerol, menthol, hypromellose 4500, mint flavour, sodium saccharin and water.

**What bonjela Cool Mint Gel looks like and contents of the pack** Each tube contains 15g of clear, mint flavoured oral gel.

**Marketing Authorisation holder/Manufacturer:** Reckitt Benckiser Healthcare (UK) Limited, Hull HU8 7DS

Licence number: PL00063/0122

Date of last revision January 2018

0000000

Ink &  
Varnish Free