

**BLOODSHOT EYE DROPS**  
*Hamamelis Virginiana*  
**Naphazoline Hydrochloride**  
 Package leaflet: Information for the patient

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.

- Keep this leaflet. You might need to read it again.
- Ask your doctor or pharmacist if you need more information or advice
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to your doctor or pharmacist if you do not feel better or feel worse after 2 days.

1. What Optrex Bloodshot Eye Drops is and what it is used for
2. What you need to know before you use Optrex Bloodshot Eye Drops
3. How to use Optrex Bloodshot Eye Drops
4. Possible side effects
5. How to store Optrex Bloodshot Eye Drops
6. Contents of the pack and other information

**1. What Optrex Bloodshot Eye Drops is and what it is used for**

Dusty, smoky atmospheres, eye strain and chlorine in swimming pools can make the eyes red. When the eyes are red it is because more blood than usual is flowing through the small blood vessels of the eye.

Optrex Bloodshot Eye Drops:

- are mildly astringent to temporarily remove redness of the eye caused by minor eye irritation.
- contain an active ingredient which restores the normal flow of blood to remove redness.

**2. What you need to know before you use Optrex Bloodshot Eye Drops**

Optrex Bloodshot Eye Drops are recommended for **adults and children over 12 years**

**Do not use this product if:**

- you are sensitive to Hamamelis water, naphazoline, or to any of the ingredients listed in Section 6.
- you have **glaucoma, other eye disease** or have had **eye surgery**.

You have taken monoamine oxidase inhibitors (MAOIs) in the last 14 days (medicine used to treat conditions such as depression).

**Warnings and precautions**  
 Talk to your doctor or pharmacist before using this medicine if you have:

- high blood pressure
- depression
- heart disease
- diabetes
- thyroid disorders
- hardening and narrowing of blood vessels

Or experience eye pain or changes in your eyesight

**Taking other medicines**  
 If you are taking any other medicines you should speak to your doctor or pharmacist before using this product, particularly if you:

- are taking **digoxin, quinidine or tricyclic antidepressants**
- have taken **monoamine oxidase inhibitors (MAOIs)** within the last 14 days
- are taking antihypertensive (high blood pressure) medication
- are using any other eye treatments

**Pregnancy and Breastfeeding**  
 If you are pregnant, planning to become pregnant or breastfeeding speak to your doctor or pharmacist **before using this product.**

**Important information about some of the ingredients in Optrex Bloodshot Eye Drops**  
 This medicine contains 0.0025 mg benzalkonium chloride in each drop. Benzalkonium chloride may be absorbed by soft contact lenses and may change the colour of the contact lenses. You should remove contact lenses before using this medicine and put them back 15 minutes afterwards.

Benzalkonium chloride may also cause eye irritation, especially if you have dry eyes or disorders of the cornea (the clear layer at the front of the eye). If you feel abnormal eye sensation, stinging or pain in the eye after using this medicine, talk to your doctor.

**Driving and using machines**  
 Do not drive or operate machinery if the vision is blurred.

Please turn over



Pharmacode reads this way



Pharmacode reads this way

3. How to use Optrex Bloodshot Eye Drops

**Applying the drops**

To be used in the eye(s) only.

1. Check the seal is not broken before first use.
2. Wash and dry your hands.
3. Remove the plastic cap from the eye drop tip.
4. Tilt your head back while seated or lie down on your back.
5. Gently pull the lower eyelid downwards and look up.
6. Place the tip of the bottle close to your eye. Squeeze the bottle and let one drop fall into the space between the lower eyelid and the eye. Take care that the tip of the bottle does not touch your eye, the skin around your eye or your fingers.
7. Close your eye for a moment.
8. When both eyes are to be treated, repeat for the other eye.
9. Replace the cap straight away.

Whilst applying the drops, **do not**

- breathe on or touch the dropper nozzle
- touch the eyes or eyelids with the dropper nozzle
- share your eye drops with anyone else.

**For external use only.**  
 If anyone accidentally swallows the contents of the bottle seek medical advice.

**Dose**  
 Gently squeeze **one or two drops** into each eye. Repeat as necessary. **Do not use more than four times in 24 hours.**

Continued use may increase redness of the eye. Do not use the medicine if it changes colour or becomes cloudy.

If you have increased redness of the eye or no improvement after 24 hours, stop taking the product and see a doctor.

**If you take more medicine than you should:**  
 Speak to your doctor or pharmacist immediately. Symptoms of overdose by mouth may include nausea, headache, low body temperature, bradycardia, sweating, drowsiness, coma, particularly in children.

**If you forget to take the medicine:**  
 Take the next dose when needed. Do not take more than 4 doses in 24 hours.

**4. Possible side effects**  
 Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Stop using the product and seek medical help immediately if you experience:**

- **allergic reactions** such as itching, rashes and other
- **pain in the eye or change in vision**
- **stinging or irritation** in the eye caused by using too many drops or using them too often

**Other side effects may include:**

- Eye irritation, eye pain. Continued and excessive use may lead to increased redness of the eye and to developing of other eye conditions.
- nausea
- headache, dizziness

**Reporting of side effects**  
 If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

**5. How to store Optrex Bloodshot Eye Drops**

Keep all medicines out of the sight and reach of children.

**Store below 25°C.** Do not freeze.  
 Do not use the drops after the expiry date (printed on the dropper bottle and carton).


You must throw away unused drops 28 days after first opening the bottle.  
 The preservative can only keep them sterile in use for a limited time.  
 Medicines should not be disposed of via waste water or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help protect the environment.

**6. Contents of the pack and other information**

Optrex Bloodshot Eye Drops contain the active ingredients Naphazoline and Hamamelis Virginiana. These are present as Naphazoline Hydrochloride PhEur 0.01% w/v and Distilled Witch Hazel BPC 12.5% v/v.

Other ingredients are Purified Water, Glycerin, the preservative Benzalkonium Chloride, Borax and Boric Acid (to balance the pH to help avoid stinging) and Disodium Edetate. The pack contains 10ml of the drops.

Licence holder: Optrex Limited, Slough, SL1 3UH, UK.  
 Manufacturer: Reckitt Benckiser Healthcare International Ltd, Nottingham, NG90 2DB. PL 00062/0024.  
 Leaflet revised August 2020.



00000000



**CUSTOMER INFO:**  
 Minimum Point Size = 9.00pt