

# Buspirone 5mg and 10mg tablets

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

## What is in this leaflet

- 1 What Buspirone tablets are and what they are used for**
- 2 What you need to know before you take Buspirone tablets**
- 3 How to take Buspirone tablets**
- 4 Possible side effects**
- 5 How to store Buspirone tablets**
- 6 Contents of the pack and other information**

## 1 What Buspirone tablets are and what they are used for

Buspirone tablets belong to a group of medicines called anxiolytics. These medicines work on the central nervous system, altering levels of chemicals in the brain.

Buspirone tablets may be used for the:

- short term management of anxiety disorders
- relief of symptoms of anxiety with or without symptoms of depression.

## 2 What you need to know before you take Buspirone tablets

**Do not take** Buspirone tablets if you:

- are **allergic** to buspirone hydrochloride or any of the other ingredients of this medicine (listed in section 6).
- are **pregnant** or **breast-feeding**.
- have **epilepsy**.
- have **severely** impaired **liver** or **kidney** function.

### Warnings and precautions

Talk to your doctor or pharmacist before you take Buspirone tablets if you have:

- had impaired **liver** or **kidney** function in the past.
- been prescribed a **benzodiazepine** e.g. nitrazepam or temazepam or another common **sedative** or **hypnotic** medicine. You should be gradually withdrawn from these medicines before taking Buspirone tablets.
- **acute narrow-angle glaucoma**
- **myasthenia gravis**, a disorder characterised by muscle weakness, difficulty chewing or swallowing and slurred speech
- have had **drug dependence**

## Other medicines and Buspirone tablets

**Tell your doctor or pharmacist** if you are taking, have recently taken or might take any other medicines. Especially:

- monoamine-oxidase inhibitors (MAOIs) such as phenelzine and tranylcypromine (for depression)
- St. John's Wort, nefazodone and L-tryptophan, fluvoxamine, trazodone (for depression)
- selective serotonin re-uptake inhibitors (SSRIs) e.g. fluoxetine and paroxetine (for depression)
- haloperidol and lithium (for mental illness)
- calcium channel blockers such as diltiazem and verapamil (to treat high blood pressure)
- rifampicin (to treat tuberculosis)
- triptan drugs e.g. sumatriptan (to treat migraine)
- tramadol (a painkiller)
- baclofen (a muscle relaxant)
- lofexidine (to manage drug withdrawal)
- nabilone (to treat nausea and vomiting)
- antihistamines (to treat allergic reactions)
- erythromycin, itraconazole and linezolid (to treat infections)
- benzodiazepines e.g. nitrazepam or temazepam or another common sedative or hypnotic medicine
- diltiazem (to treat angina)
- digoxin (to treat heart failure)
- phenobarbital, phenyton, carbamazepine (to treat epilepsy)
- cimetidine (to treat stomach ulcers)
- diazepam (to treat anxiety)
- warfarin (to treat blood clots)

## Buspirone tablets with food or drink

**Talk to your doctor** before eating or drinking products containing **grapefruit juice**, whilst taking Buspirone tablets. You should not drink **alcohol** whilst taking Buspirone tablets.

## Pregnancy and breast-feeding

Do not take Buspirone tablets if you are pregnant or are breast-feeding. Talk to your doctor before taking any medicine.

## Driving and using machines

Buspirone tablets may make you feel drowsy or dizzy. Make sure you are not affected before you drive or operate machinery.

The medicine can affect your ability to drive as it may make you sleepy or dizzy.

- Do not drive while taking this medicine until you know how it affects you.
- It is an offence to drive if this medicine affects your ability to drive.
- However, you would not be committing an offence if:
  - The medicine has been prescribed to treat a medical or dental problem and
  - You have taken it according to the instructions given by the prescriber or in the information provided with the medicine and -
- It was not affecting your ability to drive safely

Talk to your doctor or pharmacist if you are not sure whether it is safe for you to drive while taking this medicine.

## Sugar intolerance

If you have been told you have an intolerance to some sugars, contact your doctor before taking this medicine, as it contains a sugar called lactose.

## 3 How to take Buspirone tablets

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Swallow the tablets **with water**, at the same time each day. Buspirone should be taken consistently with or without food. However, the medicine is taken on the day one should be continued thereafter.

### Doses:

#### Adults (including the elderly)

The recommended starting dose is 5mg two to three times a day, which may be increased every two to three days. The usual dose you will be maintained on is 15mg to 30mg a day in divided doses. The maximum daily dosage should not exceed 60mg per day.

**Children:** Not recommended.

If you have impaired liver or kidney function, your doctor may prescribe you a lower dose.

### If you take more than you should

If you (or someone else) swallow a lot of the tablets at the same time, or if you think a child has swallowed any, contact your nearest hospital casualty department or tell your doctor immediately. Symptoms of an overdose include feeling or being sick, headache, dizziness, drowsiness, ringing or buzzing in the ears, restlessness, restriction of the pupils, stomach problems, slow heart beat, low blood pressure, fits and extrapyramidal symptoms (difficulty in speaking or swallowing, loss of balance control, mask-like face, shuffling walk, stiffness of arms and legs, trembling or shaking of hands or fingers).

### If you forget to take the tablets

Do not take a double dose to make up for a forgotten dose. If you forget to take a dose take it as soon as you remember it and then take the next dose at the right time.

### If you stop taking the tablets

Talk to your doctor before you stop taking the tablets and follow their advice.

## 4 Possible side effects

Like all medicines, Buspirone tablets can cause side effects although not everybody gets them.

**Stop taking Buspirone tablets and contact your doctor immediately** if you have:

- **Serotonin syndrome**, (feeling confused, feeling restless, sweating, shaking, shivering, hallucinations (strange visions or sounds), sudden jerks of the muscles or a fast heartbeat).
- **Allergic reaction**, itching, difficulty breathing or swelling of the face, lips, throat or tongue. Allergic conditions (causing itchy and swollen skin).

**Very common:** may affect more than 1 in 10 people

- Dizziness, headache, sleepiness

**Common:** may affect up to 1 in 10 people

- Nervousness, difficulty in sleeping, disturbance in attention, depression, confused state, sleeping problems, anger, pins and needles, tingling, blurred vision, abnormal coordination, tremor, ringing or buzzing in the ears, racing heart (abnormally fast heart rhythm), chest pain, blocked nose, sore throat, feeling and being sick, stomach pain, dry mouth,

diarrhoea, constipation, cold sweats, rash, pain in the muscles, tendons, ligaments and bone, mental and physical tiredness

**Rare:** may affect up to 1 in 1,000 people

- Bruising, itchy skin rash caused by allergic reaction - pale or red irregular raised patches with severe itching (hives)

**Very rare:** may affect up to 1 in 10,000 people

- Psychosis, hallucinations, feeling of unreality to the self, that the mind is separated from the body, effect on your liability, fits, tunnel vision, extrapyramidal disorder (tremor, slurred speech, restlessness (akathisia), dystonia (affecting posture - muscle spasm or the neck, shoulders or body), anxiety, distress), muscle rigidity, difficulty controlling movement/jerky movements, disorder affecting posture (muscle spasm of neck, shoulders and body), fainting or passing out, total or partial memory loss, lack of muscle control, co-ordination, difficulty in controlling movements, tremor, stiffness and shuffling, restless leg syndrome, restlessness, difficulty passing water, over production of breast milk

## Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

## 5 How to store Buspirone tablets

Keep out of the sight and reach of children. Do not store above 25°C. Store in the original container.

Do not use Buspirone tablets after the expiry date stated on the label/carton/bottle. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

## 6 Contents of the pack and other information

### What Buspirone tablets contain

- The active substance is buspirone hydrochloride. Each tablet contains either 5mg or 10mg of the active substance.
- The other ingredients are lactose monohydrate, magnesium stearate, purified talc, polyvidone K-25 and potato starch.

### What Buspirone tablets look like and contents of the pack

Buspirone tablets are white, uncoated, flat bevelled edge tablets. Pack size is 30.

### Marketing Authorisation Holder and Manufacturer

Accord, Barnstaple, EX32 8NS, UK

This leaflet was last revised in August 2019



If you would like a leaflet with larger text, please contact 01271 385257.