

PACKAGE LEAFLET – INFORMATION FOR THE USER

Citalopram 10mg Tablets **Citalopram 20mg Tablets** **Citalopram 40mg Tablets** (Citalopram Hydrobromide)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet., See section 4.

What is in this leaflet:

1. What are Citalopram Tablets and what are they used for
2. What you need to know before you take Citalopram Tablets
3. How to take Citalopram Tablets
4. Possible side effects
5. How to store Citalopram Tablets
6. Contents of the pack and other information

1. What are Citalopram tablets and what are they used for

Citalopram is a Selective Serotonin Reuptake Inhibitors (SSRIs) and belongs to a group of medicines called antidepressants. These medicines help to correct certain chemical imbalances in the brain that are causing the symptoms of your illness.

Citalopram Tablets are used for the treatment of depression and when you feel better, to help prevent these symptoms recurring. Citalopram tablets is also used for long-term treatment to prevent the occurrence of new episodes of depression or if you have recurrent depression.

Citalopram Tablets are also used to relieve symptoms if you are suffering from panic attacks.

2. What you need to know before you take Citalopram tablets

Do not take Citalopram Tablets and talk to your doctor:

- if you are allergic (hypersensitive) to citalopram or to any of the other ingredients of Citalopram Tablets (see **What Citalopram Tablet contains**, section 6).
- if you are also taking a medicine containing pimozide (Antipsychotic).

- if you are taking medicines known as monoamine oxidase inhibitors (MAOIs).
- MAOIs include medicines such as phenelzine, iproniazid, isocarboxazid, nialamide, tranylcypromine, moclobemide (used for the treatment of depression), selegiline (used in the treatment of Parkinson's disease) and linezolid (an antibiotic). Even if you have finished taking one of the following MAOIs: phenelzine, iproniazid, isocarboxazid, nialamide or tranylcypromine then you will need to wait 2 weeks before you start taking your Citalopram Tablets. One day must elapse after you have finished taking moclobemide. After stopping Citalopram Tablets you must allow 1 week before taking any MAOI.
- if you are born with or have had an episode of abnormal heart rhythm (seen at ECG; an examination to evaluate how the heart is functioning)
- if you take medicines for heart rhythm problems or that may affect the heart's rhythm. Also refer to the section "Taking other medicines" below.

Warnings and precautions

- If you suffer or have suffered from heart problems or have recently had a heart attack
- If you have a low resting heart-rate and/or you know that you may have salt depletion as a result of prolonged severe diarrhoea and vomiting (being sick) or usage of diuretics (water tablets)
- If you experience a fast or irregular heartbeat, fainting, collapse or dizziness on standing up which may indicate abnormal functioning of the heart rate.
- Medicines like Citalopram tablets (so called SSRIs/SNRIs) may cause symptoms of sexual dysfunction (see section 4). In some cases, these symptoms have continued after stopping treatment.

Please tell your doctor if you are suffering from

- Liver disease.
- Severe kidney disease.
- Diabetes (you may need an adjustment of your antidiabetic therapy).
- Epilepsy or a history of seizures or fits.
- A bleeding disorder or have ever suffered from bleeding in the stomach or intestine or if you are pregnant (see 'pregnancy').
- Eye problems, such as certain types of glaucoma.
- Mania or panic disorder.
- Low blood levels of sodium.
- ECT (electroconvulsive therapy).
- Problems with your eyes, such as certain kinds of glaucoma.
- Suffered or suffer from heart problems or have recently had a heart attack.
- A low resting heart-rate and/or you know that you may have salt depletion as a result of
- which may indicate abnormal functioning of the heart rate.

Please consult your doctor, even if these statements were applicable to you at any time in the past.

Please note:

Some patients with manic-depressive illness may enter into a manic phase. This is characterized by unusual and rapidly changing ideas, inappropriate happiness and excessive physical activity. If you experience this, contact your doctor.

Symptoms such as restlessness or difficulty in sitting or standing still can also occur during the first weeks of the treatment. Tell your doctor immediately if you experience these symptoms.

Special information relating to your disease

As with other medicines used to treat depression or related diseases, the improvement is not achieved immediately. After the start of Citalopram treatment it may take several weeks before you experience any improvement. In the beginning of the treatment certain patients may experience increased anxiety, which will disappear during continued treatment. Therefore, it is very important that you follow exactly your doctor's orders and do not stop the treatment or change the dose without consulting your doctor.

Thoughts of suicide and worsening of your depression or anxiety disorder

If you are depressed and/or have anxiety disorders you can sometimes have thoughts of harming or killing yourself. These may be increased when first starting antidepressants, since these medicines all take time to work, usually about two weeks but sometimes longer.

You may be more likely to think like this:

- if you have previously had thoughts about killing or harming yourself.
- if you are a young adult. Information from clinical trials has shown an increased risk of suicidal behaviour in adults aged less than 25 years with psychiatric conditions who were treated with an antidepressant.

If you have thoughts of harming or killing yourself at any time, **contact your doctor or go to a hospital straight away.**

You may find it helpful to tell a relative or close friend that you are depressed or have an anxiety disorder, and ask them to read this leaflet. You might ask them to tell you if they think your depression or anxiety is getting worse, or if they are worried about changes in your behaviour.

Use in children and adolescents under 18 years of age

Citalopram tablets should normally not be used for children and adolescents under 18 years. Also, you should know that patients under 18 years have an increased risk of side-effects such as suicide attempt, suicidal thoughts and hostility (predominantly aggression, oppositional behaviour and anger) when they take this class of medicines. Despite this, your doctor may prescribe citalopram for patients under 18 years because he/she decides that this is in their best interests. If your doctor has prescribed Citalopram tablets for a patient under 18 years and if you want to discuss this, please go back to your doctor. You should inform your doctor if any of the symptoms listed above develop or

worsen when patients under 18 years are taking Citalopram tablets. Also, the long-term safety effects concerning growth, maturation and cognitive and behavioural development of Citalopram in this age group have not yet been demonstrated.

Other medicines and Citalopram tablets

Medicines may affect the action of other medicines and this can sometimes cause serious adverse reactions. Please tell your doctor if you are taking or have taken any other medicines (including those purchased without prescription) during the last 14 days. This includes other medicines for depression (see **Do not take Citalopram tablets** above).

- The herbal remedy St John's Wort (*Hypericum perforatum*). This should not be taken at the same time as Citalopram.
- Monoamine oxidase inhibitors (MAOIs). These should not be taken at the same time as Citalopram (see Do not take Citalopram tablets above).
- Buprenorphine (a type of opioid medicine). Concomitant use increases the risk of serotonin syndrome, a potentially life-threatening condition. See "Other Warnings"

Tell your doctor if you are taking any of the following medicines:

- Sumatriptan (used to treat migraine) or tramadol (a pain killer). If you feel unwell when using these medicines with Citalopram tablets then you should see your doctor.
- Lithium (used to prevent and treat mania) and tryptophan (an anti-depressant)
- Medicines known to affect the blood platelets (e.g. anticoagulant drugs used to treat or prevent blood clots; aspirin and non-steroidal anti-inflammatory drugs (NSAIDs) such as ibuprofen and diclofenac used as painkillers and some antipsychotic drugs and tricyclic antidepressants).
- Metoprolol, a beta blocker used to treat migraine, some heart conditions and high blood pressure. The effects of either drug could be increased, decreased or altered.
- Linezolid (an antibiotic).
- Imipramine and desipramine (used to treat depression).
- Medicines containing selegiline (used to treat Parkinson's disease)
- Cimetidine (used to treat stomach ulcers).
- Fluconazole (used to treat fungal infections).
- Mefloquine (used to treat malaria).
- Bupropion (used to treat depression).
- Neuroleptics (used in the treatment of schizophrenia).

DO NOT TAKE CITALOPRAM TABLETS if you take medicines for heart rhythm problems or medicines that may affect the heart's rhythm, e.g. such as Class IA and III antiarrhythmics, antipsychotics (e.g. phenothiazine derivatives, fentiazine derivatives, pimozide, haloperidol), tricyclic antidepressants, certain antimicrobial agents (e.g. sparfloxacin, moxifloxacin, erythromycin IV, pentamidine, anti-malarial treatment particularly halofantrine), certain antihistamines (astemizole, mizolastine). If you have any further questions about this you should speak to your doctor.

Citalopram tablets with food and drink

Citalopram tablets can be taken with or without food (see section 3 “How to take Citalopram tablets”).

As with all antidepressants, it is sensible to avoid drinking alcohol whilst receiving treatment although Citalopram tablets has not been shown to increase the effects of alcohol.

Pregnancy

If you are pregnant, think you might be pregnant, or are trying to become pregnant, tell your doctor. Do not take Citalopram tablets if you are pregnant unless you and your doctor have discussed the risks and benefits involved.

Make sure your midwife and/or doctor know you are on Citalopram Tablets. When taken during pregnancy, particularly in the last 3 months of pregnancy, medicines like Citalopram Tablets may increase the risk of a serious condition in babies, called persistent pulmonary hypertension of the new born (PPHN), making the baby breathe faster and appear bluish. These symptoms usually begin during the first 24 hours after the baby is born. If this happens to your baby you should contact your midwife and/or doctor immediately.

Also, if you take Citalopram during the last 3 months of your pregnancy and until the date of birth you should be aware that the following effects may be seen in your newborn: fits, being too hot or cold, feeding difficulties, vomiting, low blood sugar, stiff or floppy muscles, overactive reflexes, tremor, jitteriness, irritability, lethargy, constant crying, sleepiness or sleeping difficulties. If your newborn baby gets any of these symptoms please contact your midwife and/or doctor immediately.

If you take Citalopram Tablets near the end of your pregnancy there may be an increased risk of heavy vaginal bleeding shortly after birth, especially if you have a history of bleeding disorders. Your doctor or midwife should be aware that you are taking Citalopram Tablets so they can advise you.

Ask your doctor or pharmacist for advice before taking any medicine.

Breast-feeding

If you are breast-feeding, ask your doctor for advice. You should not breast-feed your baby while taking Citalopram tablets because small amounts of the medicine can pass into the breast milk.

Fertility

Citalopram has been shown to reduce the quality of sperm in animal studies. Theoretically, this could affect fertility, but impact on human fertility has not been observed as yet.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Citalopram tablets does not usually affect the ability to carry out normal daily activities. However if you feel dizzy or sleepy when you start to take this medicine, you should be careful when driving, operating machinery or performing jobs that need you to be alert until these effects wear off.

Citalopram Tablets contains Lactose

These tablets contain lactose. If you have been told by your doctor that you have intolerance to some sugars, contact your doctor before taking Citalopram tablets.

3. How to take Citalopram tablets

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Adults

Depression

The usual dose is 20 mg per day. This may be increased by your doctor to a maximum of 40 mg per day.

Panic disorder

The starting dose is 10 mg per day for the first week before increasing the dose to 20-30 mg per day. The dose may be increased by your doctor to a maximum of 40 mg per day.

Elderly patients (above 65 years of age)

The starting dose should be decreased to half of the recommended dose, e.g. 10-20 mg per day. Elderly patients should not usually receive more than 20 mg per day.

Children and adolescents (less than 18 years)

Citalopram tablets is not recommended for children or adolescents. For further information, please see section 2, **What you need to know before you take Citalopram tablets.**

Patients with special risks

Patients with liver complaints should not receive more than 20 mg of Citalopram tablets per day.

How and when to take Citalopram tablets

Citalopram tablets is taken every day as a single daily dose. Citalopram tablets can be taken any time of the day with or without food. Swallow the tablets with a drink of water. Do not chew them (they have a bitter taste).

Duration of treatment

Like other medicines for depression and panic disorder these tablets may take a few weeks before you feel any improvement. Continue to take Citalopram tablets even if it takes some time before you feel any improvement in your condition.

The duration of treatment is individual, usually at least 6 months. Continue to take Citalopram tablets for as long as your doctor recommends. Do not stop taking them even if you begin to feel better, unless you are told to do so by your doctor. The underlying illness may persist for a long time and if you stop your treatment too soon your symptoms may return.

Patients who have recurrent depression benefit from continued treatment, sometimes for several years, to prevent the occurrence of new depressive episodes.

Never change the dose of the medicine without talking to your doctor first.

If you have taken more Citalopram Tablets than you should,

If you or someone else swallows several of these tablets all together, or you think a child has swallowed any of these tablets, contact your doctor or pharmacist or hospital emergency department immediately. Always take any tablets left over with you, also the box and leaflet as this will allow easier identification of the tablets.

Symptoms of overdose may include:

- Nausea (feeling sick)
- Vomiting
- Sweating
- Drowsiness
- Seizures
- Unconsciousness
- Fast heart beats
- Tremor
- Irregular heart beat
- Changes in heart rhythm
- Changes in blood pressure
- Serotonin syndrome (see Section 4)
- Agitation
- Dizziness
- Enlarged eye pupils
- Bluish skin
- Breathing too quickly

If you forget to take Citalopram Tablets

If you forget to take a dose, take it as soon as you remember. If it is almost time for your next dose, do not take the missed dose and just carry on as before. Do not take a double dose to make up for a forgotten dose.

If you stop taking Citalopram Tablets Stopping this medicine quickly may cause symptoms such as dizziness, nausea and numbness or tingling in hands or feet, sleep disturbances (vivid dreams, nightmares, inability to sleep), feeling anxious, headaches, feeling or being sick, sweating, feeling restless or agitated, tremor, feeling confused or disorientated, feeling emotional or irritable, diarrhoea (loose stools), visual disturbances, fluttering or pounding heartbeat (palpitations). These symptoms are generally non-serious and disappear within a few days. When you have completed your course of treatment, the dose of Citalopram is usually reduced gradually over a couple of weeks.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

Like all medicines, this medicine can have side effects and some people may experience unwanted effects (side effects) whilst taking Citalopram tablets. Several of the effects listed below can also be symptoms of your illness and may disappear as you start to get better.

Serious side effects

Stop taking Citalopram tablets and seek medical advice immediately if you suffer from any of the following allergic reactions:

- Difficulty in breathing.
- Swelling of the face, lips, tongue or throat that causes difficulty in swallowing or breathing.
- Severe itching of the skin (with raised lumps)
- Fast, irregular heart beat, fainting which could be symptoms of a life-threatening condition known as torsades de pointes.

If you notice any of the following symptoms you should stop taking Citalopram tablets and see your doctor immediately:

- if you start having fits for the first time or fits that you have suffered from in the past become more frequent.
- if your behaviour changes because you feel elated or over excited.
- if you experience high fever, agitation, confusion, trembling or abrupt contractions of muscles. These may be signs of a rare condition called serotonin syndrome.
- Tiredness, confusion and twitching of your muscles. These may be signs of a low blood level of sodium.

If you have thoughts of harming or killing yourself at any time, **contact your doctor or go to a hospital straight away.**

The following side effects are often mild and usually disappear after a few days' treatment.

Very common side effects (likely to affect more than 1 in 10 people)

- Feeling sick (Nausea),
- Dry mouth (a dry mouth increases the risk of tooth decay, so be sure to clean your teeth more often than usual),
- Increased sweating,
- Sleepiness,
- Difficulty in sleeping
- Headache
- Changes in your sleeping pattern
- Loss of body strength, weakness

Common side effects (likely to affect up to 1 in 10 people)

- Lack of appetite
- Agitation
- Decreased sex drive
- Anxiety
- Nervousness
- Confusion
- Abnormal dreams
- Reduced emotions, indifference (apathy)
- Tremor
- Tingling or numbness in the hands or feet
- Dizziness
- Problems concentrating
- Migraine
- Loss of memory (amnesia)
- Ringing in the ears (tinnitus)
- Palpitations
- Yawning
- Diarrhoea
- Vomiting
- Constipation
- Stomach pain
- Flatulence (wind)
- Itching
- Pain in muscles and joints
- For men, problems with ejaculation and erection
- For females, failing to reach an orgasm
- Tiredness
- Prickling of the skin
- Loss of weight

Uncommon (likely to affect up to 1 in every 100 people)

- Bruising easily
- Increased appetite
- Aggression
- Hallucinations
- Mania
- Fainting
- Large pupils (the dark centre of the eye)
- Fast heart beat
- Slow heart beat
- Nettle rash
- Loss of hair
- Rash
- Sensitivity to sunlight
- Difficulties urinating

- Excessive menstrual bleeding
- Swelling of the arms or legs
- Increased weight

Rare (likely to affect up to 1 in every 1000 people)

- Convulsions
- Increased sex drive
- Involuntary movements
- Taste disturbances
- Bleeding
- Coughing
- Hepatitis
- Feeling unwell (malaise)

Some patient have reported (frequency not known)

- Thoughts of harming or killing themselves
- An increase in bleeding or bruising caused by a decrease in blood platelets
- Rash (hypersensitivity)
- Low potassium levels in the blood (hypokalaemia), which can cause muscle weakness, twitching or abnormal heart rhythms
- Panic attack
- Grinding teeth
- Restlessness
- Unusual muscle movements or stiffness
- Involuntary movements of the muscles (akathisia)
- Low blood pressure
- Nosebleed
- Bleeding disorders including skin and mucosal bleeding (ecchymosis)
- Sudden swelling of skin or mucosa
- In men, painful erections
- Flow of breast milk in men or in women who are not breast-feeding (galactorrhoea)
- Abnormal liver function tests
- An increased risk of bone fractures has been observed in patients taking this type of medicines.
- Abnormal heart rhythm
- Heavy vaginal bleeding shortly after birth (postpartum haemorrhage), see Pregnancy in section 2 for more information

SSRIs can, very rarely, increase the risk of bleeding, including stomach or intestinal bleeding. Let your doctor know if you vomit blood or develop black or blood stained stools.

Also let your doctor know if you continue to have other symptoms associated with your depression. This might include hallucinations, anxiety, mania or confusion.

Any side effects that do occur will usually disappear after a few days. If they are troublesome or persistent, or if you develop any other unusual side effects while taking Citalopram tablets, please tell your doctor.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Citalopram tablets

Keep out of the sight and reach of children.

This medicinal product does not require any special storage conditions.

Do not use your tablets after the expiry date stated on the label or carton.

Medicines should not be disposed of via waste water or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Contents of the pack and other information

What Citalopram Tablet contains?

- The active substance is citalopram hydrobromide.
- The inactive ingredients are lactose, microcrystalline cellulose, maize starch, copovidone, croscarmellose sodium and magnesium stearate. The ingredients of the film coating material, Opadry White 20H 58983 (comprise of hypromellose, titanium dioxide (E171), propylene glycol, hydroxypropyl cellulose and talc).(see section 2 for Important information about some of the ingredients of Citalopram tablets).

What Citalopram Tablet looks like and contents of the pack?

Citalopram 10mg tablets are white to off white, round, biconvex, film coated tablets debossed with 'DT' on one side and plain on the other side.

Citalopram 20mg tablets are white to off white, round, biconvex, film coated tablets debossed with 'DU' on one side and lip shaped breakline on the other side.

Citalopram 40mg tablets are white to off white, round, biconvex, film coated tablets debossed with 'DZ' on one side and lip shaped breakline on the other side.

Citalopram tablets are available as blister strips in pack sizes of 1, 14, 20, 28, 30, 50, 56, 98, 100 or 250 tablets. (Not all pack sizes may be marketed).

Marketing Authorisation Holder and Manufacturer

Accord Healthcare Limited,
Sage House, 319 Pinner Road,
North Harrow, Middlesex,
HA1 4HF, United Kingdom

This leaflet was last approved in 12/2020.