

Package leaflet: Information for the patient

ENALAPRIL MALEATE 5 mg TABLETS
ENALAPRIL MALEATE 10 mg TABLETS
ENALAPRIL MALEATE 20 mg TABLETS
(Enalapril Maleate)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist. If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Enalapril Tablets are and what they are used for
2. What you need to know before you take Enalapril Tablets
3. How to take Enalapril Tablets
4. Possible side effects
5. How to store Enalapril Tablets
6. Contents of the pack and other information

1. What Enalapril Tablets are and what they are used for

Enalapril Tablets belong to a group of medicines called ‘ACE inhibitors’(angiotensin converting enzyme inhibitors).

Enalapril tablets are used

- to treat high blood pressure (hypertension).
- to treat heart failure (weakening of heart function). It can lower the need to go to hospital and can help some patients live longer.
- to prevent the signs of heart failure. The signs include: shortness of breath, tiredness after light physical activity such as walking, or swelling of the ankles and feet.

Enalapril works by widening your blood vessels. This lowers your blood pressure. The medicine usually starts to work within an hour, and the effect lasts for at least 24 hours. Some people will require several weeks of treatment until the best effect on your blood pressure is seen.

2. What you need to know before you take Enalapril Tablets

Do not take Enalapril Tablets if:

- you are allergic (hypersensitive) to enalapril maleate or any of the other ingredients of Enalapril tablets (listed in Section 6) or to any other ACE inhibitors similar to Enalapril tablets.
- you ever had swelling of your face, lips, mouth, tongue or throat which caused difficulty in swallowing or breathing (angioedema) when the reason why was not known or it was inherited.
- you have diabetes or impaired kidney function and you are treated with a blood pressure lowering medicine containing aliskiren
- you are more than 3 months pregnant. (It is also better to avoid Enalapril Tablets in early pregnancy- see pregnancy section.)

Do not take Enalapril Tablets if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Enalapril Tablets.

Warnings and Precautions

Check with your doctor or pharmacist before taking Enalapril Tablets if any of the following apply to you.

- If you think you are (or might become) pregnant. Enalapril Tablet is not recommended in early pregnancy, and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see pregnancy section).
- If you are breast-feeding or about to start breast-feeding (see breast-feeding section).
- If you have a kidney problem (including kidney transplantation). These may lead to higher levels of potassium in your blood which can be serious. Your doctor may need to adjust your dose of Enalapril tablets or monitor your blood level of potassium.
- If you are having dialysis.
- If you have been very sick suffered from excessive vomiting or had bad diarrhoea recently.
- If you are on a salt-restricted diet, are taking potassium supplements, potassium-sparing agents, or potassium-containing salt substitutes.
- If you are over 70 years of age.
- If you have a heart problem.
- If you have a condition involving the blood vessels in the brain.

- If you suffer from blood problems such as low or lack of white blood cells (neutropenia / agranulocytosis), low blood platelet count (thrombocytopenia) or a decreased number of red blood cells (anaemia).
- If you have a liver problem.
- If you have collagen vascular disease (e.g. lupus erythematosus, rheumatoid arthritis or scleroderma), are on therapy that suppress your immune system, are taking drugs like allopurinol (used for the treatment of gout) or procainamide (used to treat abnormal heart rhythms) or any combinations of these. Your doctor may periodically take a blood sample to check your white cell count. If any of these apply to you and you develop an infection (symptoms may be high temperature, feverish), you should let your doctor know immediately.
- If you are taking an mTOR inhibitor (e.g., temsirolimus, sirolimus, everolimus: medicines used to treat certain types of cancer or to prevent the body's immune system from rejecting a transplanted organ). You may be at increased risk for an allergic reaction called angioedema.
- If you have a history of angioedema (symptoms such as itching, nettle rash, wheezing or swelling of the hands, throat, mouth and eye lids) while taking other medicines. You should be aware that black patients are at increased risk of these types of reactions to ACE inhibitors.
- If you have low blood pressure (you may notice this as faintness or dizziness, especially when standing).
- If you have diabetes and are taking antidiabetic agents or insulin to control your diabetes; you should closely monitor your blood for low blood glucose levels especially during the first month of treatment. The level of potassium in your blood can also be higher.
- if you are taking any of the following medicines used to treat high blood pressure:
 - an angiotensin II receptor blocker (ARBs) (also known as sartans - for example valsartan, telmisartan, irbesartan), in particular if you have diabetes-related kidney problems.
 - aliskiren

Your doctor may check your kidney function, blood pressure, and the amount of electrolytes (e.g. potassium) in your blood at regular intervals. See also information under the heading "Do not take Enalapril Tablets"

You must tell your doctor if you think you are (or might become) pregnant. This medicine is not recommended in early pregnancy and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see Pregnancy section).

You should be aware that Enalapril Tablets lowers the blood pressure in black patients less effectively than in non-black patients.

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking Enalapril Tablets.

Please let your doctor know immediately if you develop any of the following symptoms while taking Enalapril Tablets

- Jaundice (yellowing of the skin and whites of the eyes)
- A dry cough which is persistent for a long time

If you going to undergo any of the below mentioned procedures, please tell the doctor who is treating you that you are taking Enalapril Tablets.

- a treatment called LDL apheresis, which is removal of cholesterol from your blood.
- a desensitisation treatment to reduce the effect of an allergy to bee or wasp stings.
- any surgery or receive anaesthetics (even at the dentist).

If any of the above applies to you, talk to your doctor or dentist before the procedure.

Other medicines and Enalapril Tablets

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This includes herbal medicines. This is because Enalapril Tablets can affect the way some medicines work. Also some other medicines can affect the way Enalapril Tablets works. Your doctor may need to change your dose and/or to take other precautions:

In particular tell your doctor or pharmacist if you are taking any of the following medicines:

- an angiotensin II receptor blocker (ARB) or aliskiren (see also information under the headings “Do not take Enalapril Tablets if” and “Warnings and precautions”)
- other medicines to lower blood pressure, such as beta-blockers or water tablets (diuretics)
- medicines containing potassium (including dietary salt substitutes)
- medicines for diabetes (including oral antidiabetic medicines and insulin)
- lithium (a medicine used to treat a certain kind of depression)
- medicines for depression called ‘tricyclic antidepressants’
- medicines for mental problems called ‘antipsychotics’
- certain cough and cold medicines and weight reducing medicines which contain something called a ‘sympathomimetic agent’
- certain pain or arthritis medicines including gold therapy
- an mTOR inhibitor (e.g., temsirolimus, sirolimus, everolimus; medicines used to treat certain types of cancer or to prevent the body’s immune system from rejecting a transplanted organ). See also information under the heading “Warnings and precautions”
- non-steroidal anti-inflammatory drugs, including COX-2-inhibitors (medicines that reduce inflammation, and can be used to help relieve pain)
- aspirin (acetylsalicylic acid)
- medicines used to dissolve blood clots (thrombolytics)
- alcohol.

If you are not sure if any of the above applies to you, talk to your doctor or pharmacist before taking Enalapril tablets.

Taking Enalapril Tablets with food and drink

Enalapril Tablets can be taken with or without food. Most people take Enalapril tablets with a drink of water.

Preganancy and breast-feeding

Pregnancy

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor will normally advise you to stop taking Enalapril Tablets before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Enalapril Tablets. Enalapril Tablets is not recommended in early pregnancy, and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

Breast-feeding

Tell your doctor if you are breast-feeding or about to start breast-feeding. Breast-feeding new born babies (first few weeks after birth), and especially premature babies, is not recommended whilst taking Enalapril Tablets.

In case of an older baby your doctor should advise you on the benefits and risks of taking Enalapril Tablets whilst breast-feeding, compared to other treatments.

Driving and using machines

You may feel dizzy or sleepy while taking Enalapril tablets. If this happens, do not drive or use any tools or machines.

Enalapril tablets contain lactose

These tablets contain Lactose, which is a type of sugar. If your doctor has told you that you have intolerance to some sugars, contact your doctor before taking Enalapril Tablets.

3. How to take Enalapril Tablets

Always take this medicine exactly as your doctor as told you. You should check with your doctor or pharmacist if you are not sure.

- It is very important to continue taking this medicine for as long as your doctor prescribes it
- Do not take more tablets than prescribed.
- The score line is only there to help you break the tablet if you have difficulty swallowing it whole

The dose you take will depend on your condition and whether you are taking any other treatment. The usual dose is:

High Blood Pressure

- The usual starting dose ranges from 5 to 20 mg taken once a day.
- Some patients may need a lower starting dose.

- The usual long term dose is 20 mg taken once a day.
- The maximal long term dose is 40 mg taken once a day.

Heart Failure

- The usual starting dose is 2.5 mg taken once a day.
- Your doctor will raise this amount step by step until the dose that is right for you has been achieved.
- The usual long term dose is 20 mg each day, taken in one or two doses.
- The maximal long term dose is 40 mg each day, divided in two doses.

Patients with kidney problems

Your dose of medicine will be changed depending on how well your kidneys are working:

- Moderate kidney problems - 5 mg to 10 mg each day.
- Severe kidney problems - 2.5 mg each day.
- If you are having dialysis - 2.5 mg each day. On days you are not having dialysis, your dose may be changed depending on how low your blood pressure is.

Elderly patients

Your dose will be decided by your doctor, and will be based on how well your kidneys are working.

Use in Children

Use of Enalapril tablets in children with high blood pressure is limited. If the child can swallow tablets, the dose will be worked out using the child's weight and blood pressure.

The usual starting doses are:

- between 20 kg and 50 kg – 2.5 mg each day.
- more than 50 kg – 5 mg each day.

The dose can be changed according to the needs of the child:

- a maximum of 20 mg daily can be used in children who are between 20 kg and 50 kg.
- a maximum of 40 mg daily can be used in children who are more than 50kg.

This medicine is not recommended in new-born babies (first few weeks after birth) and in children with kidney problems.

If you take more Enalapril Tablets than you should

If you take more Enalapril Tablets than you should, talk to your doctor or go to a hospital straight away. Take the medicine pack with you. The following effects may happen: feeling of light-headed or dizziness. This is due to a sudden or excessive drop in blood pressure.

- **If you forget to take Enalapril Tablets** If you forget to take a tablet, skip the missed dose.
- - Take the next dose as usual.
- Do not take a double dose to make up for a forgotten dose.

If you stop taking Enalapril Tablets

Do not stop taking your medicine unless your doctor tells you to.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The following side effects may happen with this medicine:

Stop taking Enalapril Tablets and talk to a doctor straight away, if you notice any of the following:

- swelling of your face, lips, tongue or throat which may cause difficulty in breathing or swallowing
- swelling of your hands, feet or ankles
- if you develop a raised red skin rash (hives).

You should be aware that black patients are at increased risk of these types of reactions. If any of the above happen, stop taking Enalapril Tablets and talk to a doctor straight away.

When you start taking this medicine you may feel faint or dizzy. If this happens, it will help to lie down. This is caused by your blood pressure lowering. It should improve as you continue to take the medicine. If you are worried, please talk to your doctor.

Other side effects include:

Very Common (may affect more than 1 in 10 people)

- Feeling dizzy, weak or sick.
- blurred vision
- Cough.

Common (may affect up to 1 in 10 people)

- Light-headedness due to low blood pressure, changes in heart rhythm, fast heartbeat, angina or chest pain.
- Headache, depression, fainting (syncope),
- Change in sense of taste,
- Shortness of breath
- Diarrhoea, abdominal pain
- Tiredness (fatigue), depression,
- rash, allergic reactions with swelling of the face, lips, tongue or throat with difficulty in swallowing or breathing
- High levels of potassium in the blood, increased levels of creatinine in your blood (both are usually detected by a test).

Uncommon (may affect up to 1 in 100 people)

- flushing
- Sudden fall in blood pressure,
- Fast or uneven heart beats (palpitations),
- Anaemia (including aplastic and haemolytic),

- heart attack (possibly due to very low blood pressure in certain high-risk patients, include those with blood flow problems of the heart or brain)
- stroke (possibly due to very low blood pressure in high-risk patients)
- Confusion, sleeplessness or sleepiness, nervousness,
- feeling your skin prickling or being numb
- Pins and needles,
- Vertigo (spinning sensation)
- Ringing in your ears (tinnitus),
- Runny nose, sore throat or hoarseness,
- Asthma- associated tightness in chest
- Slow movement of food through your intestine (ileus), inflammation of your pancreas
- Being sick (vomiting), indigestion, constipation, anorexia,
- Irritated stomach (gastric irritations), dry mouth, ulcer,
- impaired kidney function, kidney failure
- increased sweating
- itching or nettle rash
- Increased perspiration,
- Loss of hair,
- Muscle cramps, generally feeling unwell (malaise), high temperature (fever), Impotence,
- High level of proteins in your urine (measured in a test),
- Low level of blood sugar or sodium, high level of blood urea (all measured in a blood test).

Rare (may affect up to 1 in 1,000 people)

- 'Raynaud's phenomenon' where your hands and feet may become very cold and white due to low blood flow
- Changes in blood values such as a lower number of white and red blood cells, lower haemoglobin, lower number of blood platelets
- Bone marrow depression
- swollen glands in neck, armpit or groin
- Autoimmune diseases
- Strange dreams or sleep problems
- accumulation of fluid or other substances in the lungs (as seen on X-rays)
- Inflammation of your nose
- inflammation of the lungs causing difficulty breathing (Pneumonia)
- Inflammation of the cheeks, gums, tongue, lips, throat
- Lower amount of urine produced
- rash that looks like targets (erythema multiforme)
- rash that looks like targets (Erythema multiforme)
- 'Stevens-Johnson syndrome' and 'toxic epidermal necrolysis' (serious skin conditions where you have reddening and scaling of your skin, blistering or raw scores), exfoliative dermatitis/erythroderma (severe skin rash with flaking or peeling of the skin), pemphigus (small fluid-filled bumps on the skin)
- liver or gallbladder problems such as lower liver function, inflammation of your liver, jaundice (yellowing of the skin or eyes), high levels of liver enzymes or bilirubin (measured in a blood test)
- Enlargement of the mammary glands in males (gynaecomastia).

Very Rare (may affect up to 1 in 10,000 people)

- Swelling in your intestine (intestinal angioedema).

Not known (frequency cannot be estimated from the available data)

- overproduction of antidiuretic hormone, which causes fluid retention, resulting in weakness, tiredness or confusion
- A symptom complex has been reported which may include some or all of the following: fever, inflammation of the blood vessels (serositis/vasculitis), muscle pain (myalgia/myositis), joint pain (arthralgia/arthritis). Rash, photosensitivity or other skin manifestations may occur.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme. Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow card in the Google play or Apple App Store By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Enalapril Tablets

- Do not store above 25°C. Store in the original package.
- **Keep this medicine out of the sight and reach of children.**
- Do not use this medicine after the expiry date which is stated on the label or carton. The expiry date refers to the last day of that month.
- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Enalapril Tablet contains

- The **active** substance is Enalapril maleate.
- The **other** ingredients are lactose, maize starch and glycerol distearate. (*see section 2 for Important information about some of the ingredients of Enalapril tablets*).

What Enalapril Tablets looks like and contents of the pack

- Enalapril 5 mg tablets are white, circular, biplaner, uncoated, with '5' embossed on one face and score line on the other.
- Enalapril 10 mg tablets are white, circular, biplaner, uncoated, with '10' embossed on one face and score line on the other.
- Enalapril 20 mg tablets are white, circular, biplaner, uncoated, with '20' embossed on one face and score line on the other.

Enalapril tablets are available in blister pack of 28 tablets.

Marketing Authorisation Holder and Manufacturer

Accord Healthcare Limited,

Sage House, 319 Pinner Road,
North Harrow, Middlesex,
HA1 4HF, United Kingdom

Enalapril 5mg Tablets (PL 20075/0279)
Enalapril 10mg Tablets (PL 20075/0280)
Enalapril 20mg Tablets (PL 20075/0281)

POM

This leaflet was last revised in 03/2018.