




LAMISIL CREAM 10MG/G TUBE 15G GB Artwork		karo healthcare
Reason for change:	Haleon to Karo artwork - MAH/logo/manuf.	Proof version 2
Correction date:	20-12-2023	Print version X
Artworker:	JC	
Artwork No:	2003473_1	
Prev Artwork component no:		
Haleon SKU No:	531608	
Component Type:	LEAFLET	
Pharma code:	1027	
Language:	English	
Manufacturing Site:	CH Nyon	
Approving Market:	Great Britain	
Product Market Name:	LAMISIL	
Site Component No:	62000000208851	
Prev Site Component No:	62000000075737	
Site Change Control No:	201282200	
Body Text Size:	8.0pt	
Leading:	9.6pt	
Horizontal Scale:	85%	
Smallest Text Size:	6.0pt	
Micro Text:	No	
Dimensions (WxHxD):	145 x 72 mm	
Machine Drawing ID:	N-PN P08 v2.1	
Material Spec No:	P08	
Print Process:	Offset lithography	
Print Colours: Colourmatch to Pantone Matching System.		
		
Technical (Non Printing).		
		
OVERPRINT PREVIEW <input type="checkbox"/> OFF		
<p>Karo Healthcare AB owns this artwork and is not responsible for any changes made by the CMO/printing house. Changes must be communicated with us and we will supply a new print PDF file. Both CMO and printing house are responsible for the printed end result to match the intended colours.</p>		

Recto



Terbinafine Hydrochloride

Read all of this leaflet carefully because it contains important information for you.

This medicine is available without prescription. However, you still need to use Lamisil AT 1% Cream carefully to get the best results from it.

- Please keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must contact a doctor if your symptoms worsen or do not improve.
- If any of the side effects get serious, or if you notice any side effect not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

- What Lamisil AT 1% Cream is and what it is used for
- Before you use Lamisil AT 1% Cream
- How to use Lamisil AT 1% Cream
- Possible side effects
- How to store Lamisil AT 1% Cream
- Further information.

1. WHAT LAMISIL AT 1% CREAM IS AND WHAT IT IS USED FOR

Lamisil AT 1% Cream contains the active ingredient terbinafine hydrochloride which is an antifungal used to treat fungal skin infections such as:

- Athlete's foot (tinea pedis)** – Athlete's foot appears only on the feet (usually both, but not always), usually in between the toes. It can also appear on the instep, soles, or other areas of the feet. The most common type of athlete's foot causes cracking or scaling of the skin, but you may also have mild swelling, blisters, or weeping ulcers. Lamisil AT 1% Cream is recommended for athlete's foot when it occurs between the toes. If you have a fungal nail infection (fungus inside and under the nail) with discolouration of the nails and changes in nail texture (thick or flaky), you should consult your doctor because Lamisil AT 1% Cream is not appropriate for this type of infection.
- Dhobie itch/jock itch (tinea cruris)** – Dhobie itch appears in the groin and inner thigh area, usually on both sides but often with one side more than the other, where it can spread further between the legs to the buttocks, or up towards the stomach. Dhobie itch looks like a clearly bordered rash, which may have blisters.

2. BEFORE YOU USE LAMISIL AT 1% CREAM

DO NOT use Lamisil AT 1% Cream if you:

- are allergic (hypersensitive) to any of the ingredients in the product (see Section 6).

The cream is NOT recommended for use on children under 16 years of age.

Take special care with Lamisil AT 1% cream

The cream is for external use only. Do not use on the face. Avoid contact with the eyes. In case of accidental contact with eyes, rinse thoroughly with running water.

Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including those obtained without a prescription.

Pregnancy and breast-feeding

Do not use the cream if you are pregnant or breastfeeding, unless advised to by your doctor.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Lamisil AT 1% Cream is not expected to affect your ability to drive or operate machinery.

Important information about some of the ingredients of Lamisil AT 1% Cream

- Lamisil AT 1% Cream contains 36 mg benzyl alcohol in each daily dosage which is equivalent to 10 mg/g. Benzyl alcohol may cause allergic reactions.
- Cetyl alcohol and stearyl alcohol: may cause local skin reactions (e.g. contact dermatitis).

3. HOW TO USE LAMISIL AT 1% CREAM

Always use this medicine exactly as described in this leaflet or as advised by your doctor or pharmacist.

Adults

For athlete's foot and dhobie itch, apply the cream once a day, for one week.

- Cleanse and dry the affected areas thoroughly and wash your hands. Treatment can be helped by keeping the affected areas clean by regular washing and careful drying with your own clean towels and clothes, and not rubbing or scratching the skin.
- Unscrew the cap then gently squeeze out a small amount of the cream onto your finger.
- Apply just enough cream to form a thin layer on the affected skin and surrounding areas.
- Rub in gently. When used between the toes, buttocks or on the groin, the treated area may be covered with a light, fresh gauze strip, especially at night.
- Replace the cap on the tube and wash your hands.

Even though you will not be using Lamisil AT 1% Cream during the second week, full skin healing after the

2003473_1/62000000208851

11.5

11.5

PHARMA CODE REF IS: 1027

72 mm

9

9

9

