Tums Assorted Fruit Antacid Tablets

Read all of the information carefully because it contains

Read all of the Information careturing because it commiss jumportant information for you. How to take Tums For oral use. Adults and children 12 years and over:
Suck or chew 1 to 2 tablets when needed, up to a maximum of 16 tablets (7 if you are pregnant) in 24 hours. Keep out of

children.

UT.

the sight and reach of

Do not store above 25°C.

Do not use after EXP date

Each tablet contains Calcium

Braunton, North Devon, EX33

2DL. U.K. The distributor is

PEEL HERE for more information about Tums Do not remove

Do **not** take this product within three hours of taking other medicines by mouth, as it can interfere with the action of some other medicines. Do not take Tums if you:

• have ever had an alleraic reaction to calcium carbonate or to any of the other

or can a low phosphate diet
have a high blood calcium level

 are taking heart tablets called
 cardiac glycosides (eg digoxin)
 have Zollinger-Ellison Syndrome (a tumour that secretes gastrin)

• are under 12 years. Ask your doctor before you take Tums if you:

• have been told by your doctor that you have an intolerance to some sugars

have diabetes ore taking other medicines, such as

tetracycline antibiotics, bisphosphonates (for your bones) or thiazide diuretics (water tablets).

Do not give to children under 12 years. Do not take more than the recommended dose. If symptoms persist, see your doctor.

If you take too many tablets Contact your doctor or casualty department. .37mm

90.

Possible side effects

You may aet constipation, wind or changes in your blood while using Tums. You may get neartburn after you stop taking the tablets. Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/vellowcard By reporting side effects you can help provide more information on the safety of this medicine. Active ingredient Each tablet contains Calcium Carbonate 500 ma.

Other ingredients Each tablet also contains dextrose, maize starch, microcrystalline cellulose, magnesium stearate and sodium saccharin, Flavours; blackcurrant, cherry, lemon i and orange, Colours; patent blue V (E 131). ponceau 4R (E 124), quinoline yellow (E 104) and sunset yellow (E 110).

Colours E 110 & E 124 can cause alleraic reactions.

Each tablet contains between 521 and 525 ma of alucose (dextrose). Information prepared: March 2015

Tums is a registered trade mark of the GSK aroup of companies.

L367030/03 6109833i

69.85mm