
PLEASE CHECK ALL ARTWORK AND SPECIFICATION DETAILS BEFORE SIGNING

BY SIGNING THIS APPROVAL YOU ARE AUTHORISING US TO PRINT USING THESE DETAILS. PRODUCTION WILL NOT COMMENCE UNTIL APPROVAL OF PROOFS AND SPECIFICATION DETAILS ARE IN PLACE

	ARTWORK DETAILS		SPECIFICATION DETAILS	
We ask you to check fully and sign below.	Customer	Dendron Brands Ltd	Previous Version	N/A
Every care is taken to ensure correctness. We cannot be held responsible for	Product Name	LFT DENTINOX RELAUNCH 2016	Pharmacode No.	P0507
mistakes if passed by you.		TEETHING GEL	Material	50gsm
Please mark clearly any changes you	Item Code	LTD0181	Flat Size	165 (27.5) x 120
require and return the original visual to us. (Colours on the visual are representative and not exact colour matches).	Our Ref	SO55385	Fold Size	27.5 x 120
	Proof No.	Two	View Panel	DATAMATRIX CODE & TITLE TO VIEW
We will re-proof by return unless instructed otherwise. Major alterations	Date	11/09/2020	Fold Detail	PARALLEL
from the original copy are chargeable as 'Author's Corrections'.	ARTWORK APPROVED		Packing Details	Shrink Wrap in Trays
Author's Corrections .			Colours Front	CYAN, PMS 3285
CLO64/ISSUE 8 AMENDMENTS R		EQUIRED	Colours Back	PMS 3285
	ication and Artwork approved for printing by			
Tel: 01638 660063 Fax: 01638 667128 ISDN: 01638 560030	T			



3. HOW YOU USE DENTINOX TEETHING GEL DOSAGE SIZE Apply a pea sized blob of gel (see circle shown above), to a clean fingertip and spread gently onto the sore area of the gum. If necessary, repeat the dose after 3 hours. Do not use more than 6 times in one day (24 hour period). Stop treatment when your child's symptoms settle. Do not use for more than 7 days for each teething episode. If your child's symptoms worsen, do not improve after 7 days of treatment or you are concerned that your child is unwell, talk to a health care professional. If your child vomits, spits or swallows the medicine, do not use any more product immediately. Wait 3 hours before using the medicine again. Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you to. Do not give more medicine than the label tells you to. You should check with your doctor or pharmacist if you are not sure. 4. POSSIBLE SIDE EFFECTS There are no known side effects from Dentinox Teething Gel. Reporting of side effects. If your baby experiences any side effects, consult your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine. 5. HOW TO STORE DENTINOX TEETHING GEL Dentinox Teething Gel should be kept out of the reach and sight of children. Do not store above 25°C. Do not use Dentinox Teething Gel after the expiry date as indicated on the box. The expiry date refers to the last day of the month. Medicine should not be disposed of in waste-water or with household waste. Ask your pharmacist what you should do with medications you no longer need. These measures will help to protect the environment. 6. FURTHER INFORMATION What Dentinox Teething Gel contains: The active substances are lidocaine hydrochloride 0.33% w/w and cetylpyridinium chloride 0.10% w/w. The other ingredients are: Sorbitol solution 70% (non-crystallising), xylitol, ethanol, glycerol, hydroxethylcellulose, polyoxyl 40 hydrogenated castor oils, pharmaceutical liquid flavour, hydroxypolyethoxydodecane, macrogol 300, sodium saccharin, caramel (E150), levomenthol and purified water. What Dentinox Teething Gel looks like and contents of the pack: Dentinox Teething Gel is available in aluminium tubes containing 10g. Marketing Authorisation Holder and Manufacturer: Dendron Brands Limited, 94 Rickmansworth Road, Watford, Hertfordshire, United Kingdom, WD18 7JJ. This leaflet was last revised in September 2020.