

Torem 5 mg Tablets

Torem 10 mg Tablets

Torasemide

Please read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them even if their symptoms are the same as yours.
- If any of the side effects gets serious or troublesome, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Torem is and what it is used for
2. Before you take Torem
3. How to take Torem
4. Possible side effects
5. How to store Torem
6. Further information

1. What Torem is and what it is used for

Torem contains a medicine called torasemide. This belongs to a group of medicines called 'diuretics' or 'water tablets'.

- Torem 5 mg tablets are used to treat high blood pressure (hypertension).
- Torem 5 mg and 10 mg tablets are used to treat swelling caused by too much water in the body (oedema).

The tablets work by making you pass more water (urine).

2. Before you take Torem

Do not take Torem if you are allergic (hypersensitive) to:

- Torasemide or any of the other ingredients of Torem (see Section 6: Further information).
- Similar medicines called 'sulphonylureas'. These are used to treat diabetes (high blood sugar). These medicines include chlorpropamide, glibenclamide, glipizide and tolbutamide.

Do not take Torem if any of these apply to you. If you are not sure, talk to your doctor or pharmacist before taking Torem.

Do not take Torem if:

- You have kidney problems (renal failure).
- You have kidney problems that have been caused by medicines.
- You have liver problems.
- You have a low volume of blood (hypovolaemia).
- You have low blood pressure (hypotension).
- You have heart beat problems (cardiac arrhythmia).
- You are taking an antibiotic called an 'aminoglycoside' or 'cephalosporin' (for an infection). These medicines include streptomycin, gentamycin, cephalixin and ceftriaxone. See the section on 'Taking other medicines'.
- You are pregnant or breast-feeding a baby. See the section on 'Pregnancy and breast-feeding'.

Do not take Torem if any of these apply to you. If you are not sure, talk to your doctor or pharmacist before taking Torem.

Take special care with Torem

Check with your doctor or pharmacist before taking Torem if:

- You have low levels of potassium or sodium in your blood (shown in blood tests).
- You have difficulty with micturition (passing urine) including prostatic hypertrophy.
- You have diabetes (diabetes mellitus).

If any of these apply to you, or if you are not sure, talk to your doctor or pharmacist before you take Torem.

Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes medicines that you buy without a prescription and herbal medicines.

This is because Torem can affect the way some other medicines work.

Also some other medicines can affect the way Torem works.

Do not take Torem if you are already taking an antibiotic called an 'aminoglycoside' or 'cephalosporin' (for an infection). These medicines include streptomycin, gentamycin, cephalixin and ceftriaxone. If this applies to you, ask your doctor or pharmacist for advice.

It is particularly important to **tell your doctor or pharmacist** if you are taking any of the following medicines:

- Other medicines to treat high blood pressure.
- Medicines called 'ACE inhibitors' (used to treat heart disease).
- Medicines like digoxin or digitoxin (used to treat heart disease).
- Adrenaline (also called epinephrine) or noradrenaline (also called norepinephrine). These are used to treat low blood pressure.
- Cholestyramine, or other 'ionexchange resins' (used to treat high levels of cholesterol in your blood).
- High doses of medicines called 'salicylates', such as aspirin.
- Medicines called 'non-steroidal antiinflammatory drugs' (NSAIDs). These include indomethacin and ibuprofen.
- Laxatives.
- Steroid medicines, such as hydrocortisone, prednisolone and dexamethasone.
- Medicines to relax your muscles (muscle relaxants).
- Cisplatin (used to treat cancer).
- Lithium (used to treat mental health problems).
- Probenecid (used to treat gout).
- Theophylline (used to treat asthma).

Pregnancy and breast-feeding

Do not take Torem if you are pregnant, might become pregnant or are breast-feeding. This is because it is not known if Torem will affect your baby.

Driving and using machines

You may feel dizzy while taking Torem. If this happens, do not drive or use any tools or machines.

Important information about some of the ingredients of Torem

This medicine contains lactose, which is a type of sugar. If you have been told by your doctor that you cannot tolerate or digest some sugars (have an intolerance to some sugars), talk to your doctor before taking this medicine.

3. How to take Torem

Always take Torem exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

High blood pressure

- The usual dose for adults and elderly people is between 2.5 mg and 5 mg once a day.

Water retention (oedema)


- The usual dose for adults and elderly people is 5 mg once a day.
- Your doctor may increase the dose to 20 mg a day if needed.

If you take more Torem than you should

- If you take more Torem than you should, talk to a doctor or go to a hospital straight away. Take the medicine pack with you.
- If you take too many tablets, you may pass a lot of urine, become sleepy, or feel confused, faint or dizzy. You may also have stomach upset.

If you forget to take Torem

- If you forget to take a dose, take it as soon as you remember it.
- However, if it is nearly time for the next dose, skip the missed dose.

SPECIFICATION BOX	Manufacturer	Meda Pharma GmbH	Used template	N/A	Fonts used	Times New Roman
	Product (+Pack Size)	UK1956228 Torem 5 + 10 mg tablets, 28 tabs	Code (PZN, EAN, Datamatrix)	N/A	Min. font-size text	Frutiger LT Std 9,5 pt
 Typopharma GmbH Mühlstr. 57 D-79539 Leirach produktion@typopharma.com www.typopharma.com typopharma no: # 131928	Code / no.	56UK1956228-00	Comments	N/A	Dieses Korrektur-PDF wurde von uns mit größter Sorgfalt erstellt und einer internen Qualitätssicherungsprüfung unterzogen. Dennoch bedeutet jeder Eingriff in die Datei auch ein Risiko. Wir bitten deshalb, den vorliegenden Korrekturabzug eingehend und umfänglich zu prüfen. Für Fehler, die erst nach einer Freigabe festgestellt werden, übernimmt die Typopharma GmbH keine Haftung.	
	Replaced no.	T5008914	Colours used	■ black		
	Order no.	705886	Technical colours	Cutting		
	Country/language	UK, english	Corrections / No. of proofs	4		
	Component Size	148 x 420 mm	1	23.08.2018 sm		
	Page no.	Page 1 of 2	2	Correction required		
	Drawing no.	WZ_209_2	3			
	Laetus no. SEM	101	7			

56UK1956228-00
705886


- Do not take a double dose to make up for a forgotten dose.

If someone else takes your Torem tablets by mistake, they should talk to a doctor or go to a hospital straight away.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, Torem can cause side effects, although not everyone will get them.

Within the system organ classes, adverse reactions are listed under headings of frequency (number of patients expected to experience the reaction), using the following categories:

Very common ($\geq 1/10$)

Common ($\geq 1/100$ to $< 1/10$)

Uncommon ($\geq 1/1,000$ to $< 1/100$)

Rare ($\geq 1/10,000$ to $< 1/1,000$)

Very rare ($< 1/10,000$)

Not known (cannot be estimated from available data)

The following undesirable effects were observed whereas the frequency of undesirable effect is not known:

See your doctor as soon as possible if you get the following side effects:

Blood and lymphatic system disorders
Frequency not known: Thrombocytopenia, Leukopenia, Anaemia

Immune system disorders
Very rare: Allergic skin reactions (e.g. Pruritus, Exanthema), Photosensitivity reaction

Frequency not known: Serious skin reactions (e.g. Stevens-Johnson syndrome, Toxic epidermal necrolysis)

Metabolism and nutrition disorders
Common: Metabolic alkalosis, Fluid and electrolyte imbalance (e.g. Hypovolaemia, Hyponatraemia)

Nervous system disorders
Common: Headache, Dizziness
Frequency not known: Cerebral ischaemia, Parosmia, Confusional state

Eye disorders
Frequency not known: Visual impairment

Ear and labyrinth disorders
Frequency not known: Tinnitus, Deafness

Cardiac disorders
Frequency not known: Acute myocardial infarction, Myocardial ischaemia, Angina pectoris, Syncope, Hypotension

Vascular disorders
Frequency not known: Embolism

Gastrointestinal disorders
Common: Gastrointestinal disorder (e.g. Loss of appetite, Abdominal pain upper, Nausea, Vomiting, Diarrhoea, Constipation)
Frequency not known: Dry mouth, Pancreatitis

Hepatobiliary disorders
Uncommon: Hepatic enzyme increased (e.g. Gamma-glutamyltransferase increased)

Skin and subcutaneous tissue disorders
Very rare: Allergic skin reactions (e.g. Pruritus, Exanthema), Photosensitivity reaction
Frequency not known: Serious skin reactions (e.g. Stevens-Johnson syndrome, Toxic epidermal necrolysis)

Musculoskeletal and connective tissue disorders
Common: Muscle spasms

Renal and urinary disorders
Uncommon: Urinary retention, Bladder dilatation
Rare: Blood urea increased, Blood creatinine increased

General disorders and administration site conditions
Common: Fatigue, Asthenia

Investigations
Uncommon: Blood uric acid increased, Blood glucose increased, Lipids increased (e.g. Blood triglycerides increased, Blood cholesterol increased)

Other possible side effects:

Blood

- An increase in the levels of certain substances in your blood, including sugar, uric acid and fats (lipids).

Your doctor may do blood tests from time to time to make sure that Torem is working safely.

Stomach and gut

- Stomach upset.

- Inflammation of the pancreas causing pain in the stomach (pancreatitis).
- Changes in how your liver is working (shown by blood tests).

Skin

- Skin rashes and itchy skin.
- Skin being more sensitive to the sun (photo-sensitivity).

Eyes and ears

- Blurred eyesight.
- Ringing in the ears (tinnitus) and loss of hearing.

Other

- Feeling numb or tingling in the legs or arms.
- Dry mouth.
- Some people may have difficulty passing water (urinating).

If any of the side effects become serious or troublesome, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

5. How to store Torem

- Torem tablets do not need special storage conditions.
- Keep out of the reach and sight of children.
- Do not use Torem after the expiry date printed on the pack.
- Do not throw away any left over tablets. Instead, return them to your pharmacist so that they can be disposed of carefully. Only keep them if your doctor tells you to.

6. Further information

What Torem contains

The active substance in Torem is torasemide. There are two different strengths of tablet:

- Torem 5 mg Tablets (each tablet contains 5 milligrams of torasemide).
- Torem 10 mg Tablets (each tablet contains 10 milligrams of torasemide).

Other ingredients are lactose monohydrate, maize starch, colloidal silicon dioxide and magnesium stearate (E572).

What Torem looks like and contents of the pack

Torem 5 mg tablets are round. They are white to off-white round tablets with the imprint "T 5.0" and break mark on one side and plain on the other side. Torem 10 mg tablets are round. They are white to off-white round tablets with the imprint "T 10.0" and break mark on one side and plain on the other side. Torem tablets are supplied in blister packs of 14, 28, 100 or 112 tablets. Not all pack sizes may be marketed.

The Marketing Authorisation holder is:

Mylan Products Ltd.
Station Close, Potters Bar,
Hertfordshire,
EN6 1TL, United Kingdom


The manufacturers responsible for batch release is:

MEDA Pharma GmbH & Co. KG
Benzstr. 1
61352 Bad Homburg, Germany

MEDA Manufacturing GmbH
Neurather Ring 1
51063 Köln, Germany

Madaus GmbH
51101 Köln, Germany

This leaflet was last revised in August 2018.

SPECIFICATION BOX		Manufacturer	Meda Pharma GmbH	Used template	N/A
 Typopharma GmbH Mühlstr. 57 D-79539 Leirach produktion@typopharma.com www.typopharma.com typopharma no: # 131928		Product (+Pack Size)	Torem 5 + 10 mg tablets, 28 tabs	Code (PZN, EAN, Datamatrix)	N/A
		Code / no.	56UK1956228-00	Comments	N/A
		Replaced no.	T5008914	Colours used	■ black
		Order no.	705886	Technical colours	Cutting
		Country/language	UK, english	Corrections / No. of proofs	4
		Component Size	148 x 420 mm	1	23.08.2018 sm
		Page no.	Page 2 of 2	2	
		Drawing no.	WZ_209_2	3	
		Laetus no. SEM	101 N/A		

Times New Roman
Frutiger LT Std
9,5 pt

Dieses Korrektur-PDF wurde von uns mit größter Sorgfalt erstellt und einer internen Qualitätssicherungsprüfung unterzogen. Dennoch bedeutet jeder Eingriff in die Datei auch ein Risiko. Wir bitten deshalb, den vorliegenden Korrekturabzug eingehend und umfänglich zu prüfen. Für Fehler, die erst nach einer Freigabe festgestellt werden, übernimmt die Typopharma GmbH keine Haftung.

Release
 Correction required

! PLEASE TURN OVERPRINTING ON !