


**PRINTED PACKAGING MATERIAL MASTER**

Product Code: <b>28063194</b>	Product Description: Proveblue 5 X 10 ml UK - Leaflet	
Dimensions : 210 mm x 315 mm 2 pages	Technical info : Packaging line : FB320	1 COLOR  BLACK CMYK (Black 100%)
PROVEPHARM		
Reason for change : Deletion of IE PV contact + Update of UK distributor + update of revision date + Insertion of UK Marketing authorisation number	Version 2 Date : 20.Dec.2021	

Package leaflet Information for the user

**Methylthioninium chloride Proveblue 5 mg/ml  
solution for injection**

Methylthioninium chloride

**Read all of this leaflet carefully before you are given this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or nurse.
- If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What Methylthioninium chloride Proveblue is and what it is used for
2. What you need to know before you are given Methylthioninium chloride Proveblue
3. How Methylthioninium chloride Proveblue is given
4. Possible side effects
5. How to store Methylthioninium chloride Proveblue
6. Contents of the pack and other information

**1. What Methylthioninium chloride Proveblue is and what it is used for**

Methylthioninium chloride (also called methylene blue) belongs to a group of medicines called antidotes.

Methylthioninium chloride Proveblue will be given to you or your child (0-17 years old) to treat problems with your blood resulting from exposure to some medicines or chemicals that can cause a disease called methaemoglobinaemia.

In methaemoglobinaemia, your blood contains too much methaemoglobin (an abnormal form of haemoglobin that is not able to transport oxygen around your body effectively). This medicine will help your haemoglobin return to normal and restore the transport of oxygen in the blood.

**2. What you need to know before you are given Methylthioninium chloride Proveblue**

**You must not be given Methylthioninium chloride Proveblue:**

- if you are allergic to methylthioninium chloride or other thiazine dyes
- if your body does not produce enough of the enzyme G6PD (glucose-6-phosphate dehydrogenase)
- if your body does not produce enough of the enzyme NADPH (nicotinamide adenine dinucleotide phosphate) reductase
- if your blood disorder has been caused by nitrite during treatment of cyanide poisoning
- if your blood disorder has been caused by chlorate poisoning.

**Warnings and precautions**

Talk to your doctor or nurse before you are given Methylthioninium chloride Proveblue

- if you have moderate or severe renal disease; lower doses (< 1 mg/kg) may be needed
- if your blood disorder has been caused by a chemical called aniline, which is contained in dyes; lower doses may be needed and total cumulative dose should not exceed 4 mg/kg (see section 3 of this package leaflet)
- if your blood disorder has been caused by a medicine called dapson (used to treat leprosy and other skin conditions); lower doses may be needed and total cumulative dose should not exceed 4 mg/kg (see section 3)
- if you suffer from hyperglycaemia or diabetes mellitus, as these conditions may be worsened by the glucose solution used for the dilution of the medicine
- your urine and stools may turn a blue-green colour; and skin may possibly turn a blue colour when you are treated with Methylthioninium chloride Proveblue. This discolouration is expected and will disappear after the treatment has ended

If any of the above applies to you, please consult your doctor.

**Photosensitivity**

Methylthioninium chloride may cause a photosensitivity reaction in the skin (sunburn-like reaction) when exposed to strong light sources, such as light therapy, lights in operating rooms and pulse oximeters.

Protective measures against exposure to light should be taken.

**Monitoring tests**

You will undergo monitoring tests during and after treatment with Methylthioninium chloride Proveblue.

**Children**

**Special care must be taken with Methylthioninium chloride Proveblue:**

- in newborns and infants 3 months old or younger, lower doses are recommended (see section 3 of this package leaflet).

**Other medicines and Methylthioninium chloride Proveblue**

Tell your doctor, pharmacist or nurse if you are taking, have recently taken or might take any other medicines.

You should not be given methylthioninium chloride at the same time you are taking certain medicines to treat depression or anxiety which affect a brain chemical called serotonin. When used in combination with these medicines methylthioninium chloride may cause serotonin syndrome, which can be potentially life-threatening. Such medicines include:

- Selective serotonin reuptake inhibitors (SSRIs) such as citalopram, escitalopram, fluoxetine, fluvoxamine, paroxetine, sertraline and zimelidine
- bupropion
- buspirone
- clomipramine
- mirtazapine
- venlafaxine
- Monoamine oxidase inhibitors

However, if the intravenous use of methylthioninium chloride cannot be avoided, you should be administered the lowest possible dose and observed closely for up to 4 hours after administration.

If you have any doubts about whether this medicine should be given to you, consult your doctor.

**Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before you are given this medicine.

The use of Methylthioninium chloride Proveblue during pregnancy is not recommended unless it is clearly necessary, for example in a life-threatening situation.

Due to a lack of available data on whether methylthioninium chloride passes into human breast milk, breast-feeding should be discontinued for up to 8 days after treatment with this medicine.

**Driving and using machines**

Do not drive or use any tools or machines as methylthioninium chloride has moderate influence on the ability to drive and use machines.

**3. How Methylthioninium chloride Proveblue is given**

Your doctor will inject this medicine into a vein (intravenously) slowly over a period of 5 minutes.

**Adults, children above 3 months and elderly**

The usual dose is 1 to 2 mg per kilogram of your body weight, i.e. 0.2 to 0.4 ml per kilogram given over a period of 5 minutes. A second dose may be given after one hour if required.

The maximum recommended cumulative dose for the course of treatment is 7 mg/kg.

If your blood disorder has been caused by aniline or dapson, total cumulative dose should not exceed 4 mg/kg (see section 2).

Usually, treatment should not exceed one day.

**Infants 3 months old or younger**


The recommended dose is 0.3 to 0.5 mg/kg body weight, i.e. 0.06 to 0.1 ml/kg, over a period of 5 minutes.

A repeat dose (0.3 to 0.5 mg/kg body weight, i.e. 0.06-0.1 ml/kg) may be given after one hour in case of persistence or recurrence of symptoms. Usually, treatment should not exceed one day.

This medicine may be diluted in 50 ml glucose 50 mg/ml (5%) solution for injection to avoid local pain, in particular in children.



28063194

PRINTED PACKAGING MATERIAL MASTER		
Product Code: <b>28063194</b>	Product Description: Proveblue 5 X 10 ml UK - Leaflet	
Dimensions : 210 mm x 315 mm 2 pages	Technical info : Packaging line : FB320	1 COLOR  BLACK CMYK (Black 100%)
PROVEPHARM		
Reason for change : Deletion of IE PV contact + Update of UK distributor + update of revision date + Insertion of UK Marketing authorisation number	Version 2 Date : 20.Dec.2021	

#### If you are given more Methylthionium chloride Proveblue than you should

As this medicine will be given to you whilst you are in hospital, it is unlikely that you will be given too much or too little, however, tell your doctor if you notice one of the following adverse reactions:

- feeling sick,
- stomach pain,
- chest pain,
- dizziness,
- headache,
- sweating,
- confusion,
- an increase in methaemoglobin (an abnormal form of haemoglobin in the blood),
- high blood pressure,
- shortness of breath,
- abnormally fast beating of the heart,
- tremor,
- skin discolouration. Your skin may turn blue
- reduction in red blood cells which may turn your skin pale and make you breathless and weak,
- jaundice (yellowing of the skin and eyes), this has only been reported in infants.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

#### 4. Possible side effects

Like all medicines, Methylthionium chloride Proveblue can cause side effects, although not everybody gets them.

These effects are the same in adults and children except jaundice which has only been reported in infants.

- **Very common side effects** (may affect more than 1 in 10 people)

- pain in extremity
- dizziness
- sweating
- skin discolouration. Your skin may turn blue
- blue or green urine
- numbness and tingling
- abnormal taste in mouth
- nausea

- **Common side effects** (may affect up to 1 in 10 people):

- stomach pain
- chest pain
- headache
- anxiety
- injection site pain
- vomiting

- **Not known** (frequency cannot be estimated from the available data):

- serotonin syndrome when Methylthionium chloride Proveblue has been taken with certain medicines to treat depression or anxiety, see section 2
- decreased haemoglobin (protein in red blood cells that carry oxygen in the blood) levels may be reported during blood tests
- reduction in red blood cells which may turn your skin pale and make you breathless and weak
- local tissue damage at the injection site
- jaundice (yellowing of the skin and eyes) – this has only been reported in infants
- problems with speech
- high or low blood pressure
- agitation
- lack of oxygen
- irregular heartbeat, including an abnormally slow or fast beating of the heart
- severe allergic reactions (so called anaphylactic reaction which may cause your throat or face to swell, difficulty breathing or a severe rash)
- an increase in methaemoglobin (an abnormal form of haemoglobin in the blood)
- shortness of breath
- confusion
- shaking

- hives
- fever
- rapid breathing
- dilated pupils
- discoloured stools. They may appear green or blue
- increased sensitivity of your skin to light (photosensitivity)

#### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (See details below). By reporting side effects you can help provide more information on the safety of this medicine.

#### United Kingdom

Yellow Card Scheme  
Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard)  
or search for MHRA Yellow Card in the Google Play or Apple App Store

#### 5. How to store Methylthionium chloride Proveblue

Keep this medicine out of the sight and reach of children.

You should not be given this medicine after the expiry date which is printed on the carton and the ampoule labels after EXP. The expiry date refers to the last day of that month. The doctor or nurse will check that the expiry date on the label has not been passed before administering the injection to you.

Do not refrigerate or freeze. Keep the ampoule in the original package in order to protect from light.

The medicine must be used immediately after opening or dilution.

Do not use Methylthionium chloride Proveblue if the solution is discoloured, cloudy, turbid, or a precipitate or particles are present.  
Any unused product or waste material should be disposed of in accordance with local requirements.

#### 6. Contents of the pack and other information

##### What Methylthionium chloride Proveblue contains

- The active substance is methylthionium chloride.
- Each ml of solution contains 5 mg methylthionium chloride.
- Each 10 ml ampoule contains 50 mg methylthionium chloride.
- The other ingredient is water for injections.

##### What Methylthionium chloride Proveblue looks like and contents of the pack

Methylthionium chloride Proveblue is a clear dark blue solution for injection (injection) and is supplied in clear glass ampoules.  
Each box contains a tray with 5 ampoules of 10 ml.

##### Marketing Authorisation Holder

Provepharm SAS  
22 rue Marc Donadille, 13013 Marseille, France

##### Manufacturer responsible for batch release

Cenexi  
52, Rue Marcel et Jacques Gaucher, 94120 Fontenay-sous-Bois, France

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

##### United Kingdom

Macarthy's Laboratories Limited  
T/A Martindale Pharma  
Romford, Essex, RM3 8UG, UK  
Tel: + 44 (0)1277 266600

##### This leaflet was last revised in 12/2021

Marketing Authorisation number : PLGB 40051/0002

The following information is intended for healthcare professionals only:

#### Preparation for intravenous administration

Use immediately on opening. Inject very slowly over a period of 5 minutes.

Methylthionium chloride Proveblue is hypotonic and may be diluted in 50 ml glucose 50 mg/ml (5%) solution for injection to avoid local pain, in particular in paediatric population.

It must not be diluted with sodium chloride 9 mg/ml (0.9%) solution for injection because it has been demonstrated that chloride reduces the solubility of methylthionium chloride.

Additional information on how Methylthionium chloride Proveblue can be given is provided in section 3 of the Package Leaflet.

Any unused product or waste material should be disposed of in accordance with local requirements.



28063194

28063194