

Package leaflet: Information for the user

**Atorvastatin 10 mg film-coated tablets**  
**Atorvastatin 20 mg film-coated tablets**  
**Atorvastatin 30 mg film-coated tablets**  
**Atorvastatin 40 mg film-coated tablets**  
**Atorvastatin 60 mg film-coated tablets**  
**Atorvastatin 80 mg film-coated tablets**

Atorvastatin

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

**What is in this leaflet**

1. What Atorvastatin is and what it is used for
2. What you need to know before you take Atorvastatin
3. How to take Atorvastatin
4. Possible side effects
5. How to store Atorvastatin
6. Contents of the pack and other information

**1. What Atorvastatin is and what it is used for**

Atorvastatin belongs to a group of medicines known as statins, which are lipid (fat) regulating medicines.

Atorvastatin is used to lower lipids known as cholesterol and triglycerides in the blood when a low fat diet and life style changes on their own have failed. If you are at an increased risk of heart disease, Atorvastatin can also be used to reduce such risk even if your cholesterol levels are normal. You should maintain a standard cholesterol lowering diet during treatment.

**2. What you need to know before you take Atorvastatin**

**Do not take Atorvastatin**

- if you are allergic to atorvastatin or any of the other ingredients of this medicine (listed in section 6)
- if you have or have ever had a disease which affects the liver
- if you have had any unexplained abnormal blood tests for liver function
- if you are a woman able to have children and not using reliable contraception
- if you are pregnant or trying to become pregnant
- if you are breast-feeding.
- If you use the combination of glecaprevir/pibrentasvir in the treatment of hepatitis C

**Warnings and precautions**

Talk to your doctor, pharmacist or nurse before taking Atorvastatin.

- if you have severe respiratory failure
- if you are taking or have taken in the last 7 days a medicine called fusidic acid, (a medicine for bacterial infection) orally or by injection. The combination of fusidic acid and Atorvastatin can lead to serious muscle problems (rhabdomyolysis)
- if you have had a previous stroke with bleeding into the brain, or have small pockets of fluid in the brain from previous strokes
- if you have kidney problems
- if you have an under-active thyroid gland (hypothyroidism)
- if you have had repeated or unexplained muscle aches or pains, a personal history or family history of muscle problems
- if you have had previous muscular problems during treatment with other lipid-lowering medicines (e.g. other '-statin' or '-fibrate' medicines)
- if you regularly drink a large amount of alcohol
- if you have a history of liver disease
- if you are older than 70 years.

If any of these apply to you, your doctor will need to carry out a blood test before and possibly during your Atorvastatin treatment to predict your risk of muscle related side effects. The risk of muscle related side effects e.g rhabdomyolysis is known to increase when certain medicines are taken at the same time (see Section 2 "Other medicines and Atorvastatin").

Also tell your doctor or pharmacist if you have a muscle weakness that is constant. Additional tests and medicines may be needed to diagnose and treat this.

While you are on this medicine your doctor will monitor you closely if you have diabetes or are at risk of developing diabetes. You are likely to be at risk of developing diabetes if you have high levels of sugars and fats in your blood, are overweight and have high blood pressure.

**Other medicines and Atorvastatin**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

There are some medicines that may change the effect of Atorvastatin or their effect may be changed by Atorvastatin. This type of interaction could make one or both of the medicines less effective. Alternatively it could increase the risk or severity of side-effects, including the important muscle wasting condition known as rhabdomyolysis described in Section 4:

- Medicines used to alter the way your immune system works, e.g. ciclosporin
- Certain antibiotics or antifungal medicines, e.g. erythromycin, clarithromycin, telithromycin, ketoconazole, itraconazole, voriconazole, fluconazole, posaconazole, rifampin, fusidic acid
- Other medicines to regulate lipid levels, e.g. gemfibrozil, other fibrates, colestipol

- Some calcium channel blockers used for angina or high blood pressure, e.g. amlodipine, diltiazem,; medicines to regulate your heart rhythm e.g. digoxin, verapamil, amiodarone
- Medicines used in the treatment of HIV e.g. ritonavir, lopinavir, atazanavir, indinavir, darunavir, the combination of tipranavir/ritonavir etc.
- Some medicines used in the treatment of hepatitis C e.g. telaprevir, boceprevir and the combination of elbasvir/grazoprevir
- Other medicines known to interact with Atorvastatin include ezetimibe (which lowers cholesterol), warfarin (which reduces blood clotting), oral contraceptives, stiripentol (an anti-convulsant for epilepsy), cimetidine (used for heartburn and peptic ulcers), phenazone (a painkiller), colchicine (used to treat gout) and antacids (indigestion products containing aluminium or magnesium)
- Medicines obtained without a prescription: St John's Wort
- If you need to take oral fusidic acid to treat a bacterial infection you will need to temporarily stop using this medicine. Your doctor will tell you when it is safe to restart Atorvastatin. Taking Atorvastatin with fusidic acid may rarely lead to muscle weakness, tenderness or pain (rhabdomyolysis). See more information regarding rhabdomyolysis in section 4

Tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

**Atorvastatin with food, drink and alcohol**

See Section 3 for instructions on how to take Atorvastatin. Please note the following:

*Grapefruit juice*

Do not take more than one or two small glasses of grapefruit juice per day because large quantities of grapefruit juice can change the effects of Atorvastatin.

*Alcohol*

Avoid drinking too much alcohol while taking this medicine. See Section 2 "Warnings and precautions" for details

**Pregnancy, breast-feeding and fertility**

Do not take Atorvastatin if you are able to become pregnant unless you use reliable contraceptive measures. The safety of Atorvastatin during pregnancy and breast-feeding has not yet been proven. Ask your doctor or pharmacist for advice before taking any medicine.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine

**Driving and using machines**

Normally this medicine does not affect your ability to drive or operate machines. However, do not drive if this medicine affects your ability to drive. Do not use any tools or machines if your ability to use them is affected by this medicine.

**Atorvastatin contains lactose monohydrate and soya lecithin.**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

This medicine contains soya lecithin. If you are allergic to peanut or soya, do not take this medicinal product.

*Sodium*

Atorvastatin contains less than 1 mmol (23 mg) of sodium per tablet, that is to say it is essentially 'sodium-free.'

**3. How to take Atorvastatin**

Before starting treatment, your doctor will place you on a low-cholesterol diet, which you should maintain also during therapy with Atorvastatin.

The recommended starting dose of Atorvastatin is 10 mg once a day in adults and children aged 10 years or older. This may be increased if necessary by your doctor until you are taking the amount you need. Your doctor will adapt the dose at intervals of 4 weeks or more. The maximum dose of Atorvastatin is 80 mg once a day.

Atorvastatin tablets should be swallowed whole with a drink of water, and can be taken at any time of day, with or without food. However, try to take your tablet at the same time every day.

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

**The duration of treatment with Atorvastatin is determined by your doctor.**

Please ask your doctor if you think that the effect of Atorvastatin is too strong or too weak.

**If you take more Atorvastatin than you should**

If you accidentally take too many Atorvastatin tablets (more than your usual daily dose), contact your doctor or nearest hospital for advice.

**If you forget to take Atorvastatin**

If you forget to take a dose, just take your next scheduled dose at the correct time. Do not take a double dose to make up for a forgotten dose.

**If you stop taking Atorvastatin**

If you have any further questions on the use of this medicine or wish to stop your treatment, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**If you experience any of the following serious side effects, stop taking your tablets and tell your doctor immediately or go to the nearest hospital accident and emergency department.**

**Rare** (may affect up to 1 in 1,000 people):

- Serious allergic reaction which causes swelling of the face, tongue and throat that can cause great difficulty in breathing.
- Serious illness with severe peeling and swelling of the skin, blistering of the skin, mouth, eyes genitals and fever. Skin rash with pink-red blotches especially on palms of hands or soles of feet which may blister.
- Muscle weakness, tenderness or pain and particularly, if at the same time, you feel unwell or have a high temperature it may be caused by an abnormal muscle breakdown (rhabdomyolysis). The abnormal muscle breakdown does not always go away, even after you have stopped taking atorvastatin, and it can be life-threatening and lead to kidney problems.

**Very rare** (may affect up to 1 in 10,000 people):

- If you experience problems with unexpected or unusual bleeding or bruising, this may be suggestive of a liver complaint. You should consult your doctor as soon as possible.
- lupus-like disease syndrome (including rash, joint disorders and effects on blood cells).

**Other possible side effects with Atorvastatin:**

**Common** (may affect up to 1 in 10 people):

- inflammation of the nasal passages, pain in the throat, nose bleed
- allergic reactions
- increases in blood sugar levels (if you have diabetes continue careful monitoring of your blood sugar levels), increase in blood creatine kinase
- headache
- nausea, constipation, wind, indigestion, diarrhoea
- joint pain, muscle pain and back pain
- blood test results that show your liver function can become abnormal

**Uncommon** (may affect up to 1 in 100 people):

- anorexia (loss of appetite), weight gain, decreases in blood sugar levels (if you have diabetes you should continue careful monitoring of your blood sugar levels)
- having nightmares, insomnia
- dizziness, numbness or tingling in the fingers and toes, reductions of sensation to pain or touch, change in sense of taste, loss of memory
- blurred vision
- ringing in the ears and/or head
- vomiting, belching, abdominal pain upper and lower, pancreatitis (inflammation of the pancreas leading to stomach pain)
- hepatitis (liver inflammation)
- rash, skin rash and itching, hives, hair loss
- neck pain, muscle fatigue
- fatigue, feeling unwell, weakness, chest pain, swelling especially in the ankles (oedema), raised temperature
- urine tests that are positive for white blood cells

**Rare** (may affect up to 1 in 1,000 people):

- visual disturbance
- unexpected bleeding or bruising
- cholestasis (yellowing of the skin and whites of the eyes)
- tendon injury

**Very rare** (may affect up to 1 in 10,000 people):

- an allergic reaction - symptoms may include sudden wheezing and chest pain or tightness, swelling of the eyelids, face, lips, mouth, tongue or throat, difficulty breathing, collapse
- hearing loss
- gynecomastia (breast enlargement in men).

**Not known (frequency cannot be estimated from the available data):**

- Muscle weakness that is constant.

**Possible side effects reported with some statins (medicines of the same type):**

- Sexual difficulties
- Depression
- Breathing problems including persistent cough and/or shortness of breath or fever
- Diabetes. This is more likely if you have high levels of sugars and fats in your blood, are overweight and have high blood pressure. Your doctor will monitor you while you are taking this medicine.

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine.

**Malta**

ADR Reporting  
Website: [www.medicinesauthority.gov.mt/adrportal](http://www.medicinesauthority.gov.mt/adrportal)

**The United Kingdom**

Yellow Card Scheme  
Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

## 5. How to store Atorvastatin

Keep this medicine out of the sight and reach of children.

**Blister:**

For 10 mg, 20 mg, 30 mg, 40 mg, 60 mg and 80 mg (PA/AI/PVC-AI):  
This medicine does not require any special storage conditions.

For 10 mg and 20 mg (PVC/PE/PVdC/AI):

This medicine does not require any special storage conditions.

For 40 mg and 80 mg (PVC/PE/PVdC-AI):

Store below 30 °C

**HDPE:**

This medicine does not require any special storage conditions

Use within 9 months after first opening the HDPE container.

Do not use this medicine after the expiry date which is stated on the label, carton and blister after EXP. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

**What Atorvastatin contains**

- The active substance is atorvastatin  
Each film coated tablet contains 10 mg atorvastatin (as atorvastatin calcium trihydrate).  
Each film coated tablet contains 20 mg atorvastatin (as atorvastatin calcium trihydrate).  
Each film coated tablet contains 30 mg atorvastatin (as atorvastatin calcium trihydrate).  
Each film coated tablet contains 40 mg atorvastatin (as atorvastatin calcium trihydrate).  
Each film coated tablet contains 60 mg atorvastatin (as atorvastatin calcium trihydrate).  
Each film coated tablet contains 80 mg atorvastatin (as atorvastatin calcium trihydrate).
- The other ingredients are.  
**Tablet core:** Mannitol, copovidone, sodium carbonate anhydrous, croscarmellose sodium, silicified microcrystalline cellulose (contains Silica, colloidal anhydrous) and microcrystalline cellulose, lactose monohydrate, sodium lauryl sulfate, silica colloidal anhydrous, magnesium stearate.  
**Tablet coat (Ready to use coating material):** Poly vinyl alcohol – part hydrolyzed, titanium dioxide (E171), talc (E553b), lecithin (soya) (E322), xanthan gum (E415).

**What Atorvastatin looks like and contents of the pack**

Film-coated tablet

**Atorvastatin 10 mg film-coated tablets**  
White, elliptical [9.8 mm x 5.2 mm], film-coated tablets, debossed with “Y” on one side and “77” on other side.

**Atorvastatin 20 mg film-coated tablets**  
White, elliptical [12.3 mm x 6.5 mm], film-coated tablets, debossed with “Y” on one side and “78” on other side.

**Atorvastatin 30 mg film-coated tablets**  
White, round [10.1 mm], film coated tablets, debossed with “N” on one side and “30” on other side.

**Atorvastatin 40 mg film-coated tablets**  
White, elliptical [15.5 mm x 8.1 mm], film-coated tablets, debossed with “Y” on one side and “79” on other side.

**Atorvastatin 60 mg film-coated tablets**  
White, oval [17.6 mm x 9.3 mm], film coated tablets, debossed with “N” on one side and “60” on other side.

**Atorvastatin 80 mg film-coated tablets**  
White, elliptical [19.4 mm x 10.4 mm], film-coated tablets, debossed with “Y” on one side and “72” on other side.

**For 10 mg, 20 mg, 30 mg, 40 mg, 60 mg and 80 mg:**

Atorvastatin film-coated tablets are available in polyamide/ Aluminium foil/ PVC - Aluminium foil blisters packs and HDPE bottle packs with polypropylene closure. Bottle pack contains silica gel as desiccant.

**For 10 mg, 20 mg, 40 mg and 80 mg:**  
Atorvastatin film-coated tablets are also available in PVC/PE/PVdC- Aluminium foil blister as alternate blister pack.

**Pack sizes:**

**For 10 mg, 20 mg, 40 mg and 80 mg:** 14, 28, 30, 50, 56, 90, 100 and 500 film-coated tablets

**For 30 mg and 60 mg:** 28, 30, 50, 60, 100 film-coated tablets.

**HDPE bottle pack:**

30, 90, 100, 200 and 250 film-coated tablets (for 10 mg, 20 mg and 40 mg)  
30 and 200 film-coated tablets (for 80 mg).

Not all pack sizes may be marketed.

**Marketing Authorisation Holder**

**For 10 mg, 20 mg, 30 mg, 40 mg, 60 mg, 80 mg:**

UK\_ Milpharm Limited  
Ares Block, Odyssey Business Park  
West End Road  
Ruislip HA4 6QD  
United Kingdom

**For 10 mg, 20 mg, 40 mg, 80 mg:**

MT\_ Aurobindo Pharma (Malta) Limited  
Vault 14, Level 2, Valletta Waterfront  
Floriana FRN 1913  
Malta

**Manufacturer**

APL Swift Services (Malta) Limited  
HF26, Hal Far Industrial Estate, Hal Far Birzebbugia, BBG 3000  
Malta

or

Milpharm Limited  
Ares Block, Odyssey Business Park  
West End Road  
Ruislip HA4 6QD  
United Kingdom

**This leaflet was last revised in 02/2019.**