

Package leaflet: Information for the user

Tramadol 50 mg capsules, hard

tramadol hydrochloride

This medicine contains tramadol which is an opioid, which can cause addiction. You can get withdrawal symptoms if you stop taking it suddenly.

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Tramadol is and what it is used for
2. What you need to know before you take Tramadol
3. How to take Tramadol
4. Possible side effects
5. How to store Tramadol
6. Contents of the pack and other information

1. What Tramadol is and what it is used for

Tramadol hydrochloride - the active substance in Tramadol belongs to a class of medicines called opioids, which are pain relievers. This medicine has been prescribed to you and should not be given to anyone else.

Tramadol is used for the treatment of moderate to severe pain.

Opioids can cause addiction and you may get withdrawal symptoms if you stop taking it suddenly.

Your prescriber should have explained how long you will be taking it for and when it is appropriate to stop, how to do this safely.

2. What you need to know before you take Tramadol

Do not take Tramadol

- if you are allergic to active substance or any of the other ingredients of this medicine (listed in section 6);
- in acute poisoning with alcohol, sleeping pills, pain relievers or other psychotropic medicines (medicines that affect mood and emotions);
- if you are also taking MAO inhibitors (certain medicines used for treatment of depression) or have taken them in the last 14 days before treatment with Tramadol (see "Other medicines and Tramadol");
- if you are an epileptic and your fits are not adequately controlled by treatment;
- as a substitute in drug withdrawal.

Warnings and precautions

Talk to your doctor or pharmacist before taking Tramadol, if:

- you suffer from consciousness disorders (if you feel that you are going to faint);
- you are in a state of shock (cold sweat may be a sign of this);
- you suffer from increased pressure in the brain (possibly after a head injury or brain disease);
- you have difficulty in breathing;
- you have a tendency towards epilepsy or fits because the risk of a fit may increase;
- you suffer from a liver or kidney disease;
 - you suffer from depression and you are taking antidepressants as some of them may interact with tramadol (see 'Other medicines and Tramadol').

There is a small risk that you may experience a so-called serotonin syndrome that can occur after having taken tramadol in combination with certain antidepressants or tramadol alone. Seek medical advice immediately if you have any of the symptoms related to this serious syndrome (see section 4 'Possible side effects').

Tolerance, dependence, and addiction
This medicine contains tramadol which is an opioid medicine. Repeated use of opioids can result in drug being less effective (you become accustomed to it, known as tolerance). Repeated use of tramadol can also lead to dependence, abuse and addiction, which may result in life-threatening overdose. The risk of these side effects can increase with a higher dose and longer duration of use.

Dependence or addiction can make you feel that you are no longer in control of how much medicine you need to take or how often you need to take it.

The risk of becoming dependent or addicted varies from person to person. You may have a greater risk of becoming dependent on or addicted to tramadol if:

- You or anyone in your family have ever abused or been dependent on alcohol, prescription medicines or illegal drugs ("addiction").
- You are a smoker.
- You have ever had problems with your mood (depression, anxiety, or a personality disorder) or have been treated by a psychiatrist for other mental illnesses.

If you notice any of the following signs whilst taking tramadol, it could be a sign that you have become dependent or addicted:

- You need to take the medicine for longer than advised by your doctor
- You need to take more than the recommended dose
- You are using the medicine for reasons other than prescribed, for instance, 'to stay calm' or 'help you sleep'
- You have made repeated, unsuccessful attempts to quit or control the use of the medicine
- When you stop taking the medicine you feel unwell, and you feel better once taking the medicine again ('withdrawal effects')

If you notice any of these signs, speak to your doctor to discuss the best treatment pathway for you, including when it is appropriate to stop and how to stop safely (See section 3, If you stop taking Tramadol).

Tramadol is transformed in the liver by an enzyme. Some people have a variation of this enzyme and this can affect people in different ways. In some people, they may not get enough pain relief but other people are more likely to get serious side effects. If you notice any of the following side effects, you must stop taking this medicine and seek immediate medical advice: slow or shallow breathing, confusion, sleepiness, small pupils, feeling or being sick, constipation, lack of appetite.

Sleep-related breathing disorders

Tramadol belongs to the group of opioids. Opioids can cause sleep-related breathing disorders, for example central sleep apnoea (shallow/pause of breathing during oxygen) and sleep-related hypoxemia (low oxygen level in the blood).

The risk of experiencing central sleep apnea is dependent on the dose of opioids. Your doctor may consider decreasing your total opioid dosage if you experience central sleep apnea.

Epileptic fits have been reported in patients taking tramadol at the recommended dose level. The risk may be increased when doses of tramadol exceed the recommended upper daily dose limit (400 mg).

Taking this medicine regularly, particularly for a long time, can lead to addiction. Your prescriber should have explained how long you will be taking it for and when it is appropriate to stop, how to do this safely.

Rarely, increasing the dose of this medicine can make you more sensitive to pain. If this happens, you need to speak to your prescriber about your treatment.

Addiction can cause withdrawal symptoms when you stop taking this medicine.

Withdrawal symptoms can include restlessness, difficulty sleeping, irritability, agitation, anxiety, feeling your heartbeat (palpitations), increased blood pressure, feeling or being sick, diarrhoea, loss of appetite, shaking, shivering or sweating. Your prescriber will discuss with you how to gradually reduce your dose before stopping the medicine. It is important that you do not stop taking the medicine suddenly as you will be more likely to experience withdrawal symptoms.

Opioids should only be used by those they are prescribed for. Do not give your medicine to anyone else. Taking higher doses or more frequent doses of opioid, may increase the risk of addiction. Overuse and misuse can lead to overdose and/or death.

Talk to your doctor or pharmacist if you experience any of the following symptoms while taking Tramadol:

Extreme fatigue, lack of appetite, severe abdominal pain, nausea, vomiting or low blood pressure. This may indicate that you have adrenal insufficiency (low cortisol levels). If you have these symptoms, contact your doctor, who will decide if you need to take hormone supplement.

Children and adolescents
Use in children with breathing problems: Tramadol is not recommended in children with breathing problems, since the symptoms of tramadol toxicity may be worse in these children.

Other medicines and Tramadol
Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Do not take Tramadol at the same time as medicines called "monoamine oxidase inhibitors" or MAOIs (which are used to treat e.g. depression), or if you have taken MAOIs in the past 2 weeks.

The pain-relieving effect of Tramadol may be reduced and the length of time it acts may be shortened if you also take medicines containing:

- carbamazepine (used to treat epilepsy)
- ondansetron (used to stop you feeling sick).

Your doctor will tell you whether you should take Tramadol, and which dose.

The risk of side effects increases if you take Tramadol at the same time as:

- if you are taking medicines which may cause convulsions (fits), such as certain antidepressants or antipsychotics. The risk having a fit may increase if you take Tramadol at the same time. Your doctor will tell you whether Tramadol is suitable for you.
- if you are taking certain antidepressants. Tramadol may interact with these medicines and you may experience serotonin syndrome (see section 4 'Possible side effects').
- other pain relievers such as morphine and codeine (also as cough medicine), and alcohol while you are taking Tramadol. You may feel drowsy or feel that you might faint. If this happens tell your doctor.
- coumarin anticoagulants (medicines for blood thinning), e.g. warfarin, together with Tramadol. The effect of these medicines on blood clotting may be affected and bleeding may occur.
- medicines used to treat allergies, travel sickness or nausea (antihistamines or antiemetics).
- medicines to treat psychiatric disorders (antipsychotics or neuroleptics).
- muscle relaxants.
- medicines to treat Parkinson's disease.
- concomitant use of Tramadol and tranquillizers, sleeping pills (e.g. benzodiazepines) or other sedatives or medicines that impair breathing activity (e.g. opioids) or gabapentin or pregabalin to treat epilepsy or pain due to nerve problems (neuropathic pain) increases the risk of drowsiness, difficulties in breathing (respiratory depression), coma and may be life-threatening. Because of this, concomitant use should only be considered when other treatment options are not possible.

However if your doctor does prescribe Tramadol together with sedative medicines the dose and duration of concomitant treatment should be limited by your doctor. Please tell your doctor about all sedative medicines you are taking, and follow your doctor's dose recommendation closely. It could be helpful to inform friends or relatives to be aware of the signs and symptoms stated above. Contact your doctor when experiencing such symptoms.

Tramadol with food and drink and alcohol
You should not drink alcohol during treatment with Tramadol, as its effect may be intensified.

Food does not influence the effect of Tramadol.

Pregnancy, breast-feeding and fertility
If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. Do not take Tramadol if you are pregnant or think you might be pregnant and you have discussed this with your prescriber and the benefits of treatment are considered to outweigh the potential harm to the baby.

If you use Tramadol during pregnancy, your baby may become dependent and experience withdrawal symptoms after the birth which may need to be treated.

Do not take Tramadol while you are breastfeeding as tramadol passes into breast milk and will affect your baby.

Based on human experience tramadol is suggested not to influence female or male fertility.

Driving and using machines
Tramadol may cause drowsiness, dizziness and blurred vision and therefore may impair your reactions. If you feel that your reactions are affected, do not drive a car or other vehicle, do not use electric tools or operate machinery.

The medicine can affect your ability to drive as it may make you sleepy or dizzy.

- Do not drive while taking this medicine until you know how it affects you.
- It is an offence to drive if this medicine affects your ability to drive.
- However, you would not be committing an offence if:
 - The medicine has been prescribed to treat a medical or dental problem and
 - You have taken it according to the instructions given by the prescriber or in the information provided with the medicine and
 - It was not affecting your ability to drive safely

Talk to your doctor or pharmacist if you are not sure whether it is safe for you to drive while taking this medicine.

Tramadol contains Sodium:

Tramadol capsule contains Sodium. This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

3. How to take Tramadol

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Before starting treatment and regularly during treatment, your doctor will discuss with you what you may expect from using tramadol, when and how long you need to take it, when to contact your doctor, and when you need to stop it (see also section 2).

The dosage should be adjusted to the intensity of your pain and your individual pain sensitivity. In general the lowest pain-relieving dose should be taken. Do not take more than 400 mg tramadol hydrochloride daily, except if your doctor has instructed you to do so.

Unless otherwise prescribed by your doctor, the usual dose is:

Adults and adolescents from the age of 12 years

One or two Tramadol (equivalent to 50 mg – 100 mg tramadol hydrochloride)

Depending on the pain the effect lasts for about 4-8 hours.

Your doctor may prescribe a different, more appropriate dosage of Tramadol if necessary.

Children below 12 years of age

Tramadol is not recommended for use in children below age 12.

Elderly patients

In elderly patients (above 75 years) the excretion of tramadol may be delayed. If this applies to you, your doctor may recommend prolonging the dosage interval.

Severe liver or kidney disease (insufficiency)/ dialysis patients

Patients with severe liver and/or kidney insufficiency should not take Tramadol.

If in your case the insufficiency is mild or moderate, your doctor may recommend prolonging the dosage interval.

How and when should you take Tramadol?

Tramadol are for oral use.

Always swallow Tramadol whole, not divided or chewed, with sufficient liquid, preferably in the morning and evening. You may take the capsule on an empty stomach or with meals.

How long should you take Tramadol?

Your prescriber should have discussed with you, how long the course of capsules will last. They will arrange a plan for stopping treatment. This will outline how to gradually reduce the dose and stop taking the medicine.

If you take more Tramadol than you should

If you have taken an additional dose by mistake, this will generally have no negative effects. You should take your next dose as prescribed.

If you forget to take Tramadol

If you forget to take Tramadol take it as soon as you remember and then carry on as before.

Do not take a double dose to make up for a forgotten dose.

If you stop taking Tramadol

If you stop taking Tramadol, your pain may return.

You should not suddenly stop taking this medicine unless your doctor tells you to. If you want to stop taking your medicine, discuss this with your

doctor first, particularly if you have been taking it for a long time. Your doctor will advise you when and how to stop, which may be by lowering the dose gradually to reduce the chance of developing unnecessary side effects (withdrawal symptoms).

If you would like to stop treatment because you have unpleasant side effects, please talk to your doctor. If you have been taking this medicine for a very long time, you may get the following side effects if you suddenly stop treatment: restlessness, difficulty sleeping, irritability, agitation, anxiety, feeling your heartbeat (palpitations), increased blood pressure, feeling or being sick, diarrhoea, shaking, shivering or sweating. If you get any of these effects after stopping treatment with Tramadol please talk to your doctor.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

You should see a doctor immediately if you experience symptoms of an allergic reaction such as swollen face, tongue and/or throat, and/or difficulty swallowing or hives together with difficulties in breathing.

The most common side effects during treatment with Tramadol are nausea and dizziness, which occur in more than 1 in 10 people.

The frequency of side effects is classified as follows:

Common: may affect up to 1 in 10 people

Uncommon: may affect up to 1 in 100 people

Rare: may affect up to 1 in 1,000 people

Very rare: may affect up to 1 in 10,000 people

Not known: frequency cannot be estimated from the available data

The following side effects may occur:

Very common: feeling sick (nausea); dizziness.

Common: headache, drowsiness, fatigue, vomiting (being sick), constipation, dry mouth; sweating (hyperhidrosis).

Uncommon: effects on the heart and blood circulation (pounding of the heart, fast heartbeat, feeling faint or collapse). These adverse effects may particularly occur in patients in an upright position or under physical strain; diarrhoea, urge to sick (retching), gastrointestinal pressure (a feeling of pressure in the stomach, bloating); skin disorders (e.g. itchiness, rash, sudden onset of skin redness).

Rare: slow heartbeat, increase in blood pressure; change in appetite,

abnormal sensations (e.g. itching, tingling, numbness); trembling, breathing difficulties, epileptic-like seizures ("fits"), uncoordinated movement, transient loss of consciousness (syncope); muscle twitches, fainting; blurred vision, dilation of pupils (mydriasis), constriction of pupils (miosis); slow breathing, shortness of breath (dyspnoea); Worsening of asthma has been reported, however it has not been established whether it was caused by tramadol. If the recommended doses are exceeded, or if other medicines that depress brain function are taken at the same time, breathing may slow down; difficulty in passing urine or pain, passing less urine than normal (dysuria). Muscle weakness. Allergic reactions (e.g. difficulty in breathing, wheezing, swelling of skin) and shock (sudden circulation failure) have occurred in very rare cases.). Hallucinations, confusion, anxiety, sleep disturbance and nightmares, changes in mood (high or low spirits), changes in activity (slowing down but sometimes an increase in activity) and being less aware and less able to make decisions, which may lead to errors in judgement.

Very rare: hepatic enzyme increased

Not known: decrease in blood sugar level, Hiccups, Serotonin syndrome, that can manifest as mental status changes (e.g. agitation, hallucinations, coma), and other effects, such as fever, increase in heart rate, unstable blood pressure, involuntary twitching, muscular rigidity, lack of coordination and/or gastrointestinal symptoms (e.g. nausea, vomiting, diarrhoea) (see section 2 'What you need to know before you take Tramadol'), dependence and addiction (see section 'How do I know if I am addicted').

Drug Withdrawal

When you stop taking Tramadol, you may experience drug withdrawal symptoms, which include restlessness, difficulty sleeping, irritability, agitation, anxiety, feeling your heartbeat (palpitations), increased blood pressure, feeling or being sick, diarrhoea, shaking, shivering or sweating

How do I know if I am addicted?

If you notice any of the following signs whilst taking Tramadol, it could be a sign that you have become addicted.

- You need to take the medicine for longer than advised by your prescriber

- You feel you need to use more than the recommended dose

- You are using the medicine for reasons other than prescribed

- When you stop taking the medicine you feel unwell, and you feel better once taking the medicine again

If you notice any of these signs, it is important you talk to your prescriber.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme, Website: www.mhra.gov.uk/yellowcard or search for [MHRA Yellow Card in the Google Play or Apple App Store](http://www.mhra.gov.uk/yellowcard). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Tramadol

Keep this medicine out of the sight and reach of children.

Store this medicine in a safe and secure storage space, where other people cannot access it. It can cause serious harm and be fatal to people when it has not been prescribed for them.

Do not use this medicine after the expiry date which is stated on the carton, bottle after EXP. The expiry date refers to the last day of that month.

Store below 30°C.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Tramadol contains

The active substance is tramadol hydrochloride. Each hard capsule contains 50 mg tramadol hydrochloride.

The other ingredients are:

Capsule contents: Cellulose, microcrystalline, silica colloidal anhydrous, sodium starch glycolate (Type A), magnesium stearate.

Capsule shell: Gelatin, sodium lauryl sulphate, indigo carmine, iron oxide yellow (E172) and titanium dioxide (E171).

Printing ink: Shellac and black iron oxide (E172).

What Tramadol looks like and contents of the pack

filled /Yellow, size '4' hard gelatin capsules filled with white to off-white powder and imprinted with 'T' on green cap and '02' on yellow body with black ink.

PVC/PVDC/Aluminium foil blister: 10, 20, 30, 50, 60, 90, 100 and 500 Capsules

HDPE bottle pack with polypropylene closure: 30,200 and 500 Capsules

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Milpharm Limited
Ares Block
Odyssey Business Park
West End Road
Ruislip HA4 6QD
United Kingdom

Manufacturer

APL Swift Services (Malta) Limited
HF26, Hal Far Industrial Estate, Hal Far,
Birzebbugia, BBG 3000
Malta

or

Milpharm Limited
Ares Block
Odyssey Business Park
West End Road
Ruislip HA4 6QD
United Kingdom

This leaflet was last revised in 10/2025.