

Applying the ointment

To be used in the eye(s) only.

- 1. Check the tube seal is not broken before first using the eve ointment.
- 2. Wash and dry your hands
- 3. Remove the cap. Take special care that the nozzle of the tube does not touch your eye,
- the skin around your eye or your fingers. 4. Use a mirror, this will help you see what you are doing
- 5. Place the nozzle of the tube close to your eye. Gently pull the lower eyelid downwards and look up.
- 6. Squeeze the tube gently and apply about 1cm of the ointment to the space between the lower eyelid and the eye.
- 7. Close for a moment
- 8. When both eyes are to be treated, repeat
- 9. Replace the cap securely after use.

Whilst applying the ointment, do not

- Breathe on or touch the nozzle of the tube • Touch the eyes or eyelids with the nozzle of the
- Share your ointment with anyone else

For external use only

The dose is a small amount of ointment (approx 1cm) to be applied to the affected eye(s) If you are using **drops during the day** and the ointment at night - apply the ointment at night, before going to bed.

If you are using **just the ointment** - apply **3 to 4** times a day.

Length of treatment

The course of treatment is 5 days. **Keep using** the eye ointment for 5 days, do not stop just because vou feel better - this could make your condition

If your symptoms do not improve within 48 hours, consult your doctor. If your symptoms get worse, 'seek medical advice at once.

Do not use Optrex Bacterial Conjunctivitis 1% w/w Eye Ointment for **more than 5 days** without consulting your doctor. Discard any remaining ointment after the 5 day course of treatment. (see section 5).

If you forget to use your ointment

If you have only just missed a dose and it is a long time before the next dose is due, put in the missed dose of ointment straight away. If it is nearly time for your next dose, just put in the next dose of ointment at the usual time. Do not take a double dose to make up for a forgotten dose.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Some people may get one or more of these side effects. Some effects happen straight away but do not last long - others may only happen after several days of

Possible side effects include :

* allergic reactions such as itching or rashes. If you

""" - or bysothing

9-COLOP-36-GB-PV2 get a severe reaction, with swelling or breathing problems get medical help straight away.

3. How to use Optrex Bacterial Conjunctivitis • changes in the blood (anaemia) leading to severe tiredness or easy bruising. Stop using the treatment and consult your doctor if you

E

• The use of topical chloramphenicol may occasionally result in the overgrowth of other non susceptible organisms including funghi. If any new infection appears during treatment you should tell your doctor.

• blurred vision or mild burning or stinging when you put the ointment in. These should subside

f you do not feel well or are worried about your héalth, talk to a pharmacist or doctor. If any of the side effects get serious or if you experience any side effect not listed in this leaflet please tell your doctor or pharmacist.

Reporting of side effects

experience these effects.

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/vellowcard By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Optrex Bacterial Conjunctivitis 1% w/w Eye Ointment

Replace the cap securely after use. Store below 25°C.

Keep all medicines out of the sight and reach of children. Do not use after the 'use by' date shown on the tube and end of the carton.

Discard any unused ointment after completing your 5 day course of treatment Do not throw away any medicines via wastewater

or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other

What Optrex Bacterial Conjunctivitis 1% w/w Eye Ointment contains

The active substance is chloramphenicol. One (1) gram of ointment contains 10 mg of chloramphenicol. The other ingredients are liquid paraffin and white petroleum.

What Optrex Bacterial Conjunctivitis 1% w/w Eye Ointment looks like and contents of the pack

Each pack contains one tube with 4g of ointment.

Marketing Authorisation Holder

Medicom Healthcare Ltd., Lynton House, 7-12 Tavistock Square Kings Cross, London, WC1H oLT, U.K.

Manufacture

TUBILUX PHARMA S.P.A, via Costarica 20/22, 00071 Pomezia (RM), Italy

Distributed by

Optrex Limited, Slough, SL1 3UH

PL number: 18956/0009

Leaflet revised December 2019.

🚤 10 mm

VOLTA BACK PANEL

> **CUSTOMER INFO:** Minimum Point Size = 9.00pt