

Package leaflet: Information for the user

Decapeptyl® SR 11.25 mg Powder and solvent for suspension for injection Triptorelin

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, please ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Decapeptyl SR 11.25 mg is and what it is used for
2. What you need to know before you use Decapeptyl SR 11.25 mg
3. How to use Decapeptyl SR 11.25 mg
4. Possible side effects
5. How to store Decapeptyl SR 11.25 mg
6. Contents of the pack and other information

1. What Decapeptyl SR 11.25 mg is and what it is used for

The active ingredient in Decapeptyl SR 11.25 mg is triptorelin. Triptorelin belongs to a group of medicines called gonadotropin releasing hormone (GnRH) agonists. Triptorelin is similar to the gonadotropin releasing hormone which occurs naturally in your body.

In men, triptorelin lowers the levels of the hormone testosterone.

In women, it reduces oestrogen levels.

Decapeptyl SR 11.25 mg has three different uses. It is used in men, women and children to treat completely different conditions.

Decapeptyl SR is available in two other strengths: Decapeptyl SR 3 mg is used once a month and Decapeptyl SR 22.5 mg is used once every 6 months. Not all dose strengths are approved for all indications. Ask your doctor if you would like to discuss changing your treatment.

This leaflet gives information for all three uses of Decapeptyl SR 11.25 mg. Please read all the sections that are about you and your condition.

MEN

In men, Decapeptyl SR 11.25 mg is used to treat prostate cancer.

WOMEN

In women, Decapeptyl SR 11.25 mg is used to treat Endometriosis - a condition in which the tissue that normally lines the uterus (endometrium) grows in other places.

CHILDREN

In children, Decapeptyl SR 11.25 mg is used to treat puberty that occurs at a very young age, i.e. before 8 years in girls and 10 years in boys (Central Precocious Puberty). This is called 'early puberty' in the rest of this leaflet.

2. What you need to know before you use Decapeptyl SR 11.25 mg

MEN

Do not use Decapeptyl SR 11.25 mg:

- If you are **allergic** to triptorelin or similar types of drugs (other GnRH agonists) or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions:

Talk to your doctor or pharmacist before taking Decapeptyl SR 11.25 mg.

- There have been reports of depression in patients taking Decapeptyl SR 11.25 mg which may be severe. If you are taking Decapeptyl SR 11.25 mg and develop depressed mood, **inform your doctor**. Your doctor may want to monitor your depression during treatment.
- If you are using medicines for delaying your normal blood clotting, you may experience bruising at the site of the intramuscular injection.
- In adults, triptorelin may cause thinning of the bones (osteoporosis) with an increased risk of bone fractures. You should therefore tell your doctor if you have any of the below risk factors as he/she might give you bisphosphonate (drugs used to treat weak bones) to treat bone loss. Risk factors may include:
 - If you or any of your close family have thinning of the bones.
 - If you drink excessive amounts of alcohol, and/or smoke heavily.
 - If you take medicines over a long period of time that may cause thinning of the bones, for example medicines for epilepsy or steroids (such as hydrocortisone or prednisolone).
- If any convulsions occur, inform immediately your doctor. There have been reports of convulsions in patients receiving triptorelin or similar medicines. These occurred in patients with or without medical history of epilepsy.
- When you first start treatment with Decapeptyl SR 11.25 mg it actually **increases** the level of your hormones for a short time. This means that you may feel worse to begin with (see section 4 'Possible side effects' for more information). The doctor may give you some medicine (an anti-androgen) to prevent your symptoms from getting worse. After a short time the amount of hormone will drop and your symptoms will get better.

If you suffer from urinary obstruction or spinal cord (nerves in your backbone) compression due to your prostate cancer spreading, your doctor will supervise you closely for the first few weeks of treatment. If you experience difficulty passing urine, bone pain, weakness of lower limbs or pins and needles sensation, contact your doctor immediately, who will assess and treat you appropriately.
- **Tell your doctor** if you have diabetes.
- **Tell your doctor** if you have any heart or blood vessel conditions, including heart rhythm problems (arrhythmia), or are being treated with medicines for these conditions. The risk of heart rhythm problems may be increased when using Decapeptyl SR 11.25 mg.
- If you have an enlargement (benign tumour) of the pituitary gland that you were unaware of, this may be discovered during treatment. Symptoms include sudden headache, problems with eyesight and paralysis of the eye muscles.
- After surgical castration triptorelin does not induce any further decrease in serum testosterone levels.

- Diagnostic tests of pituitary gonadal function or sex organs conducted during treatment or after discontinuation of therapy with Decapeptyl SR 11.25 mg may be misleading.
- **Tell your doctor** if you have back pain, weakness, numbness or tingling in your legs.
- Treatment with GnRH analogues including Decapeptyl SR 11.25mg might increase the risk of anaemia (defined as a decrease in the count of red blood cells).

Your doctor may give you another drug to help you feel better during this time.

Other medicines and Decapeptyl SR 11.25 mg:

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Decapeptyl SR 11.25 mg might interfere with some medicines used to treat heart rhythm problems (e.g. quinidine, procainamide, amiodarone and sotalol) or might increase the risk of heart rhythm problems when used with some other drugs (e.g. methadone (used for pain relief and part of drug addiction detoxification), moxifloxacin (an antibiotic), antipsychotics used for serious mental illnesses).

Drugs which increase the level of a hormone called prolactin may react with Decapeptyl SR 11.25 mg. Many different kinds of drugs may increase prolactin levels.

Driving and using machines:

You may feel dizzy, tired or have problems with your sight such as blurred vision. These are possible side effects of treatment or from the underlying disease. If you experience any of these side effects, you should not drive or use machines.

Decapeptyl SR 11.25 mg contains sodium

This medicine contains less than 1 mmol (23 mg) sodium per dose, i.e., essentially 'sodium-free'.

WOMEN

Do not use Decapeptyl SR 11.25 mg:

- If you are **allergic** to triptorelin or similar types of drugs (other GnRH agonists) or any of the other ingredients of this medicine (listed in section 6).
- If you are pregnant or breast-feeding.

Warnings and precautions:

Talk to your doctor or pharmacist before taking Decapeptyl SR 11.25 mg.

- Due to lack of clinical experience in women under 18 years of age, Triptorelin is not recommended in adolescent and young women as it might cause thinning of bone.
- There have been reports of depression in patients taking Decapeptyl SR 11.25 mg which may be severe. If you are taking Decapeptyl SR 11.25 mg and develop depressed mood, **inform your doctor**. Your doctor may want to monitor your depression during treatment.
- If you are using medicines for delaying your normal blood clotting, you may experience bruising at the site of the intramuscular injection.
- In adults, triptorelin may cause thinning of the bones (osteoporosis) with an increased risk of bone fractures. You should therefore tell your doctor if you have any of the below risk factors as he/she might give you bisphosphonate (drugs used to treat weak bones) to treat bone loss. Risk factors may include:
 - If you or any of your close family have thinning of the bones.
 - If you drink excessive amounts of alcohol, and/or smoke heavily.
 - If you take medicines over a long period of time that may cause thinning of the bones, for

example medicines for epilepsy or steroids (such as hydrocortisone or prednisolone).

- If any convulsions occur, inform immediately your doctor. There have been reports of convulsions in patients receiving triptorelin or similar medicines. These occurred in patients with or without medical history of epilepsy.
- When you first start treatment with Decapeptyl SR 11.25 mg it actually **increases** the level of your hormones for a short time. This means that you may feel worse to begin with (see section 4 'Possible side effects' for more information). After a short time the amount of hormone will drop and your symptoms will get better.
- **Tell your doctor** if you have diabetes.
- **Tell your doctor** if you have any heart conditions.
- If you have an enlargement (benign tumour) of the pituitary gland that you were unaware of, this may be discovered during treatment. Symptoms include sudden headache, problems with eyesight and paralysis of the eye muscles.
- You may have some vaginal bleeding in the first month of treatment. After that your periods normally stop.
- **Tell your doctor** if you have bleeding after the first month of treatment.
- Your periods should start approximately 5 months after the last injection.

You must use some form of contraception other than the 'pill' while you are having treatment and until you start your next period. Your doctor may suggest using a barrier method of contraception such as a condom or diaphragm (cap).

Other medicines and Decapeptyl SR 11.25 mg:

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Drugs which increase the level of a hormone called prolactin may react with Decapeptyl SR 11.25 mg. Many different kinds of drugs may increase prolactin levels.

Pregnancy and breast-feeding:

Do not take Decapeptyl SR 11.25 mg if you are pregnant or breast-feeding.

Driving and using machines:

You may feel dizzy, tired or have problems with your sight such as blurred vision. These are possible side effects of treatment or from the underlying disease. If you experience any of these side effects, you should not drive or use machines.

Decapeptyl SR 11.25 mg contains sodium

This medicine contains less than 1 mmol (23 mg) sodium per dose, i.e., essentially 'sodium-free'.

CHILDREN

Do not use Decapeptyl SR 11.25 mg:

- If you are **allergic** to triptorelin or similar types of drugs (other GnRH agonists) or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions:

Talk to your doctor or pharmacist before taking Decapeptyl SR 11.25 mg.

- There have been reports of depression in patients taking Decapeptyl SR 11.25 mg which may be severe. If you are taking Decapeptyl SR 11.25 mg and develop depressed mood, **inform your doctor**. Your doctor may want to monitor your depression during treatment.
- If you are using medicines for delaying your normal blood clotting, you may experience bruising at the site of the intramuscular injection.
- If you have a progressive brain tumour, **tell your doctor**. This may affect the way your doctor decides to treat you.
- If your child suffers from a bad or recurrent headache, problems with eyesight and ringing or buzzing in the ears, contact a doctor immediately (see section 4).
- Girls who have an early puberty may have some vaginal bleeding in the first month of treatment.
- In girls, menstrual bleeding will start on average one year after stopping treatment.
- Early puberty caused by other diseases should be ruled out by your doctor.
- **Tell your doctor** if you have diabetes.
- **Tell your doctor** if you have any heart conditions.
- If you have an enlargement (benign tumour) of the pituitary gland that you were unaware of, this may be discovered during treatment. Symptoms include sudden headache, problems with eyesight and paralysis of the eye muscles.
- A pathology of the hip may occur after stopping treatment (slipped capital femoral epiphysis of the hip). It results in stiffness of the hip, a limp and / or severe pain in the groin radiating to the thigh. If this occurs, you should consult your doctor.
- If any convulsions occur, inform immediately your doctor. There have been reports of convulsions in patients receiving triptorelin or similar medicines. These occurred in patients with or without medical history of epilepsy.

Other medicines and Decapeptyl SR 11.25 mg:

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Drugs which increase the level of a hormone called prolactin may react with Decapeptyl SR 11.25 mg. Many different kinds of drugs may increase prolactin levels.

Decapeptyl SR 11.25 mg contains sodium

This medicine contains less than 1 mmol (23 mg) sodium per dose, i.e., essentially ‘sodium-free’.

3. How to use Decapeptyl SR 11.25 mg

MEN

Decapeptyl SR 11.25 mg will be injected into a muscle, usually your bottom, by a doctor or nurse. On this leaflet there are instructions for them that explain how to prepare the injection.

You will normally receive an injection once every 3 months.

Also read 'Other medicines and Decapeptyl SR 11.25 mg' in section 2.

If you are given more Decapeptyl SR 11.25 mg than you should

If you are given too much Decapeptyl SR 11.25 mg you may experience additional or more severe side effects (see section 4 'Possible side effects').

If you forget to take a dose of Decapeptyl SR 11.25 mg

As soon as you realise that you have missed an injection you should **tell your doctor**. You will then be given your next injection.

If you stop receiving Decapeptyl SR 11.25 mg

If you stop receiving your Decapeptyl SR 11.25 mg injection before your doctor tells you to then your symptoms are likely to return.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

WOMEN

Decapeptyl SR 11.25 mg will be injected into a muscle, usually your bottom, by a doctor or nurse. On this leaflet there are instructions for them that explain how to prepare the injection.

You will normally receive two injections, the second one three months after the first. Each injection will be given in the first five days of your period.

Also read 'Other medicines and Decapeptyl SR 11.25 mg' in section 2.

If you are given more Decapeptyl SR 11.25 mg than you should

If you are given too much Decapeptyl SR 11.25 mg you may experience additional or more severe side effects (see section 4 'Possible side effects').

If you forget to take a dose of Decapeptyl SR 11.25 mg

As soon as you realise that you have missed an injection you should **tell your doctor**. You will then be given your next injection.

If you stop receiving Decapeptyl SR 11.25 mg

If you stop receiving your Decapeptyl SR 11.25 mg injection before your doctor tells you to then your symptoms are likely to return.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

CHILDREN

Decapeptyl SR 11.25 mg will be injected into a muscle, usually your bottom, by a doctor or nurse. On this leaflet there are instructions for them that explain how to prepare the injection.

You will normally receive an injection once every 3 months.

Your doctor will decide when treatment should be stopped (normally when you are about 12-13 if you are a girl and about 13-14 if you are a boy).

Also read 'Other medicines and Decapeptyl SR 11.25 mg' in section 2.

If you are given more Decapeptyl SR 11.25 mg than you should

If you are given too much Decapeptyl SR 11.25 mg you may experience additional or more severe side effects (see section 4 'Possible side effects').

If you forget to take a dose of Decapeptyl SR 11.25 mg

As soon as you realise that you have missed an injection you should **tell your doctor**. You will then be given your next injection.

If you stop receiving Decapeptyl SR 11.25 mg

If you stop receiving your Decapeptyl SR 11.25 mg injection before your doctor tells you to then your symptoms are likely to return.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, Decapeptyl SR 11.25 mg can have side effects although not everybody gets them.

In rare cases you may experience a severe allergic reaction (angioedema, anaphylactic reaction). Tell your doctor immediately if you develop symptoms such as swallowing or breathing problems, dizziness, a rash, swelling of your lips, face, throat or tongue.

MEN

Many of the side effects are expected, due to the change in the level of testosterone in your body. These effects include hot flushes, impotence and decreased libido.

Side effects which are **very common** (may affect more than 1 in 10 people) are hot flushes, weakness, excessive sweating, back pain, pins and needles sensation in the legs, reduced libido and impotence.

Side effects which are **common** (may affect up to 1 in 10 people) are nausea, dry mouth, pain, bruising, redness and swelling at injection site, muscle and bone pain, pain in the arms and legs, oedema (build-up of fluid in the body tissues), lower abdominal pain, high blood pressure, allergic reaction, increase in weight, dizziness, headache, loss of libido, depression and mood changes.

Side effects which are **uncommon** (may affect up to 1 in 100 people) are increase of blood platelets, feeling your heartbeat, ringing in the ears, vertigo, blurred vision, pain in abdomen, constipation, diarrhoea, vomiting, drowsiness, severe shivering associated with sweating and a fever, sleepiness, pain, swelling of the ankles, feet or fingers, some blood tests affected (including raised liver function tests), blood pressure

increased, weight loss, loss of appetite, increase of appetite, gout (severe pain and swelling in the joints usually in the big toe), diabetes, excessive lipids in the blood, joint pain, muscle cramp, muscle weakness, muscle pain, swelling and tenderness, bone pain, tingling or numbness, inability to sleep, feeling of irritability, development of enlarged breasts in men, breast pain, reduction in testicular size, pain in testicles, difficulty in breathing, acne, hair loss, itching, rash, redness of skin, hives, waking up to pass urine, problems passing urine and nosebleeds.

Side effects which are **rare** (may affect up to 1 in 1,000 people) are red or purple discolorations on the skin, abnormal sensation in the eye, blurring or disturbance in vision, sensation of fullness in the abdomen, flatulence, abnormal sense of taste, chest pain, difficulty in standing, flu-like symptoms, fever, anaphylactic reaction (serious allergic reaction which can cause dizziness or difficulty in breathing, swelling of the face or throat), inflammation of the nose/throat, increased body temperature, stiff joints, joint swelling, musculoskeletal stiffness, osteoarthritis, memory loss, feeling confused, decreased activity, having a feeling of elation, shortness of breath when lying flat, blisters and low blood pressure.

During post-marketing surveillance the following side effects have also been reported (their frequency cannot be estimated from the available data): changes in ECG (QT prolongation), serious allergic reaction which can lead to a swelling of the face, the tongue and the neck, dizziness or breathing difficulties (Quincke oedema, anaphylactic shock), convulsions, general discomfort, anxiety, rapid formation of wheals due to swelling of the skin or mucous membranes and urinary incontinence, if there is an existing pituitary tumour there is an increased risk of bleeding to the area, anaemia (decrease in the count of red blood cells).

As with other GnRH agonists, an increase in white blood cell count may be found in patients being treated with Decapeptyl SR 11.25 mg.

Patients receiving long-term treatment by GnRH analogue in combination with radiation may have more side effects especially gastrointestinal, related to radiotherapy.

WOMEN

Many of the side effects are expected due to the change in the level of oestrogens in your body.

These **very common** side effects (may affect more than 1 in 10 people) include headache, decreased libido, mood swings, difficulty in sleeping, breast disorder, ovarian hyperstimulation syndrome, pain during or after sexual intercourse, painful periods, genital bleeding, pelvic pain, dryness of the vagina, weakness, excessive sweating, acne, oily skin and hot flushes.

Side effects which are **common** (may affect up to 1 in 10 people) are breast pain, muscle cramps, painful joints, weight gain, feeling sick, depression (long term treatment), nervousness, abdominal pain or discomfort, pain, bruising, redness and swelling at injection site, swelling and tenderness, allergic reaction, pain in the arms and legs, dizziness and swelling of ankles, feet or fingers.

Side effects which are **uncommon** (may affect up to 1 in 100 people) are feeling your heartbeat, vertigo, dry eye, blurred vision, bloating, vomiting, diarrhoea, dry mouth, flatulence, mouth ulcer, weight decrease, decrease in appetite, water retention, back pain, muscle pain, abnormal taste, loss of sensations, temporary loss of consciousness, memory loss, lack of concentration, tingling or numbness, involuntary muscle movement, mood change, anxiety, disorientation, depression (short term treatment), bleeding after sex, prolapse, irregular period, painful period and heavy period, small cysts (swelling) on the ovaries which can

cause pain, discharge from the vagina, difficulty breathing, nosebleed, hair loss, dry skin, excessive bodily hair, brittle nails, itching, hives and skin rash.

During post-marketing surveillance the following side effects have also been reported (their frequency cannot be estimated from the available data): general discomfort, increased blood pressure, increased body temperature, serious allergic reaction which can lead to a swelling of the face, the tongue and the neck, dizziness or breathing difficulties (Quincke oedema, anaphylactic shock), convulsions, some blood tests affected (including raised liver function tests) muscle weakness, confusion, absence of menstrual periods, rapid formation of wheals due to swelling of the skin or mucous membranes, abnormal sensations in the eyes and/or changes in sight, hives, if there is an existing pituitary tumour there is an increased risk of bleeding to the area.

In endometriosis treatment, the disorders for which the treatment has been justified (pelvic pain, dysmenorrhea) may be exacerbated at the beginning of the treatment, but should disappear in one to two weeks. This may occur even if the treatment is producing a favorable effect. You should nevertheless immediately notify your doctor of this phenomenon.

CHILDREN

Side effects which are **very common** (may affect more than 1 in 10 people) include vaginal bleeding which may occur in girls in the first month of treatment.

Side effects which are **common** (may affect up to 1 in 10 people) include pain in abdomen, pain, bruising, redness and swelling at injection site, headache, hot flushes, weight gain, acne, allergic reactions.

Side effects which are **uncommon** (may affect up to 1 in 100 people) are blurred vision, vomiting, constipation, nausea, general discomfort, overweight, neck pain, changes in mood, pain in breast, nosebleeds, itching, rash or hives in the skin.

Long term trial (up to 4 years) did not bring any new and significant safety concerns.

During post-marketing surveillance the following side effects have also been reported (their frequency cannot be estimated from the available data): high blood pressure, abnormal vision, serious allergic reaction which can lead to a swelling of the face, the tongue and the neck, dizziness or breathing difficulties (Quincke oedema, anaphylactic shock), convulsions, some blood tests affected include hormone levels, rapid formation of wheals due to swelling of the skin or mucous membranes, muscle pain, mood disorders, depression, nervousness, Idiopathic intracranial hypertension (increased intracranial pressure around the brain characterised by headache, double vision and other visual symptoms, and ringing or buzzing in the ears).

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme. Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store Decapeptyl SR 11.25 mg

Keep this medicine out of the sight and reach of children.

Do not use the vial or ampoule after the expiry date printed on the box.

This medicine should not be stored above 25°C. The vial and ampoule should be kept in the outer box.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Decapeptyl SR 11.25 mg contains:

The active substance of Decapeptyl SR 11.25 mg is triptorelin. Each vial contains sufficient quantity of triptorelin (as triptorelin pamoate) to ensure that the minimum triptorelin quantity injected is 11.25 mg. The other ingredients are D,L lactide-glycolide copolymer, mannitol, carmellose sodium, polysorbate 80.

What Decapeptyl SR 11.25 mg looks like and contents of the pack

Each pack contains:

- 1 clear glass vial with a rubber stopper and an aluminium cap containing the powder
- 1 glass ampoule containing the suspension vehicle
- 1 syringe
- 2 needles.

Marketing Authorisation Holder

Ipsen Limited, 5th Floor, The Point, 37 North Wharf Road, Paddington, London, W2 1AF, UK.

Manufacturer

Ipsen Pharma Biotech, Signes, France.

This leaflet was last revised in July 2024.

Is this leaflet hard to see or read? Please phone +44 (0) 1753 627777 and ask for help.

The following information is intended for medical or healthcare professionals only:

INSTRUCTIONS FOR RECONSTITUTION


1. PREPARATION OF THE PATIENT BEFORE RECONSTITUTION

- Prepare the patient by disinfecting the injection site. This operation needs to be performed first because once reconstituted, the drug should be injected immediately.

2. PREPARATION OF THE INJECTION

Two needles are provided in the box :

- **Needle 1** : a 20G needle (38 mm of length) without safety device to be used for reconstitution
- **Needle 2** : a 20G needle (38 mm of length) with safety device to be used for injection




The presence of bubbles on top of the lyophilisate is a normal appearance of the product.

The following steps must be completed in a continuous sequence.







2a


- Take out the ampoule containing the solvent. Tap any solution within the tip of the ampoule back to the main body of the ampoule.
- Screw **Needle 1** (without safety device) on to the syringe. Do not remove the needle protection yet.
- Break open the ampoule with dot face up.
- Remove the needle protection from Needle 1. Insert the needle in the ampoule and draw up all the solvent into the syringe.
- Put aside the syringe containing the solvent.



2b

- Take out the vial containing the powder. Tap any powder which has accumulated at the top of the vial back to the bottom of the vial.
- Remove the plastic tab on top of the vial.
- Take back the syringe containing the solvent, and insert the needle through the rubber stopper vertically into the vial. Inject the solvent slowly, so that, if possible, it washes down the entire upper part of the vial.


	<p>2c</p> <ul style="list-style-type: none"> • Pull up Needle 1 above the liquid level. Do not remove the needle from the vial. Reconstitute the suspension, by swirling gently from side to side. Do not invert the vial. • Continue swirling long enough (at least 30 seconds) to obtain a homogeneous and milky suspension. • Important: Check there is no unsuspended powder in the vial (if any powder clumps are present, continue swirling until they disappear).
 	<p>2d</p> <ul style="list-style-type: none"> • When the suspension is homogeneous, pull down the needle and without inverting the vial, draw up all of the suspension. A small amount will remain in the vial and should be discarded. An overfill is included to allow for this loss. • Grasp the coloured hub to disconnect the needle. Remove Needle 1 used for the reconstitution from the syringe. Screw on to the syringe Needle 2. • Move the safety sheath away from the needle and towards the syringe barrel. The safety sheath remains in the position you set. • Remove the needle protection from the needle. • Prime the needle to remove air from the syringe and inject immediately.
<p>3. INTRAMUSCULAR INJECTION</p>	
	<ul style="list-style-type: none"> • To avoid sedimentation, inject immediately into the disinfected area as quickly as possible (within 1 minute from reconstitution).
<p>4. AFTER USE</p>	
<ul style="list-style-type: none"> • Activation of the safety system using a one-handed technique. • Note: Keep your finger behind the tab at all times. <p>There are two alternatives to activate the safety system:</p> <ul style="list-style-type: none"> • Method A : push the tab forward with your finger 	



Method A


or

- Method B : push the sheath to a flat surface



Method B

- In both cases press down with a firm quick motion until a distinct audible click is heard.
- Visually confirm that the needle is fully engaged under the lock.



- Used needles, any unused suspension or other waste materials should be disposed of in accordance with local requirements.