

Patient Information Leaflet



Ibuprofen 5% w/w Gel

Read all of this leaflet carefully before you start using this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions ask your doctor or pharmacist.
- This medicine may have been prescribed for you by your doctor. Do not pass it on to others. It may harm them even if their symptoms are the same as yours.
- This medicine is also available without a prescription from your pharmacy. However, you still need to use Ibuprofen Gel carefully to get the best results from it.
- You must see a doctor if your symptoms do not improve after **2 weeks**.
- If any of the side effects gets serious or if you notice any side effect not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Ibuprofen Gel is and what it is used for
2. Before you use Ibuprofen Gel
3. How to use Ibuprofen Gel
4. Possible side effects
5. How to store Ibuprofen Gel
6. Further Information

1. What Ibuprofen Gel is and what it is used for

Ibuprofen Gel is a pain relieving gel to be applied to the skin. It is used for the relief of

- Rheumatic pain (pain caused by problems with muscles tendons joints or bones)
- Muscular aches and pains
- Pain from strains and sprains
- Backache
- Lumbago (lower back pain)
- Fibrositis (muscle tenderness or stiffness)

Ibuprofen is one of a group of medicines called NSAIDs which provide effective pain relief and reduce both inflammation and swelling.

2. Before you use Ibuprofen Gel

Do not use Ibuprofen Gel if you:

- have an allergy to ibuprofen or any of the ingredients in Ibuprofen Gel (see end of Section 2 and Section 6)
- have an allergy to aspirin or other pain-relieving medicines known as non-steroidal anti-inflammatory drugs (NSAIDs) including those you take orally by mouth

- have asthma and know that aspirin or NSAIDs may bring on an asthma attack, a runny nose (rhinitis), itching or rashes
- are more than 6 months pregnant
- are breastfeeding
- have broken skin or your skin is infected or inflamed

Do not use Ibuprofen Gel on:

- children under 14 years old
- your lips, nostrils, eyes, genital or anal areas, or other sensitive areas. If this occurs accidentally, wash away with plenty of clean water
- the same area with other medicines that you put on the skin.

Ibuprofen Gel is only for use on the skin. Do not take by mouth.

Take special care with Ibuprofen Gel

See your doctor before using Ibuprofen Gel if you:

- have ever had allergic symptoms such as skin rashes or itching caused by medicines or cosmetics
- have a kidney problem (now or in the past)
- have a stomach ulcer (now or in the past)
- are up to 6 months pregnant or think you might be pregnant
- are having problems conceiving or having fertility treatment

Avoid getting Ibuprofen Gel on or near the eyes and other sensitive areas.

Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes any medicines you have bought without a prescription.

See your doctor before using Ibuprofen Gel if you are taking

- aspirin
- any other pain relieving medicine
- any other medicines regularly

Pregnancy and breastfeeding

See your doctor if you are pregnant or think you might be pregnant. Do not use Ibuprofen Gel if you are more than 6 months pregnant. Do not use Ibuprofen Gel if you are breast feeding.

Ask your doctor or pharmacist for advice before taking any medicine.

Important information about some of the ingredients of Ibuprofen Gel

Ibuprofen Gel contains:

- propylene glycol which may cause skin irritation

3. How to use Ibuprofen Gel

Ibuprofen Gel is for adults and children over 14 years old.

- Always try on a small area first.
- Use 1-4cm each time you use Ibuprofen Gel.
- Apply the gel as a thin layer over the affected area.
- Gently rub in the gel until it is absorbed.
- Do not use more gel than recommended.
- Repeat these steps up to 3 times a day but no more. Do not use more often than every 4 hours. If the pain or swelling continues after using the gel for **2 weeks** see your doctor.
- Unless you are treating your hands always wash your hands immediately after use.
- Replace the cap tightly.
- Stop using Ibuprofen Gel if you get excessive skin irritation or other unwanted effects

Do not cover the skin with bandages, plasters or any other dressing. However, it is safe to cover the skin with your clothes

If you use more Ibuprofen Gel than you should

If you accidentally squeeze out too much gel use a tissue to wipe off any excess.

If you or a child accidentally swallow the gel contact your doctor or hospital immediately.

If you forget to use Ibuprofen Gel

Apply Ibuprofen Gel again when you remember making sure you do not use the gel more than 3 times in one day or more often than every 4 hours.

4. Possible side effects

Like all medicines Ibuprofen Gel can cause side effects, although not everybody gets them.

Stop using Ibuprofen Gel straight away and tell your doctor if you experience any of the following effects:

- unexplained wheezing
- shortness of breath
- blisters or a rash on the skin
- itching, redness or bruising of the skin
- swelling of the face

A slight redness, drying, tingling or burning feeling may occur where you have applied the gel. This is not a cause for concern. However, if this gets worse stop using Ibuprofen Gel and tell your doctor.

Unwanted effects may be reduced if you use the smallest dose for the shortest time as possible.

More rarely you may suffer from:

- stomach pains or indigestion
- kidney problems if you have kidney disease

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of medicines.

5. How to store Ibuprofen Gel

Keep out of the sight and reach of children.

Close the cap tightly and store in a cool place (below 25°C).

Do not use after the expiry date which is printed on the tube and carton. The 'expiry' date refers to the last day of that month.

Take any Ibuprofen Gel that is out of date to a pharmacist for disposal.

6. Further information

What Ibuprofen Gel contains

The active substance is:

- ibuprofen (5 %w/w)

The other ingredients are propylene glycol, carbomer, diisopropanolamine, ethanol and water (see the end of Section 2).

What Ibuprofen Gel looks like and contents of the pack

Ibuprofen Gel is a clear gel.

Each tube contains 15, 35, 50 or 100g.

Not all pack sizes may be marketed.

Marketing authorisation holder and manufacturer:

The Mentholatum Company Limited, East Kilbride, G74 5PE, Scotland, UK.

Distributor:

The Boots Company PLC, Nottingham, NG2 3AA.

PL 00189/0024

For any information about this product please contact The Boots Company PLC,
Nottingham, NG2 3AA.

This leaflet was last revised January 2015.

Other formats

To request a copy of this leaflet in Braille, large print or audio please call, free of charge

0800 198 5000 (UK only)

Please be ready to give the following information:

Product Name: Boots Ibuprofen Gel 5% w/w Gel

Reference Number: PL 00189/0024

This is a service provided by the Royal National Institute of Blind People.

Artwork reference: BTC173600 version D