

Reads This Way



#### INFORMATION FOR THE USER

Read all of this leaflet carefully because it contains important information for you. This medicine is available without prescription. However, you still need to use Nurofen Express 684 mg Caplets carefully to get the best results from them. Nurofen Express 684 mg Caplets will be referred to as this medicine throughout this leaflet. Keep this leaflet. You may want to read it again. If you have any further questions after you have read it, ask your doctor or pharmacist.

**You must contact a doctor if your symptoms worsen or do not improve after 3 days for children and adolescents between 12 and 18 years and after 10 days for adults.**

If any side effects get serious, or if you notice any side effect not listed in this leaflet, please tell your doctor or pharmacist.

#### In this leaflet:

1. What this medicine is and what it is used for
2. Before you take this medicine
3. How to take this medicine
4. Possible side effects
5. How to store this medicine
6. Further Information

#### 1. What this medicine is and what it is used for

The active ingredient (which makes this medicine work) is Ibuprofen. It belongs to a group of medicines known as non-steroidal anti-inflammatory drugs (NSAIDs). NSAIDs provide relief by changing the body's response to pain, swelling, and high temperature.

This medicine is used for the relief of:

- Headaches and migraine pain

#### 2. Before taking this medicine

Signs of an allergic reaction to this medicine, including breathing problems, swelling of the face and neck region (angioedem aa), chest pain have been reported with ibuprofen. Immediately stop taking this medicine and contact your doctor or medical emergencies if you notice any of these signs.

**Do not take this medicine if you:**

- are allergic to ibuprofen or any of the other ingredients (see section 6) or to aspirin or other painkillers
- have (or have had two or more episodes of) a stomach ulcer, perforation or bleeding
- have had a worsening of asthma, skin rash, itchy runny nose or facial swelling when previously taking ibuprofen, aspirin or similar medicines
- have had gastrointestinal bleeding or perforation when previously taking NSAIDs (Non-steroidal anti-inflammatory drugs)
- are taking other NSAID painkillers or more than 75 mg aspirin a day
- have severe liver or kidney problems
- have heart problems, high blood pressure or blood coagulation disorder.
- have breathing difficulties
- are in the last 3 months of pregnancy
- are under 12 years old.

**Check with your pharmacist or your doctor before taking this product if you:**

- have or have had asthma
- have kidney, heart, liver or bowel problems
- have high cholesterol or previously have had a heart attack or stroke
- have a history of gastrointestinal disease (such as ulcerative colitis, Crohn's disease)
- have Systemic Lupus Erythematosus (a condition of the immune system causing joint pain skin changes and other organ disorders)
- are a smoker
- are in the first 6 months of pregnancy.
- **have an infection.** This medicine may hide signs of infections such as fever and pain. It is therefore possible that this medicine may delay appropriate treatment of infection, which may lead to an

increased risk of complications. This has been observed in pneumonia caused by bacteria and bacterial skin infections related to chickenpox. If you take this medicine while you have an infection and your symptoms of the infection persist or worsen, consult a doctor without delay.

#### Skin reactions

Serious skin reactions including exfoliative dermatitis, erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis, drug reaction with eosinophilia and systemic symptoms (DRESS), acute generalised exanthematous pustulosis (AGEP) have been reported in association with ibuprofen treatment. Stop using this medicine and seek medical attention immediately, if you notice any of the symptoms related to these serious skin reactions described in section 4.

#### Taking other medicines

To reduce the risk of side effects, do not take this product with other NSAID containing products (e.g. aspirin, ibuprofen...) This medicine may affect or be affected by some other medicines. For example:

- Avoid taking this product with corticosteroid tablets, quinolone antibiotics or drugs that are prescribed:
- as anti-coagulants (i.e. thin blood/prevent clotting e.g. aspirin/acetysalicylic acid, warfarin, ticlopidine)
- to stimulate your heart (e.g. glycosides)
- to reduce high blood pressure (ACE-inhibitors such as captopril, beta-blockers such as atenolol medicines, angiotensin-II receptor antagonists such as losartan)
- to help you passing water (diuretics)
- for the temporary suppression of your immune system (e.g. methotrexate, ciclosporin, tacrolimus)
- for mania or depression (e.g. Lithium or SSRIs)
- for pregnancy termination (e.g. mifepristone)
- for HIV treatment ( e.g. zidovudine)

**Some other medicines may also affect or be affected by the treatment of this medicine. You should therefore always seek the advice of your doctor or pharmacist before you use this medicine with other medicines.**

#### Other warnings

- This medicine belongs to a group of medicines which may **impair fertility in women.** This is reversible on stopping the medicine. It is unlikely that this medicine, used occasionally will affect your chances of becoming pregnant. However, tell your doctor before taking this medicine if you have problems becoming pregnant.
- Anti-inflammatory/pain-killer medicines such as ibuprofen may be associated with a small increased risk of heart attack or stroke, particularly when used at high doses. Do not exceed the recommended dose or duration of treatment.
- You should discuss your treatment with your doctor or pharmacist before taking this medicine if you:
  - have heart problems including heart failure, angina (chest pain), or if you have had a heart attack, bypass surgery, peripheral artery disease (poor circulation in the legs of feet due to narrow or blocked arteries), or any kind of stroke (including 'mini-stroke' or transient ischaemic attack 'TIA').
  - have high blood pressure, diabetes, high cholesterol, have a family history of heart disease or stroke, or if you are a smoker.
- There is a risk of renal impairment in dehydrated children and adolescents.
- If you are taking this medicine for longer than the recommended time or at higher than recommended doses you are at risk of serious harm. These include serious harm to the stomach/gut and kidneys, as well as very low levels of potassium in your blood. These can be fatal (see section 4).

#### Pregnancy and breast feeding

Do not take if you are in the last 3 months of pregnancy as it could harm your unborn child or cause problems at delivery. It can cause kidney and heart problems in your unborn baby. It may affect your and your baby's tendency to bleed and cause labour to be later or longer than expected. You should not take this medicine during the first 6 months of pregnancy unless absolutely necessary and advised by your doctor. If you need treatment during this period or while you are trying to get pregnant, the lowest dose for the shortest time possible should be used. If taken for more than a few days from 20 weeks of pregnancy onward, this medicine can cause kidney problems in your unborn baby that may lead to low levels of amniotic fluid that surrounds the baby (oligohydramnios) or narrowing of a blood vessel (ductus arteriosus) in the heart of the baby. If you need treatment for

longer than a few days, your doctor may recommend additional monitoring. Do not take if you are breastfeeding. Speak with your doctor if you are in the first 6 months of pregnancy. Speak to your doctor if you are in the first 6 months of pregnancy.

#### Important information about some of the ingredients of this medicine

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

#### 3. How to take this medicine

This product is for short term use only. The lowest effective dose should be used for the shortest duration necessary to relieve symptoms. If you have an infection, consult a doctor without delay if symptoms (such as fever and pain) persist or worsen (see section 2).

**Adults, the elderly and children & adolescents between 12 and 18 years:**

Take 1 caplet with water, up to three times a day as required. Leave at least four hours between doses. Do not take more than 3 caplets in 24 hours.

**Do not give to children under 12 years.**

**In children and adolescents between 12 and 18 years:**

If in children and adolescents this medicinal product is required for more than 3 days, or if symptoms worsen a doctor should be consulted.

**In adults:**

**Do not take for longer than 10 days** unless your doctor tells you to. If symptoms persist or the pain worsens, or if any new symptoms occur, consult your doctor or pharmacist.

**If you have taken more of this medicine than you should, or if children have taken this medicine by accident always contact a doctor or nearest hospital to get an opinion of the risk and advice on action to be taken.**

The symptoms can include nausea, stomach pain, vomiting (may be blood streaked), headache, ringing in the ears, confusion and shaky eye movement. At high doses, drowsiness, chest pain, palpitations, loss of consciousness, convulsions (mainly in children), weakness and dizziness, blood in urine, cold body feeling, and breathing problems have been reported.

**If you forgot to take this medicine.**

Simply refer to the directions above on how to take the medicine and do not take more than is advised.

#### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Some side effects may be minimised by taking the lowest dose for the shortest time necessary to relieve the symptoms. If any of the side effects get serious or if you notice any side effects not listed in this leaflet, tell your doctor or pharmacist.

**STOP TAKING the medicine and seek immediate medical help if you develop:**

- **signs of intestinal bleeding** such as: bright red faeces (stools/motions), black tarry stools, vomiting blood or dark particles that look like coffee grounds.
- **signs of serious allergic reaction** such as:
  - difficulties in breathing or unexplained wheezing
  - dizziness or faster heartbeat,
  - skin reactions such as reddish non-elevated, target-like or circular patches on the trunk, often with central blisters, skin peeling, ulcers of mouth, throat, nose, genitals and eyes. These serious skin rashes can be preceded by fever and flu-like symptoms (exfoliative dermatitis, erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis).
  - swelling of your face, tongue or throat
- **signs of kidney problems** such as:
  - passing less or more urine
  - cloudy urine or blood in urine
  - pain in the back and/or swelling (particularly in the legs)
- **signs of aseptic meningitis** with neck stiffness, headache, feeling sick, being sick, fever or disorientation. Patients with autoimmune disorders (lupus, mixed connective tissue disease) may be more likely to be affected.
- **a severe skin reaction known as DRESS syndrome** can occur. Symptoms of DRESS include: skin rash, fever, swelling of lymph nodes and an increase of eosinophils (a type of white blood cells).
- A red, scaly widespread rash with bumps under the skin and blisters mainly localised on the skin folds, trunk, and upper extremities accompanied by fever at the initiation of treatment (acute generalised exanthematous pustulosis) (frequency not known) See also section 2.

**STOP TAKING the medicine and tell your doctor if you experience the following uncommon side effects** which may affect up to 1 in 100 people:

- indigestion, heartburn or feeling sick
  - pains in your stomach (abdomen) or other abnormal stomach problems
- Like all medicines, this medicine can cause side-effects, although not everybody gets them. Tell your doctor or pharmacist if you notice any of the following:
- Liver, kidney problems or difficulty urinating
- This medicine, especially when taken at higher than recommended doses or for a prolonged period of time, can cause damage to your kidneys and affect them removing acids properly from your blood into the urine (renal tubular acidosis). It can also cause very low levels of potassium in your blood (see section 2). This is a very serious condition and will require immediate treatment. Signs and symptoms include muscle weakness and light-headedness.

**TELL YOUR DOCTOR if you have any of the following side effects, they become worse or you notice any effects not listed:**

**Uncommon:** may affect up to 1 in 100 people:

- allergic reactions, such as skin rashes (urticaria), itching, peeling
- headaches

**Rare:** may affect up to 1 in 1,000 people:

- flatulence (wind), diarrhoea, constipation and vomiting

**Very rare:** may affect up to 1 in 10,000 people:

- blood disorder resulting in unexplained or unusual bruising or bleeding, fever, sore throat, mouth ulcers, flu-like symptoms and severe exhaustion
- drop in blood pressure
- stomach or intestinal ulcers, sometimes with bleeding and perforation, inflammation of the lining of the mouth with ulceration (ulcerative stomatitis), inflammation of the stomach (gastritis)
- liver problems

**Not known:** frequency cannot be estimated from the available data

- worsening of asthma or bronchospasm
- swelling (oedema), high blood pressure, heart failure or attack
- chest pain, which can be a sign of a potentially serious allergic reaction called Kounis syndrome
- worsening of colitis and Crohn's disease
- skin becomes sensitive to light

Medicines such as this medicine may be associated with a small increased risk of heart attack ('myocardial infarction') or stroke. (See section 2 'Other warnings'.)

#### Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects, you can help provide more information on the safety of this medicine.

#### 5. How to store this medicine

**Keep all medicines out of the sight and reach of children.**

Do not use after the expiry date stated on the carton after EXP. The expiry date refers to the last day of that month. Do not store above 25° C. Store in the original pack.

#### 6. Further information

Each caplet contains the active ingredient Ibuprofen 400 mg (as Ibuprofen Lysine 684 mg).

Also contains: Povidone, Sodium Starch Glycolate Type A, Magnesium Stearate, Hypromellose, Talc, Opaspray White M-1-7111B (contains Hypromellose and Titanium Dioxide (E171) and Black Printing Ink (contains shellac, Iron oxide black (E172) and propylene glycol). This medicine is available in packs of 4, 6, 8, 12, 16 and 24 white, capsule-shaped tablets printed in black with identifying 'N684' logo. Not all packs will be marketed. Licence Holder: Reckitt Benckiser Healthcare (UK) Ltd, Slough, SL1 4AQ. Manufacturers: Reckitt Benckiser Healthcare International Ltd, 1 Thane Road, Nottingham NG90 2DB, UK. RB NL Brands B.V., WTC Schiphol Airport, Schiphol Boulevard 207, 1118 BH Schiphol, NL. Product licence number: PL 00063/0384 [P] Date of revision: April 2024

Reads This Way

#### CUSTOMER INFO

Minimum Point Size =9.00pt

#### RB Artwork and Print Specification

Trident Reference No:	RB605172
ZEN Ref:	<b>TR3154839</b>
Brand/Work Ref:	<b>RBL2402457</b>
Action:	<b>C</b>
Brand:	Nurofen
Category:	Adult
Segment Group:	Stronger Pain
Segment:	Express 684mg Caplets
Pack Size:	24 Pack
Market/Country:	UK
Date:	19/08/2024

RBH Contact: Rachel Beresfordward

<b>Artwork Type: BW Submission</b>	
Component Code (2D if applicable):	<b>0000000</b>
2" Component Code (if applicable):	<b>N/A</b>
Parent Technical Packaging Specification:	D0069114
Finished Goods Code:	<b>0000000</b>
Supply Point:	RB Nottingham
3rd Party Code:	N/A
Pharmacode No/NE:	N/A
Edgemark Position:	N/A

CAD Cam Ref:	N-Lit-FB-D0069114-296x192-Lge-Reel
Printer:	Generic RB
Substrate:	Paper White

Technical & Non Printing Items	
Dieline	Dieline 2 (if applicable)
Guides	Guides 2 (if applicable)

Colours	
Black	Thin




Cornought House, Cornought Road,  
Kingwood Business Park, Hull, HU7 3AP, England.  
T: +44 (0) 01482 828100

Please note that any low resolution paper Canon colour copies associated with this job should be referred to for content, layout and colour separation only.

UNDER NO CIRCUMSTANCES SHOULD THIS  
ARTWORK BE ALTERED WITHOUT PRIOR  
PERMISSION FROM TRIDENT.

STUDIO USE ONLY Subramanian MarudhuPandiyan v2.0



TDD1022/5