

## Package leaflet: Information for the patient

### Daptomycin 500 mg powder for solution for injection/infusion daptomycin

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or nurse.
- If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Daptomycin is and what it is used for
2. What you need to know before you take Daptomycin
3. How to take Daptomycin
4. Possible side effects
5. How to store Daptomycin
6. Contents of the pack and other information

#### **1. What Daptomycin is and what it is used for**

The active substance in Daptomycin powder for solution for injection or infusion is daptomycin. Daptomycin is an antibacterial that can stop the growth of certain bacteria. Daptomycin is used in adults and in children and adolescents (age from 2 to 17 years) to treat infections of the skin and the tissues below the skin. It is also used in adults and in children and adolescents (age from 1 to 17 years) to treat infections in the blood when associated with skin infection.

Daptomycin is also used in adults to treat infections in the tissues that line the inside of the heart (including heart valves) which are caused by a type of bacteria called *Staphylococcus aureus*. It is also used to treat infections in the blood caused by the same type of bacteria when associated with skin or heart infection.

Depending on the type of infection(s) that you have, your doctor may also prescribe other antibacterials while you are receiving treatment with Daptomycin.

#### **2. What you need to know before you take Daptomycin**

##### **You should not be given Daptomycin**

If you are allergic to daptomycin or to any of the other ingredients of this medicine (listed in section 6).

If this applies to you, tell your doctor or nurse. If you think you may be allergic, ask your doctor or nurse for advice.

##### **Warnings and precautions**

Talk to your doctor or nurse before you are given Daptomycin.

- If you have, or have previously had kidney problems. Your doctor may need to change the dose of Daptomycin (see section 3 of this leaflet).
- Occasionally, patients receiving daptomycin may develop tender or aching muscles or muscle weakness (see section 4 of this leaflet for more information). If this happens tell your doctor. Your doctor will make sure you have a blood test and will advise whether or

not to continue with Daptomycin. The symptoms generally go away within a few days of stopping Daptomycin.

- If you are very overweight. There is a possibility that your blood levels of daptomycin could be higher than those found in persons of average weight and you may need careful monitoring in case of side effects.

If any of these applies to you, tell your doctor or nurse before you are given Daptomycin.

**Tell your doctor straight away if you develop any of the following symptoms:**

- Serious, acute allergic reactions have been observed in patients treated with nearly all antibacterial agents, including daptomycin. Tell a doctor or a nurse straight away if you experience symptoms suggestive of allergic reaction, such as wheezing, difficulty breathing, swelling of the face, neck and throat, rashes and hives, fever (see section 4 of this leaflet for more information).
- Any unusual tingling or numbness of the hands or feet, loss of feeling or difficulties with movements. If this happens, tell your doctor who will decide whether you should continue the treatment.
- Diarrhoea, especially if you notice blood or mucus, or if diarrhoea becomes severe or persistent.
- New or worsening fever, cough or difficulty breathing. These may be signs of a rare but serious lung disorder called eosinophilic pneumonia. Your doctor will check the condition of your lungs and decide whether or not you should continue Daptomycin treatment.

Daptomycin may interfere with laboratory tests that measure how well your blood is clotting. The results can suggest poor blood clotting when, in fact, there is no problem. Therefore it is important that your doctor takes into account that you are receiving Daptomycin. Please inform your doctor that you are on treatment with Daptomycin.

Your doctor will perform blood tests to monitor the health of your muscles both before you start treatment and frequently during treatment with Daptomycin.

**Children and adolescents**

Daptomycin should not be administered to children below one year of age as studies in animals have indicated that this age group may experience severe side effects.

**Use in elderly**

People over the age of 65 can be given the same dose as other adults, provided their kidneys are working well.

**Other medicines and Daptomycin**

Tell your doctor or nurse if you are taking, have recently taken or might take any other medicines.

It is particularly important that you mention the following:

- Medicines called statins or fibrates (to lower cholesterol) or ciclosporin (a medicinal product used in transplantation to prevent organ rejection or for other conditions, e.g. rheumatoid arthritis or atopic dermatitis). It is possible that the risk of side effects affecting the muscles may be higher when any of these medicines (and some others that can affect muscles) is taken during treatment with Daptomycin. Your doctor may decide not to give you Daptomycin or to stop the other medicine for a while.
- Pain killing medicines called non-steroidal anti-inflammatory drugs (NSAIDs) or COX-2 inhibitors (e.g. celecoxib). These could interfere with the effects of Daptomycin in the kidney.
- Oral anti-coagulants (e.g. warfarin), which are medicines that prevent blood from clotting. It may be necessary for your doctor to monitor your blood clotting times.

**Pregnancy and breast-feeding**

Daptomycin is not usually given to pregnant women. If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before you are given this medicine.

Do not breast-feed if you are receiving Daptomycin, because it may pass into your breast milk and could affect the baby.

### **Driving and using machines**

Daptomycin has no known effects on the ability to drive or use machines.

## **3. How to take Daptomycin**

Daptomycin will usually be given to you by a doctor or a nurse.

### **Adults (18 years of age and above)**

The dose will depend on how much you weigh and the type of infection being treated. The usual dose for adults is 4 mg for every kilogram (kg) of body weight once daily for skin infections or 6 mg for every kg of body weight once daily for a heart infection or a blood infection associated with skin or heart infection. In adult patients, this dose is given directly into your blood stream (into a vein), either as an infusion lasting about 30 minutes or as an injection lasting about 2 minutes. The same dose is recommended in people aged over 65 years provided their kidneys are working well.

If your kidneys do not work well, you may receive Daptomycin less often, e.g. once every other day. If you are receiving dialysis, and your next dose of Daptomycin is due on a dialysis day, you will be usually given Daptomycin after the dialysis session.

### **Children and adolescents (1 to 17 years of age)**

The dose for children and adolescents will depend on the age of patient and the type of infection being treated: infections of the skin and the tissues below the skin (children and adolescents from 2 to 17 years of age) and infections in the blood when associated with skin infection (children and adolescents from 1 to 17 years of age). This dose is given directly into the blood stream (into a vein), as an infusion lasting about 30-60 minutes.

A course of treatment usually lasts for 1 to 2 weeks for skin infections. For blood or heart infections and skin infections your doctor will decide how long you should be treated.

Detailed instructions for use and handling are given at the end of the leaflet.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

The most serious side effects are described below:

### **Very rare serious side effects** (may affect up to 1 in 10,000 people)

A hypersensitivity reaction (serious allergic reaction including anaphylaxis, angioedema, drug rash with eosinophilia and systemic symptoms (DRESS)) has been reported, in some cases during administration of daptomycin. This serious allergic reaction needs immediate medical attention. Tell your doctor or nurse straight away if you experience any of the following symptoms:

- Chest pain or tightness,
- Rash with blistering, sometimes affecting the mouth and genitals,

- Swelling around throat,
- Rapid or weak pulse,
- Wheezing,
- Fever,
- Shivering or trembling,
- Hot flushes,
- Dizziness,
- Fainting,
- Metallic taste.

Tell your doctor straight away if you experience unexplained muscle pain, tenderness, or weakness. In very rare cases (reported in less than 1 in every 10,000 patients), muscle problems can be serious, including muscle breakdown (rhabdomyolysis), which can result in kidney damage.

**Serious side effects with frequency not known** (frequency cannot be estimated from the available data)

A rare but potentially serious lung disorder called eosinophilic pneumonia has been reported in patients given daptomycin, mostly after more than 2 weeks of treatment. The symptoms can include difficulty breathing, new or worsening cough, or new or worsening fever. If you experience these symptoms, tell your doctor or nurse straight away. If you experience raised or fluid-filled skin spots over a large area of your body, tell your doctor or nurse straight away.

The most frequently reported side effects are described below:

**Common side effects** (may affect up to 1 in 10 people)

- Fungal infections such as thrush,
- Urinary tract infection,
- Decreased number of red blood cells (anaemia),
- Dizziness, anxiety, difficulty in sleeping,
- Headache,
- Fever, weakness (asthenia),
- High or low blood pressure,
- Constipation, abdominal pain,
- Diarrhoea, feeling sick (nausea) or being sick (vomiting),
- Flatulence,
- Abdominal swelling or bloating,
- Skin rash or itching,
- Pain, itchiness or redness at the site of infusion,
- Pain in arms or legs,
- Blood testing showing higher levels of liver enzymes or creatine phosphokinase (CPK).

Other side effects which may occur following daptomycin treatment are described below:

**Uncommon side effects** (may affect up to 1 in 100 people)

- Blood disorders (e.g. increased number of small blood particles called platelets, which may increase the tendency for blood clotting, or higher levels of certain types of white blood cells),
- Decreased appetite,
- Tingling or numbness of the hands or feet, taste disturbance,
- Trembling,
- Changes in heart rhythm, flushes,
- Indigestion (dyspepsia), inflammation of the tongue,
- Itchy rash of skin,
- Muscle pain, cramping, or weakness, inflammation of the muscles (myositis), joint pain,

- Kidney problems,
- Inflammation and irritation of the vagina,
- General pain or weakness, tiredness (fatigue),
- Blood test showing increased levels of blood sugar, serum creatinine, myoglobin, or lactate dehydrogenase (LDH), prolonged blood clotting time or imbalance of salts,
- Itchy eyes.

**Rare side effects** (may affect up to 1 in 1,000 people)

- Yellowing of the skin and eyes,
- Prothrombin time prolonged.

**Frequency not known** (frequency cannot be estimated from the available data)

Antibacterial-associated colitis, including pseudomembranous colitis (severe or persistent diarrhoea containing blood and/or mucus, associated with abdominal pain or fever), easy bruising, bleeding gums, or nosebleeds..

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via Yellow Card Scheme Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Daptomycin**

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and label after EXP. The expiry date refers to the last day of the month.
- Store in a refrigerator (2°C – 8°C).

After reconstitution:

Chemical and physical in-use stability of the reconstituted solution in the vial has been demonstrated for 12 hours at 25°C and up to 48 hours at 2°C – 8°C.

After dilution:

Chemical and physical stability of the diluted solution in infusion bags is established as 12 hours at 25°C or 24 hours at 2°C – 8°C.

For the 30-minute intravenous infusion, the combined storage time (reconstituted solution in vial and diluted solution in infusion bag; see section 6.6) at 25°C must not exceed 12 hours (or 24 at 2°C – 8°C).

For the 2-minute intravenous injection, the storage time of the reconstituted solution in the vial (see section 6.6) at 25°C must not exceed 12 hours (or 48 at 2°C – 8°C).

From a microbiological point of view the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C – 8°C.

Do not use Daptomycin if you notice any change in the aspect of the product (signs of humidity or presence of particles with a different colour in the powder, or presence of particles, turbidity or precipitate when the solution is reconstituted).

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## **6. Contents of the pack and other information**

### **What Daptomycin contains**

The active substance is daptomycin. One vial of powder contains 500 mg daptomycin. One ml provides 50 mg of daptomycin after reconstitution with 10 ml of sodium chloride 9 mg/ml (0.9%) solution.

- The other ingredient is sodium hydroxide.

### **What Daptomycin looks like and contents of the pack**

Daptomycin powder for solution for injection or infusion is supplied as a pale yellow to light brown cake or powder in a glass vial. It is mixed with a solvent to form a liquid before it is administered.

Daptomycin is available in packs containing 1 vial or 5 vials.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder**

Accord Healthcare Limited  
Sage House, 319 Pinner Road  
North Harrow, Middlesex, HA1 4HF  
United Kingdom

### **Manufacturer**

Medichem, S.A.  
Mossen Jacint Verdaguer, 67A  
08970 Sant Joan Despí  
(Barcelona)  
Spain

Hikma Italia SpA  
10 Viale Certosa  
Pavia 27100  
Italy

**This leaflet was last revised in 09/2018.**

**The following information is intended for healthcare professionals only**

Important: Please refer to the Summary of Product Characteristics before prescribing.

Instructions for use and handling

500 mg presentation:

In adults, daptomycin may be administered intravenously as an infusion over 30 minutes or as an injection over 2 minutes. Unlike in adults, daptomycin should not be administered by injection over a 2 minute period in paediatric patients. Paediatric patients 7 to 17 years old should receive daptomycin infused over 30 minutes. In paediatric patients under 7 years old receiving a 9-12mg/kg dose, daptomycin should be administered over 60 minutes. Preparation of the solution for infusion requires an additional dilution step as detailed below.

**Daptomycin given as an intravenous infusion over 30 or 60 minutes**

A 50 mg/ml concentration of Daptomycin can be achieved by reconstituting the lyophilised product with 10 ml of sodium chloride 9 mg/ml (0.9%) solution for injection.

The lyophilised product takes approximately 15 minutes to dissolve. The fully reconstituted product will appear clear and may have a few small bubbles or foam around the edge of the vial.

To prepare Daptomycin for intravenous infusion, please adhere to the following instructions: Aseptic technique should be used throughout to reconstitute lyophilised Daptomycin.

1. The flip off cap should be removed to expose the central portions of the rubber stopper. Wipe the top of the rubber stopper with an alcohol swab or other antiseptic solution and allow to dry. After cleaning, do not touch the rubber stopper or allow it to touch any other surface. Draw 10 ml of sodium chloride 9 mg/ml (0.9%) solution for injection into a syringe using a sterile transfer needle that is 21 gauge or smaller in diameter, or a needleless device, then slowly inject through the centre of the rubber stopper into the vial pointing the needle towards the wall of the vial.
2. The vial should be gently rotated to ensure complete wetting of the product and then allowed to stand for 10 minutes.
3. Finally the vial should be gently rotated/swirled for a few minutes as needed to obtain a clear reconstituted solution. Vigorous shaking/agitation should be avoided to prevent foaming of the product.
4. The reconstituted solution should be checked carefully to ensure that the product is in solution and visually inspected for the absence of particulates prior to use. Reconstituted solutions of Daptomycin range in colour from pale yellow to light brown.
5. Slowly remove the reconstituted liquid (50 mg daptomycin/ml) from the vial using a sterile needle that is 21 gauge or smaller in diameter.
6. The reconstituted solution should then be diluted with sodium chloride 9 mg/ml (0.9%) (typical volume 50 ml).
7. Invert the vial of the sodium chloride solution in order to allow the solution to drain towards the stopper. Using a new syringe, insert the needle into the inverted vial. Keeping the vial inverted, position the needle tip at the very bottom of the solution in the vial when drawing the solution into the syringe. Before removing the needle from the vial, pull the plunger all the way back to the end of the syringe barrel in order to remove all of the solution from the inverted vial.
8. Replace needle with a new needle for the intravenous infusion.
9. Expel air, large bubbles, and any excess solution in order to obtain the required dose.

10. The reconstituted and diluted solution should then be infused intravenously over 30 or 60 minutes.

Daptomycin is not physically or chemically compatible with glucose-containing solutions. The following have been shown to be compatible when added to daptomycin containing infusion solutions: aztreonam, ceftazidime, ceftriaxone, gentamicin, fluconazole, levofloxacin, dopamine, heparin and lidocaine.

The combined storage time (reconstituted solution in vial and diluted solution in infusion bag) at 25°C must not exceed 12 hours (24 hours if refrigerated).

Stability of the diluted solution in infusion bags is established as 12 hours at 25°C or 24 hours if stored under refrigeration at 2°C – 8°C.

### **Daptomycin given as 2-minute intravenous injection**

Water should not be used for reconstitution of Daptomycin for intravenous injection. Daptomycin should only be reconstituted with sodium chloride 9 mg/ml (0.9%).

A 50 mg/ml concentration of Daptomycin for injection is obtained by reconstituting the lyophilised product with 10 ml of sodium chloride 9 mg/ml (0.9%) solution for injection.

The lyophilised product takes approximately 15 minutes to dissolve. The fully reconstituted product will appear clear and may have a few small bubbles or foam around the edge of the vial.

To prepare Daptomycin for intravenous injection, please adhere to the following instructions: Aseptic technique should be used throughout to reconstitute lyophilised Daptomycin.

1. The flip off cap should be removed to expose the central portions of the rubber stopper. Wipe the top of the rubber stopper with an alcohol swab or other antiseptic solution and allow to dry. After cleaning, do not touch the rubber stopper or allow it to touch any other surface. Draw 10 ml of sodium chloride 9 mg/ml (0.9%) solution for injection into a syringe using a sterile transfer needle that is 21 gauge or smaller diameter, or a needleless device, then slowly inject through the centre of the rubber stopper into the vial pointing the needle towards the wall of the vial.
2. The vial should be gently rotated to ensure complete wetting of the product and then allowed to stand for 10 minutes.
3. Finally the vial should be gently rotated/swirled for a few minutes as needed to obtain a clear reconstituted solution. Vigorous shaking/agitation should be avoided to prevent foaming of the product.
4. The reconstituted solution should be checked carefully to ensure that the product is in solution and visually inspected for the absence of particulates prior to use. Reconstituted solutions of Daptomycin range in colour from pale yellow to light brown.
5. Slowly remove the reconstituted liquid (50 mg daptomycin/ml) from the vial using a sterile needle that is 21 gauge or smaller in diameter.
6. Invert the vial in order to allow the solution to drain towards the stopper. Using a new syringe, insert the needle into the inverted vial. Keeping the vial inverted, position the needle tip at the very bottom of the solution in the vial when drawing the solution into the syringe. Before removing the needle from the vial, pull the plunger all the way back to the end of the syringe barrel in order to remove all of the solution from the inverted vial.
7. Replace needle with a new needle for the intravenous injection.
8. Expel air, large bubbles, and any excess solution in order to obtain the required dose.
9. The reconstituted solution should then be injected intravenously slowly over 2 minutes.

Chemical and physical in-use stability on the reconstituted solution in the vial has been demonstrated for 12 hours at 25°C and up to 48 hours if stored under refrigeration (2°C – 8°C).



From a microbiological point of view the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than the times stated above for the chemical and physical in-use stability.

This medicinal product must not be mixed with other medicinal products except those mentioned above.

Daptomycin vials are for single-use only. Any unused portion remaining in the vial should be discarded.