

Package leaflet: Information for the patient

Eplerenone 25 mg film-coated tablets

Eplerenone 50 mg film-coated tablets

eplerenone

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Eplerenone is and what it is used for
2. What you need to know before you take Eplerenone
3. How to take Eplerenone
4. Possible side effects
5. How to store Eplerenone
6. Contents of the pack and other information

1. What Eplerenone is and what it is used for

Eplerenone belongs to a group of medicines known as selective aldosterone blocking agents. These blocking agents inhibit the action of aldosterone, a substance produced within the body, which controls your blood pressure and heart function. High levels of aldosterone can cause changes in your body that lead to heart failure.

Eplerenone is used to treat your heart failure to prevent worsening and reduce hospitalisations if you have:

1. had a recent heart attack, in combination with other drugs that are used to treat your heart failure, or
2. have persistent, mild symptoms despite the treatment you have been receiving so far.

2. What you need to know before you take Eplerenone

Do not take Eplerenone:

- if you are allergic to eplerenone or any of the other ingredients of this medicine (listed in section 6)
- if you have high levels of potassium in your blood (hyperkalemia)
- if you are taking groups of drugs which help you to excrete excessive body fluid (potassium sparing diuretics)
- if you have severe kidney disease
- if you have severe liver disease
- if you are taking medicines that are used to treat fungal infection (ketoconazole or itraconazole)
- if you are taking antiviral medication for treating HIV (nelfinavir or ritonavir)
- if you are taking antibiotics used to treat bacterial infections (clarithromycin or telithromycin)
- if you are taking nefazodone used to treat depression
- if you are taking medicines used to treat certain heart conditions or hypertension (so called angiotensin converting enzyme (ACE) inhibitor and an angiotensin receptor blocker (ARB)) together.

Warnings and precautions

Talk to your doctor or pharmacist before taking Eplerenone:

- if you have kidney or liver disease (see also “Do not take Eplerenone”)
- if you are taking lithium (usually given for manic depressive disorder, also called bipolar disorder)
- if you are taking tacrolimus or ciclosporin (used to treat skin conditions such as psoriasis or eczema, and to prevent rejection after organ transplantation)

Children and adolescents

The safety and efficacy of eplerenone in children and adolescents have not been established.

Other medicines and Eplerenone

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

You must not take Eplerenone with the following medications (see section “Do not take Eplerenone”):

- Itraconazole or ketoconazole (used to treat fungal infections), ritonavir, nelfinavir (antiviral medication for treating HIV), clarithromycin, telithromycin (used to treat bacterial infections) or nefazodone (used to treat depression) as these drugs reduce the break-down of Eplerenone, thereby prolonging its effect on the body.
- Potassium sparing diuretics (drugs which help you to excrete excess body fluid) and potassium supplements (salt tablets) as these drugs increase the risk of high potassium levels in your blood.
- Angiotensin converting enzyme (ACE) inhibitors and angiotensin receptor blockers (ARB) together (which are used to treat high blood pressure, heart disease or particular kidney conditions) as these drugs may increase the risk of high potassium levels in your blood.

Please inform your doctor if you are taking any of the following medicines:

- Lithium (usually given for manic depressive disorder, also called bipolar disorder). Use of lithium together with diuretics and ACE inhibitors (used to treat high blood pressure and heart disease) has been shown to cause levels of lithium in the blood to become too high, which may cause side effects of: loss of appetite; visual impairment; tiredness; muscle weakness; muscle twitches.
- Ciclosporin or tacrolimus (used to treat skin conditions such as psoriasis or eczema, and to prevent rejection after organ transplantation). These drugs can cause kidney problems and therefore increase the risk of high potassium levels in your blood.
- Non-steroidal anti-inflammatory drugs (NSAIDs - certain pain killers such as ibuprofen, used to relieve pain, stiffness and inflammation). These drugs may lead to kidney problems and therefore increase the risk of high potassium levels in your blood.
- Trimethoprim (used to treat bacterial infections) may increase the risk of high potassium levels in your blood.
- Alpha I blockers, such as prazosin or alfuzosin (used to treat high blood pressure and particular prostate conditions) may lead to a fall in blood pressure and dizziness upon standing.
- Tricyclic antidepressants such as amitriptyline or amoxapine (for treatment of depressions), antipsychotics (also known as neuroleptics) such as chlorpromazine or haloperidol (for the treatment of psychiatric disorders), amifostine (used during cancer chemotherapy) and baclofen (used to treat muscle spasm). These drugs may lead to a fall in blood pressure and dizziness upon standing.
- Glucocorticoids, such as hydrocortisone or prednisone (used to treat inflammation and certain skin conditions) and tetracosactide (mainly used for diagnosing and treating disorders of the adrenal cortex) may reduce the blood pressure lowering effect of Eplerenone.
- Digoxin (used in the treatment of heart conditions). Digoxin blood levels may be increased when taken together with Eplerenone.
- Warfarin (an anti-clotting drug): Caution is warranted when taking warfarin because high levels of warfarin in the blood may cause changes in the effect of Eplerenone on the body.
- Erythromycin (used to treat bacterial infections), saquinavir (antiviral medication for treating HIV), fluconazole (used to treat fungal infections), amiodarone, diltiazem and verapamil (for the treatment of heart problems and high blood pressure) reduce the break-down of Eplerenone thereby prolonging the effect of Eplerenone on the body (see section 3 ‘How to take Eplerenone’).
- St John’s Wort (herbal medicinal product), rifampicin (used to treat bacterial infections), carbamazepine, phenytoin, and phenobarbital (used, among others, to treat epilepsy) may increase the break-down of Eplerenone and thus decrease its effect.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. The effect of Eplerenone has not been evaluated during pregnancy in humans.

It is not known if eplerenone is excreted in human breast milk. A decision should be made with your doctor, whether to discontinue breast-feeding or to discontinue the drug.

Driving and using machines

You may feel dizzy after taking Eplerenone. If this should happen, do not drive or operate machinery.

Eplerenone Tablets contain lactose and sodium

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

3. How to take Eplerenone

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Eplerenone is usually administered together with other medication for heart failure e.g. beta blockers. The recommended starting dose is one 25 mg tablet once daily, increasing after about 4 weeks to 50 mg once daily (either as one 50 mg tablet or two 25 mg tablets). The maximum dose regimen is 50 mg daily.

Blood potassium levels should be measured before starting Eplerenone therapy, within the first week and at one month after the start of treatment or after a change in dose. The dose may be adjusted by your doctor, depending on the potassium levels in your blood.

If you have mild kidney disease, you should start on one 25 mg tablet every day. And if you have moderate kidney disease, you should start on one 25 mg tablet every other day. These doses may be adjusted if your doctor recommends and according to your blood potassium levels.

In patients with severe kidney disease, Eplerenone is not recommended.

In patients with mild-to-moderate liver disease no adjustment of the starting dose is required. If you have liver or kidney problems, you may need more frequent testing of your blood potassium levels (see also "Do not take Eplerenone").

If you take some other medicines, your doctor may recommend you take a lower dose.

For the elderly: no adjustment of the starting dose is required.

For children and adolescents: Eplerenone is not recommended.

Taking your medicine

Eplerenone tablets may be taken together with food or on an empty stomach. Swallow the tablets whole with plenty of water.

If you take more Eplerenone than you should

If you take more Eplerenone than you should, tell your doctor or pharmacist immediately. If you have taken too much of your medicine, the most likely symptoms will be low blood pressure (expressed as a light feeling in your head, dizziness, blurred vision, weakness, acute loss of consciousness) or hyperkalemia, high levels of potassium in the blood (expressed by muscle cramps, diarrhoea, nausea, dizziness or headache).

If you forget to take Eplerenone

If it is almost time for your next tablet, skip the tablet you missed and take your next tablet when it is due.

Otherwise take the tablet as soon as you remember, providing there is more than 12 hours to when you are due to take your next tablet. Then go back to taking your medicine as you would normally. Do not take a double dose to make up for the forgotten tablet.

If you stop taking Eplerenone

It is important to keep taking Eplerenone as prescribed unless your doctor tells you to stop your treatment.

If you have any further questions on the use of this medicine ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If you experience any of the following **you should seek immediate medical attention:**

Common side-effects (*may affect up to 1 in 10 people*):

- heart complaints e.g. irregular heart beat and heart failure
- abnormal functioning of your kidney – you may notice little or no urine being passed, or have lower back pain.

Uncommon side-effects (*may affect up to 1 in 100 people*):

- inflammation of the gall bladder – you may have sudden sharp pain in the stomach that may spread towards the shoulders
- thrombosis (blood clot) in the leg
- kidney inflammation – you may notice lower back pain, cloudy urine or blood in the urine
- swollen face, tongue or throat, difficulty swallowing, hives and difficulties breathing. These are the symptoms of angioneurotic oedema.

Other reported side effects include:

Common side-effects (*may affect up to 1 in 10 people*):

- elevated potassium level in your blood (symptoms include muscle cramps, diarrhoea, nausea, dizziness or headache)
- dizziness
- fainting
- infection
- cough
- constipation
- low blood pressure
- diarrhoea
- feeling sick (nausea) or being sick (vomiting)
- headache
- difficulty sleeping (insomnia)
- rash
- itching
- muscle spasm and pain
- back pain
- increased urea level in the blood
- feeling unusually weak
- elevated quantity of cholesterol or triglycerides (fats) in your blood
- increased creatinine blood levels which may indicate kidney problems.

Uncommon side-effects (*may affect up to 1 in 100 people*):

- eosinophilia (increase in certain white blood cells)
- dehydration
- low sodium blood levels
- fast heart beat
- decreased blood pressure that can cause dizziness upon standing

- sore throat
- flatulence
- underactive thyroid
- increase in blood glucose
- reduced sense of touch
- increased sweating
- feeling unwell
- enlargement of breasts in men
- changes in some blood test results

Reporting of side effects

If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Eplerenone

Keep this medicine out of the sight and reach of children.

This medicine does not require any special storage conditions.

Do not use this medicine after the expiry date which is stated on the carton, bottle or blister after EXP. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Eplerenone contains

- The active substance is eplerenone. Each film-coated tablet contains 25 mg or 50 mg eplerenone.
- The other ingredients are:

Tablet Core

Lactose monohydrate microcrystalline cellulose, croscarmellose sodium, hypromellose, sodium laurilsulfate, talc, magnesium stearate (see section 2, 'Eplerenone Tablets contain lactose and sodium')

Film Coating

Hypromellose, titanium dioxide (E171), macrogol, yellow iron oxide (E172), red iron oxide (E172), polysorbate

What Eplerenone looks like and contents of the pack

Eplerenone 25 mg film-coated tablets are yellow, film-coated, round, biconvex tablets, marked with 'EP1' on one side and 'M' on other side.

Eplerenone 50 mg film-coated tablets are yellow, film-coated, round, biconvex tablets, marked with 'EP2' on one side and 'M' on other side.

Eplerenone tablets are available in blister packs containing 20, 28, 30, 50, 90 and 100 tablets and single-unit dose blisters in packs containing 30 x 1, 50 x 1 and 90 x 1 tablets, or plastic bottles containing 28, 30, 90 and 250 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

Mylan, Potters Bar, Hertfordshire, EN6 1TL, United Kingdom.

Manufacturers

Gerard Laboratories, 35/36 Baldoyle Industrial Estate, Grange Road, Dublin 13, Ireland

Mylan Hungary Kft., H-2900 Komárom Mylan útca 1, Hungary

This leaflet was last revised in 12/2020.