

**PACKAGE LEAFLET**

## PACKAGE LEAFLET: INFORMATION FOR THE USER

NALOXONE 400 microgram/ml solution for injection or infusion

naloxone hydrochloride

### **Read all of this leaflet carefully before you start using this medicine.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

### **In this leaflet:**

1. What NALOXONE is and what it is used for
2. Before you use NALOXONE
3. How to use NALOXONE
4. Possible side effects
5. How to store NALOXONE
6. Further information

## **1. WHAT NALOXONE IS AND WHAT IT IS USED FOR**

NALOXONE is a drug used to counter the effects of opioid overdose, for example morphine overdose. NALOXONE is used for reversal of unwanted effects of opioids for countering life-threatening depression of the central nervous system and respiratory system (breathing difficulties). NALOXONE is also used to diagnose an acute opioid overdose or intoxication.

## **2. BEFORE YOU USE NALOXONE**

### **Do not use NALOXONE**

if you are allergic (hypersensitive) to naloxone hydrochloride or any of the other ingredients of NALOXONE.

### **Take special care with NALOXONE**

- if you are physically dependent to morphine or similar drugs or when you have received high doses of these drugs, you may develop withdrawal symptoms like high blood pressure, rapid heartbeat, serious respiratory problems or stop of the heartbeat.
- if NALOXONE must be administered to your newborn baby, as acute withdrawal symptoms can occur.
- if you have cardiovascular complaints (because side effects like high and low blood pressure, rapid heartbeat or serious respiratory problems probably can occur sooner).
- if you take the analgesic drug buprenorphine. In that case naloxone is effective to a limited extent (see also the paragraph ‘Taking other medicines’).

*Please consult your doctor even if these statements were applicable to you at any time in the past.*

### **Taking other medicines**

Please tell your doctor if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

Please note that these statements may also apply to products used some time ago or at some time in the future.

- If you use analgesics, such as buprenorphine, the analgesic effects can be increased when you are treated with NALOXONE.
- An administration of NALOXONE in coma as a consequence of clonidine-overdose, serious high blood pressure has been reported. Clonidine is a medicine used in withdrawal symptoms occurring after stopping opioids. It is also administered in high blood pressure, migraine and menopausal flushes.

### **Using NALOXONE with food and drink**

Please inform your doctor if you drank alcohol. In patients with multiple intoxication (with opioids and sedatives or alcohol) NALOXONE onset of effect can be less rapid.

### **Pregnancy**

There are no adequate data available on the use of NALOXONE in pregnant women. During pregnancy, your doctor will outweigh the benefits of the use of NALOXONE against the possible risks for the unborn baby. NALOXONE can cause withdrawal symptoms in the baby (see paragraph Take special care with...).

*Ask your doctor or pharmacist for advice before taking any medicine.*

### **Breast-feeding**

It is not known whether NALOXONE passes into breast milk and it has not been established whether infants who are breast-fed are affected by NALOXONE. Therefore, breast-feeding is not recommended for 24 hours after treatment.

*Ask your doctor or pharmacist for advice before taking any medicine.*

### **Driving and using machines**

After receiving NALOXONE for the reversal of the effects of opioids you must not take part in road traffic, operate machinery or engage in any other physically or mentally demanding activity for at least 24 hours since the effects of opioids may possibly recur.

### **Important information about some of the ingredients of NALOXONE**

This medicinal product contains 17.7 mg sodium per 2 mg dose (5 ml) of naloxone hydrochloride. To be taken into consideration by patients on a controlled sodium diet.

## **3. HOW TO USE NALOXONE**

### **Dose**

Your doctor will determine the right dose.

#### Adults

- Overdose of opioids: 400 microgram. If needed, the dose can be repeated at 2-3 minute intervals.
- Reducing the effect of opioids used in anaesthesia for an operation: 100–200 microgram at 2-3 minute intervals.

#### Children and adolescents

- Overdose of opioids: 10–20 microgram/kg body weight. If needed, the dose can be repeated at 2-3 minute intervals.

#### Neonates

- Decreased respiration caused by opioids: 10 microgram/kg body weight. If needed, the dose can be repeated at 2-3 minute intervals.

#### Elderly

In elderly patients with heart diseases NALOXONE must be used with caution.

### **Method of administration**

NALOXONE is administered as an injection. It is injected into a vein (=intravenous) or into a muscle (=intramuscular) by a doctor or a nurse.

It also can be given as intravenous infusion after dilution with sodium chloride 0.9% or glucose 5%.

### **Duration of treatment**

Your doctor will determine the duration of treatment.

*If you have the impression that the effect of NALOXONE is too strong or too weak, talk to your doctor or pharmacist.*

### **If you use more NALOXONE than you should**

If you may have received more NALOXONE than you should, talk to your doctor or nurse immediately. He/she will take further measures, if necessary.

*If you have any further questions on the use of this product, ask your doctor or pharmacist.*

## **4. POSSIBLE SIDE EFFECTS**

Like all medicines, NALOXONE can cause side effects, although not everybody gets them.

The possible side effects are ranked according to their frequency:

*very common: affects more than 1 user in 10*

*common: affects 1 to 10 users in 100*

*uncommon: affects 1 to 10 users in 1,000*

*rare: affects 1 to 10 users in 10,000*

*very rare: affects less than 1 user in 10,000*

*not known: frequency cannot be estimated from the available data*

The following side effects may appear:

Very common: sickness, feeling of being sick

Common: dizziness, headache, rapid heartbeat, low blood pressure, high blood pressure, vomiting, postoperative pain

Uncommon: shiver, sweating, heart rhythm disturbance, slow heartbeat, diarrhoea, dry mouth, rapid and deep breathing (hyperventilation), irritation of vessel wall (after intravenous administration)

Rare: fits, tension

Very rare: rapid and irregular heartbeat, stop of heartbeat, fluid accumulation in the lungs, allergic reactions (urticaria, rhinitis, respiratory difficulties, anaphylactic shock)

When NALOXONE is administered to persons addicted to morphine or similar drugs, acute withdrawal symptoms can occur (for example high blood pressure and heart symptoms). This can also occur in babies of opioid-dependent mothers.

If a too large dose is given after an operation, you may become excited and feel pain (because the painkilling effects of the medicines you were given will have been counteracted as well as the effects on your breathing).

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

## 5. HOW TO STORE NALOXONE

Keep out of reach and sight of children.

Do not use NALOXONE after the expiry date which is stated on the carton and ampoule after “exp”. The expiry date refers to the last day of that month.

Do not store above 25°C. Keep the ampoules in the outer carton in order to protect from light.

After first opening the medicinal product must be used immediately.

After dilution, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8°C, unless dilution has taken place under controlled and validated aseptic conditions.

This medicinal product is for single use only. Discard any unused solution.

Do not use NALOXONE if you notice a discolouration, cloudiness or particles in the solution.

For i.v. infusion, NALOXONE 0.4 mg/ml is diluted with sodium chloride 0.9% w/v or glucose 5% w/v. 5 ampoules of NALOXONE 0.4 mg/ml (2 mg) diluted to 500 ml give a final concentration of 4 microgram/ml.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## 6. FURTHER INFORMATION

### What NALOXONE contains

The active substance is naloxone hydrochloride.

Each ampoule of 1 ml of solution for injection or solution contains 400 microgram/ml naloxone hydrochloride (as naloxone hydrochloride dihydrate).

The other ingredients are sodium chloride, hydrochloric acid (diluted) and water for injections.

### What NALOXONE looks like and contents of the pack

NALOXONE is a clear and colourless solution for injection or solution.

NALOXONE is available in packs with 10 ampoules of 1 ml solution of injection or solution.

### Marketing Authorisation Holder

Orpha-Devel Handels und Vertriebs GmbH  
3002 Purkersdorf, Austria

### Manufacturer

Hikma Italia S.p.A.  
27100 Pavia, Italy

G.L. Pharma GmbH  
A-1160 Vienna, Austria

Amomed Pharma GmbH  
A-1150 Vienna, Austria

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Bulgaria, Nexodal 0,4 mg/ml инжекционен разтвор или инфузия  
Denmark, Nexodal 0,4 mg/ml injektions-/infusionsvæske, opløsning  
Estonia, Nexodal 0,4 mg/ml süste-/infusioonilahus  
Finland, Nexodal 0,4 mg/ml injektio-/infusioneste, liuos Germany, Nexodal 0,4 mg/ml Injektionslösung oder Infusionslösung  
Greece, Naloxon Orpha 0,4 mg/ml διάλυμα για ένεση/έγχυση  
Hungary, Nexodal 0,4 mg/ml oldatos injekció vagy infúzió  
Ireland, Naloxone 400 microgram/ml solution for injection or infusion  
Italy, Nexodal 0,4 mg/ml soluzione iniettabile  
Latvia, NEXODAL 0,4 mg/ml šķīdums injekcijām vai infūzijām  
Lithuania, NEXODAL 0,4 mg/ml injekcinis/infuzinis tirpalas  
Netherlands, Naloxon Orpha 0,4 mg/ml oplossing voor injectie of infusie  
Norway, Nexodal 0,4 mg/ml injeksjons-/infusjonsvæske, oppløsning  
Poland, Nexodal 0,4 mg/ml roztwór do wstrzykiwań lub infuzji  
Romania, Nexodal 0,4 mg/ml soluție injectabilă sau perfuzabilă  
Slovenia, Nexodal 0,4 mg/ml raztopina za injiciranje ali infundiranje  
Sweden, Nexodal 0,4 mg/ml injektions-/infusionsvätska, lösning  
United Kingdom, Naloxone 400 microgram/ml solution for injection or infusion

**This leaflet was last approved in 05/2017**