

Product Name Slow Sodium Tablets
 MA Number PL 44490/0004
 Component Patient Leaflet
 Artwork Number PL-SOD-TAB-600 Rev 00

Second Part of Fold-Out Package Leaflet




<p>If you forget to take Slow Sodium tablets If you miss a dose, take one as soon as you can. If you have missed several doses tell your doctor. Do not take a double dose to make up for a forgotten dose.</p> <p>If you have any further questions on the use of this medicine, ask your doctor or pharmacist.</p> <p>4. POSSIBLE SIDE EFFECTS No side effects have been reported at the recommended doses.</p> <p>Reporting of side effects If you get any side effects talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the yellow card</p>	<p>scheme at www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.</p> <p>5. HOW TO STORE SLOW SODIUM TABLETS Keep this medicine out of the sight and reach of children. Do not use this medicine after the expiry date which is stated on the label after EXP. The expiry date refers to the last day of that month. Do not store above 30°C. Store in the original bottle in order to protect from moisture.</p> <p>Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.</p>	<p>6. CONTENTS OF THE PACK AND OTHER INFORMATION</p> <p>What Slow Sodium Tablets contain The active substance is sodium chloride. Each tablet contains the equivalent of 600mg of sodium chloride.</p> <p>The other ingredients are: cetostearyl alcohol, gelatin, magnesium stearate, hypromellose (E464), hydroxypropyl cellulose (E463), talc and titanium dioxide (E171).</p> <p>What Slow Sodium tablets look like and contents of the pack The tablets are round, white and polished. They are available in containers of 100 tablets.</p>	<p>Marketing Authorisation Holder ALTURIX Ltd. 287 Upper Fourth Street, Milton Keynes, MK9 1EH, U.K.</p> <p>Manufacturer Elara Pharmaceuticals Limited, Iron Farm, 7 Grimesgate Diseworth DE74 2QD United Kingdom</p> <p>This leaflet was last revised in April 2023. PL 44490/0004</p>	<p>Slow Sodium Tablets® 100 tablets</p> <p>For the treatment and prevention of sodium chloride deficiency. Read the package leaflet before use.</p> <p>Dose Adults: To prevent sodium chloride deficiency: 4-8 tablets per day. For cases of severe salt loss: Up to 20 tablets per day. To control muscle cramps during haemodialysis: 10-16 tablets. Children: As directed by a doctor.</p> <p>Each tablet should be swallowed whole with approx. 70ml of water and not chewed.</p> <p>Keep out of the sight and reach of children.</p> <p>Store in the original bottle in order to protect from moisture. Store below 30°C.</p> <p>Each tablet contains 600mg sodium chloride equivalent to approx. 10mEq of Na+ (230mg Na) in a sustained release formulation.</p>	 <p>PL 44490/0004 ALTURIX Ltd 287 Upper Fourth Street, Milton Keynes, MK9 1EH</p>  <p>5 036376 224646 1401689 a</p> <p>Peel here but do not remove</p>
--	---	---	--	---	---

Outer Bottle Label (Peeled back to reveal Package Leaflet)




First Part of Fold-Out Package Leaflet

<p>PACKAGE LEAFLET: INFORMATION FOR THE USER</p> <p>SLOW SODIUM TABLETS</p> <p>Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.</p> <p>Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist has told you.</p> <ul style="list-style-type: none"> Keep this leaflet. You may need to read it again. Ask your pharmacist if you need more information or advice. If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4. 	<p>What is in this leaflet:</p> <ol style="list-style-type: none"> What Slow Sodium tablets are and what they are used for What you need to know before you take Slow Sodium tablets How to take Slow Sodium tablets Possible side effects How to store Slow Sodium tablets Contents of the pack and other information <p>1. WHAT SLOW SODIUM TABLETS ARE AND WHAT THEY ARE USED FOR</p> <p>Slow Sodium tablets contain the active substance sodium chloride (salt) in a special slow-release form. Slow Sodium tablets are used for the treatment and prevention of sodium chloride deficiency.</p>	<p>2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE SLOW SODIUM TABLETS</p> <p>Do not take Slow Sodium tablets if you:</p> <ul style="list-style-type: none"> are allergic to sodium chloride or any of the ingredients of Slow Sodium tablets (listed in Section 6) have fluid retention have heart disease or failure have aldosteronism (a condition where too much aldosterone [a hormone] is produced) are having treatment to produce salt and water loss, usually if you have fluid retention or heart failure or have high blood pressure. 	<p>Warnings and precautions Talk to your doctor or pharmacist before taking Slow Sodium if you:</p> <ul style="list-style-type: none"> have a condition of the intestines, where narrowing may occur e.g. Crohn's disease, diverticulitis or stricture (band) formation (usually after major abdominal surgery). think you may have a problem drinking enough water. <p>Other medicines and Slow Sodium tablets Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. In particular, Slow Sodium may affect medicines (e.g. furosemide, bumetanide, metolazone) used to treat high blood pressure particularly if you also have a kidney disorder.</p>	<p>It may still be all right for you to be given Slow Sodium tablets and your doctor will be able to decide what is suitable for you.</p> <p>Pregnancy and breast-feeding If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.</p> <p>Driving and using machines Slow Sodium tablets should have no effect on your ability to drive and use machines.</p> <p>3. HOW TO TAKE SLOW SODIUM TABLETS Always take this medicine exactly as described in this leaflet or as your doctor has told you. Check with your doctor or pharmacist if you are not sure.</p>	<p>Dosage The recommended dose is:</p> <p>Adults: To prevent sodium chloride deficiency: 4-8 tablets per day. For cases of severe salt loss: Up to 20 tablets per day. To control muscle cramps during haemodialysis (treatment with a kidney machine): 10-16 tablets.</p> <p>Use in children and adolescents: The dose is adjusted to the child's needs.</p> <p>The tablets should be swallowed whole with water (about 70ml per tablet) and not chewed.</p> <p>If you take more Slow Sodium tablets than you should Tell your doctor straight away or go to the nearest casualty department immediately, taking the tablet container with you.</p>
---	--	---	---	--	---

Outer Bottle Label

<p>Slow Sodium Tablets® 100 tablets</p> <p>For the treatment and prevention of sodium chloride deficiency.</p> <p>Dose Adults: To prevent sodium chloride deficiency: 4-8 tablets per day. For cases of severe salt loss: Up to 20 tablets per day. To control muscle cramps during haemodialysis: 10-16 tablets. Children: As directed by a doctor.</p> <p>Each tablet should be swallowed whole with approx. 70ml of water and not chewed.</p>	<p>Read the package leaflet before use.</p> <p>Keep out of the sight and reach of children.</p> <p>Store in the original bottle in order to protect from moisture. Store below 30°C.</p> <p>Each tablet contains 600mg sodium chloride equivalent to approx. 10mEq of Na+ (230mg Na) in a sustained release formulation.</p>	 <p>PL 44490/0004 ALTURIX Ltd 287 Upper Fourth Street, Milton Keynes, MK9 1EH</p> 	 <p>1401689 a EXPLOT</p>
---	--	--	---

Outer Bottle Label - To display Braille

<p>Slow Sodium Tablets® 100 tablets</p> <p>For the treatment and prevention of sodium chloride deficiency. Read the package leaflet before use.</p> <p>Dose Adults: To prevent sodium chloride deficiency: 4-8 tablets per day. For cases of severe salt loss: Up to 20 tablets per day. To control muscle cramps during haemodialysis: 10-16 tablets. Children: As directed by a doctor.</p> <p>Each tablet should be swallowed whole with approx. 70ml of water and not chewed.</p>	<p>Keep out of the sight and reach of children.</p> <p>Store in the original bottle in order to protect from moisture. Store below 30°C.</p> <p>Each tablet contains 600mg sodium chloride equivalent to approx. 10mEq of Na+ (230mg Na) in a sustained release formulation.</p>	 <p>PL 44490/0004 ALTURIX Ltd 287 Upper Fourth Street, Milton Keynes, MK9 1EH</p> 	 <p>1401689 a EXPLOT</p>
--	--	--	--