



Fach 10ml of the oral suspension contains 1000mg of sodium alginate and 200mg of potassium hydrogen carbonate. Also contains parahydroxybenzoates (Parabens, F214 and F216) which may cause allergic reactions (possibly delayed). This medicinal product contains about 5.1mmol of sodium and 2mmol of notassium per 10ml dose. To be taken into consideration by patients on a controlled sodium diet or when taking Advance drugs which can increase potassium revers. Care should be also be taken if on calcium controlled diet. Read the attached leaflet carefully before use. To be taken by the controlled the patterns of the better well. druns which can increase notassium levels. Care should gension mouth after meals and at bedtime. Shake bottle well before use. Keep out of the sight and reach of children. Take special care if you are allergic to any of the potassium hydrogen ingredients of this medicine. Side effects are occasional, the potassium hydrogen ingredients of this medicine. Side effects are occasional, the potassium hydrogen in the potassium ee leaflet for further information. Store below 25°C. Do not use after the expiry date shown on the label. to not refrigerate or freeze.

Braille reads:

Braille Warning! Cirrus cannot accept responsibility for any errors in this proof after approval by the customer.

Whilst extreme care is taken in the setting of Braille, the customer must take the final responsibility for its accuracy.

There is no single European Braille authority and there are many different Braille formats in existence, with country

This Braille is set to the Marburg Medium format unless you have requested otherwise.

When you sign this proof you are signifying full approval of the Braille text and specification.

PACKAGE LEAFLET: INFORMATION FOR THE USER
Acidex Advance Oral Suspension
sodium alginate 1000mg/10ml, potassium hydrogen carbonate 200mg/10ml

Read all of this leaflet carefully before you start taking this medicine.

These are of an execute calculus belone you start canning units interesting us. Breath and the start canning units interesting units inter

- Reep this leaflet. You may need to read it again.
 If you have any further questions, ask your doctor or pharmacist.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet,

What Acidex Advance is and what it is used for

- Before you take Acidex Advance How to take Acidex Advance
- Possible side effects How to store Acidex Advance

6. Further information.
1. WHAT ACIDEX DVANICE IS AND WHAT IT IS USED FOR
Acidex Advance Oral Suspension (referred to as "Acidex Advance" throughout this leaflet) belongs to a
group of medicine called "reflux suppressants." Felfux is when the acid stomach contents flow back
up the guilet (pesophagus) and this can cause pain and discomfort known as hearthurn or acid
indigestion. This medicine contains the active ingredients sodium alignizate and potassium hydrogen
carbonate which together form a protective layer at the top of your stomach to stop reflux from

happening by keeping the stomach contents away from the lining of the food pipe to relieve the symptoms of hearburn and acid indigestion. It can therefore be taken to prevent and relieve the symptoms of acid reliux, including during pregnancy and relieve the symptoms of conditions such as hatus hernia (prottuson of muscle through a muscle wall), reflux oesophigis (inflamed foodpipe) and symptoms of hoarseness and other vioce disorders, sole throat and cough associated with reflux. If can also be taken to control hearthum symptoms which may occur when taking, or following withdrawal of medication to reduce stomach acid such as Proton Pump hinbliotic Plays or 122.

2. BEFORE YOU TAKE ACIDEX ADVANCE Do not take Acidex Advance if you:

That take Notes Advance in you.

are allergic (hypersensitive) to any of the ingredients in this medicine, including the esters of hydroxybenzoates (parabens) (see Section 6 and end of Section 2

Warnings and precautions:

Warnings and precautions:

I mmol per 10ml) potassium (2.0mmol per 10ml) and calcium.

If you have been advised to follow a diet restricted in any of these please consult you doctor before

taxing mis product.

The maximum recommended daily dose of this medicinal product contains 424mg sodium (found in table sail). This is equivalent to 21% of the adult recommended maximum daily infake for sodium. Please also talk to your doctor or pharmacist if you need this product on a daily basis for a prolonged period, especially if you have been advised to follow a low salt (sodium) diet or if you stifer or have suffered from significant kidney or heart disease, as certain salts could interfer with

Taking other medicines
Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines,

including medicines obtained without prescription. Wait at least 2 hours between taking this medicine and other medicines such as tetracyclines, fluoroquinolones (types of antibiotics), iron Salts, thyroid homones, chloroquine (a medication used to prevent malaria), bisphosphorates (a medication used to treat osteoporosis) and estramustine (a medication to treat prostate cancer).

Pregnancy and breast-feeding his medical to the product of the product of the product of the product of the pregnancy and breast-feeding, however it is recommended to limit the treatment duration as much as possible. Driving and using machines

here are no known effects of this medicine on ability to drive or operate

Important information about some of the ingredients of Acidex Advance

s medicine also contains:

- s rieducine add cotination.

 Parahydroxydenzoates: may cause allergic reactions, possibly delayed.

 Sodium: each 10ml dose of this medicine contains about 5. Intimol of sodium and therefore care should be exercised in patients on a sodium restricted diet.

 Potassium: Each 10ml dose of this medicine contains about 2mmol of potassium and therefore.
- care should be taken when taking drugs which can increase potassium levels.

HOW TO TAKE ACIDEX ADVANCE

To be taken by mouth, after meals and at bedtime. Shake bottle well before use. If you are unsure how to use this medicine, check with your doctor or pharmacist.

Adults including the elderly and children 12 years and over:	5 – 10ml (one to two 5ml spoonfuls) after meals and at bedtime.
Children under 12 years:	Should only be taken on medical advice.

WARNING: DO NOT EXCEED THE STATED DOSE

If you take more of this medicine than you should

If you take those of this product you may feel bloated and experience some abdominal discomfort. It is unlikely to cause you any harm, but please consult your doctor or pharmcist if this doesn't go away. If symptoms persist after 7 days consult your doctor.

If you forget to take this medicine
If you forget to take this medicine
If you forget to take your medicine, take it as soon as you remember, **Do not** take a double dose to make up for the forgotten dose.

4. POSSIBLE SIDE EFFECTS

PLASSINE SIDE EFFCIS
Like all medicines, this product can cause side effects, although not everybody gets them. Very rarely (ses than 1 in 10,000 patients) there is a chance of an altergic reaction to the ingredients. Side effects are districted, nauses and vomiting, if you experience these or any other side effects stop cannot be an advised of the product and consult your doctor immediately.
Reporting of side effects.

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side Infoot any activities that the state of the

HOW TO STORE ACIDEX ADVANCE Keep out of the sight and reach of children.

Do not store above 25°C. Do not refrigerate or freeze. Do not use this product after the expiry date

which is stated on the pack. The expiry date refers to the last day of that month.

Which is stated on the pack. The expiry date refers to the last day of that month wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will

FURTHER INFORMATION What Acidex Advance contains:

- he active ingredients are: Sodium alginate 1000mg/10ml, Potassium hydrogen carbonate
- 200mg/10ml
 The offer ingredients are: Calcium Carbonate, Carbomer, Sodium Hydroxide, Saccharin Sodium,
 Ethyl Parahydroxybenzoate (E214), Propyl Parahydroxybenzoate (E216), Butyl Parahydroxybenzoate,
 Isopropyl Alcohol, Star Anise Oil and Purified Water (see end of Section 2 for further information)

What Acidex Advance looks like and contents of the pack

This medicine is an aniseed flavoured white or cream coloured suspension.


It comes in amber glass bottles of 150ml, 200ml, 250ml, 300ml and 500ml with white polypropylene caps. Not all pack sizes may be marketed.

Marketing Authorisation Holder Wockhardt UK Ltd, Ash Road North, Wrexham, LL13 9UF, UK.

Manufacturer

Pinewood Laboratories Limited, Ballymacarbry, Clonmel, Co. Tipperary, Ireland. PI 29831/0697

This leaflet was last updated in 10/2022.



Each 10ml of the oral suspension contains 1000mg of sodium alginate and 200mg of potassium hydrogen carbonate. Also contains parahydroxybenzoates (Parabens, E214 and E216) which may cause allergic reactions (pbs.bb) delayed). This netricinar product contains about 5, immol of sodium and 2mmol of potassium per 10ml dose. To be taken into consideration by patients on Advance and drugs which can increase per assistant evels. See should the controlled diet. Read the attached leaflet carefully before use. To be taken by mouth after meals and at bedtime. Shake bottle well before use, Keep out of the sight and reach of children. sedium alginate and potassium hydrogen carbonate and potassium hydrogen carbonate sedium carbonate sedium alginate and potassium hydrogen carbonate sedium s See leaflet for further information. Store below 25°C. Do not use after the expiry date shown on the label. Do not refrigerate or freeze.