
### How to use the Easyhaler inhaler

### About your Easyhaler

Easyhaler Beclometasone may be different to inhalers you have used in the past. Therefore it is very important that you use it properly as using incorrectly can lead to you not receiving the right amount of medicine. This could make you very unwell or could lead to your asthma not being treated as it should.

Your doctor, nurse or pharmacist will show you how to use your inhaler properly. Make sure you understand the correct way to use the inhaler. If you are unsure contact your doctor, nurse or pharmacist. As with all inhalers, caregivers should ensure that children prescribed Easyhaler Beclometasone use correct inhalation technique, as described below. You can also use the instruction video at <u>www.oeh.fi/begb.</u>

### When you first get your Easyhaler



# HOW TO USE CORRECTLY

Step 1: SHAKE	SHAKE x 3-5	Important points to remember
• Remove the dustcap • Shake the inhaler <b>3</b> to <b>5</b> times holding it in the <b>upright</b> <b>position</b>		<ul> <li>It is important to keep the inhaler in the upright position</li> <li>If you accidentally click while you shake the inhaler, empty the powder from the mouthpiece as shown below</li> </ul>

Step 2: CLICK	CLICK x 1	Important points to remember
<ul> <li>Keep holding the inhaler upright between your forefinger and thumb</li> <li>Press down until you hear a click, and let the inhaler click back again. This releases a dose</li> <li>Only click down once</li> </ul>		<ul> <li>The inhaler will not click if the dustcap is still on</li> <li>Only click down once</li> <li>If you accidentally click more than once, empty the powder from the mouthpiece, see below</li> <li>Click to release dose before you inhale, not at the same time</li> <li>Keep the inhaler upright when you click it and when you lick it and when you inhale the dose. If you tip it, the powder could fall out before you are able to inhale it</li> </ul>

Step 3: INHALE	INHALE	Important points to remember
<ul> <li>Keep holding the inhaler upright</li> <li>Breathe out normally</li> <li>Place the mouthpiece in your mouth between your teeth and close your lips tightly around the mouthpiece</li> <li>Take a strong and deep breath in</li> <li>Take the inhaler out of your mouth, hold breath for at least</li> </ul>		<ul> <li>Make sure the whole mouthpiece is well inside your mouth, so that the medication gets into your lungs</li> <li>Make sure your lips make a good seal around the mouthpiece</li> <li>Do not breathe out into the inhaler. This is important: it could clog up the inhaler. If you breathed out into the inhaler, empty the powder from</li> </ul>


# If you need to take another inhalation, please repeat the steps 1-3 Shake-Click-Inhale.

### After you have used the inhaler:

normally.


• Put the dust cap back on the mouthpiece. It stops the inhaler going off by accident.

• After you have taken the dose, rinse your mouth with water, and spit it out and/or brush your teeth. This may help to stop you getting thrush and becoming hoarse.



# **Cleaning the Easyhaler**

Keep your inhaler dry and clean. If necessary you may wipe the mouthpiece of your inhaler with a dry cloth or tissue. Do not use water: the powder in the Easyhaler is sensitive to moisture.



# When to switch to a new Easyhaler

The dose counter shows the number of remaining doses. The counter turns after every 5th click. When the dose counter starts turning red, there are 20 doses left.

If you do not already have a new Easyhaler, contact your doctor for a new prescription. When the counter reaches 0 (zero), you need to replace the Easyhaler.

If you use the protective cover, you can keep it and insert your new inhaler into it.



# Remember

- 1. Shake 2. Click 3. Inhale.
- After you have taken the dose, rinse your mouth with water and spit it out and/or brush your teeth.
- Do not get your inhaler wet, protect it from moisture.