

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet.

You can also report side effects directly via the Yellow Card Scheme at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard), or search for MHRA Yellow Card in the Google Play or Apple App Store.

You should also report side effects to Roche Products Ltd by emailing the Roche Drug Safety Centre at [welwyn.uk\\_dsc@roche.com](mailto:welwyn.uk_dsc@roche.com) or calling +44 (0) 1707 367554.

By reporting side effects you can help provide more information on the safety of this medicine.



### Emergency Contact Phone Numbers

Office Hours

Out of Hours

## Erivedge<sup>®</sup> (vismodegib) *Patient reminder card*

This educational material is provided by Roche Products Limited and is mandatory as a condition of the Marketing Authorisation in order to further minimise important selected risks.

RXUKERIV00203a(1)

Date of preparation: January 2019

**Erivedge may harm or cause the death of an unborn child.**

**FEMALE PATIENTS** who are able to become pregnant must:

- always use **2 forms** of recommended contraception during treatment and for 24 months after your last dose (please refer to the “Erivedge® Pregnancy Prevention Programme: Important information for men and women taking Erivedge about pregnancy prevention and contraception” for more information).
- have monthly pregnancy tests during treatment to monitor for pregnancy.

**MALE PATIENTS** must:

- always use a condom (with spermicide, if available) when having sex with female partners during treatment and for 2 months from final dose.
- not donate semen during treatment and for 2 months after the final dose.

You must talk to your healthcare professional immediately if you or your sexual partner misses a period, suspect a pregnancy, or are pregnant.

- **FEMALE PATIENTS:** talk to your doctor if you suspect a pregnancy while taking Erivedge and for 24 months after your final dose.
- **MALE PATIENTS:** talk to your doctor if your female partner suspects that she is pregnant while you are taking Erivedge and for 2 months after your final dose.

**ALL PATIENTS** must:

- never give this medicine to anyone else.
- not donate blood during treatment and for 24 months after the final dose.
- return all unused capsules at the end of treatment.