

What should I know about Rivaroxaban Gedeon Richter?

- Rivaroxaban Gedeon Richter thins the blood, which prevents you from getting dangerous blood clots.
- Rivaroxaban Gedeon Richter must be taken exactly as prescribed by your doctor. To ensure optimal protection from blood clots, **never skip a dose.**
- You must not stop taking Rivaroxaban Gedeon Richter without first talking to your doctor as your risk of blood clots may increase.

- Tell your health care provider about any other medicines you are currently taking, took recently or intend to start taking, before you start Rivaroxaban Gedeon Richter.
- Tell your health care provider that you are taking Rivaroxaban Gedeon Richter before any surgery or invasive procedure.

When should I seek advice from my health care provider?

When taking a blood thinner such as Rivaroxaban Gedeon Richter it is important to be aware of its possible side effects. Bleeding is

the most common side effect. Do not start taking Rivaroxaban Gedeon Richter if you know you are at risk of bleeding, without first discussing this with your doctor. Tell your health care provider straight away if you have any signs or symptoms of bleeding such as the following:

- pain
- swelling or discomfort
- headache, dizziness or weakness
- unusual bruising, nosebleeds, bleeding of gums, cuts that take a long time to stop bleeding
- menstrual flow or vaginal bleeding that is heavier than

normal

- blood in your urine which may be pink or brown, red or black stools
- coughing up blood, or vomiting blood or material that looks like coffee grounds

Patient Alert Card

- RIVAROXABAN
GEDEON RICHTER 2.5 mg**
- RIVAROXABAN
GEDEON RICHTER 10 mg**
- RIVAROXABAN
GEDEON RICHTER 15 mg**
- RIVAROXABAN
GEDEON RICHTER 20 mg**

- Keep this card with you at all times
- Present this card to every physician or dentist prior to treatment

How do I take Rivaroxaban Gedeon Richter?

- To ensure optimal protection, Rivaroxaban Gedeon Richter
 - 2.5 mg can be taken with or without food
 - 10 mg can be taken with or without food
 - 15 mg must be taken with food
 - 20 mg must be taken with food

I am under anticoagulation treatment with Rivaroxaban Gedeon Richter (rivaroxaban)

Name:
Address:
Birth date:
Weight:
Other medicines / conditions:

In case of emergency, please notify:

Doctor's name:
Doctor's phone:
Doctor's stamp:

Please also notify:


Name:
Phone:
Relationship:

Information for health care providers:

- INR values should not be used as they are not a dependable measure of the anticoagulant activity of Rivaroxaban Gedeon Richter.

K-36132-1.0

 GEDEON RICHTER

nazwa/name	RIVAROXABAN GEDEON RICHTER 2,5 mg, 10 mg, 15 mg, 20 mg	kraj/country	UK
		rodzaj/type	pcard
numer/number	K-36132-1.0		
wymiary/size	218x68mm		
kolory/colours	 P 280		
czcionka/font size	Futura AOT (8 pt) [9.2 pt]	uwagi/comment ___ ___ ___ pac folding lines (unprinted)	
data/date	2024.05.17		
grafik/designer			