

60 mm



DXXXXXXXX-XX

Apixaban 2.5 mg and 5 mg film-coated tablets (MSND)

**Patient Alert Card**

**This card should be with you / the child / the caregiver at all times.**

**Show this card to your pharmacist, dentist and any other healthcare professionals before treatment.**

**I am under anticoagulation treatment with Apixaban to prevent blood clots.**

**Please complete this section or ask your doctor to do it**

Name:

Birth Date:

Indication:

Weight:

Dose: ..... mg twice daily

Doctor's Name:

Doctor's telephone:

**Information for patients / caregivers**

- Take / give Apixaban regularly as instructed. If you miss a morning dose, take / give it as soon as you remember and it may be taken / given together with the evening dose. A missed evening dose can only be taken / given during the same evening. Do not take / give two doses the next morning, instead continue to follow the dosing schedule twice daily as recommended on the next day.
- Do not stop taking / giving Apixaban without talking to the doctor, as you are / the patient is at risk of suffering from a stroke / blood clot or other complications.
- Apixaban helps to thin the blood. However, this may increase your risk of bleeding.
- Signs and symptoms of bleeding include bruising or bleeding under

168 mm

28 mm

Fold here

28 mm

28 mm

28 mm

28 mm

28 mm

