



My Enbrel Patient Alert Card

Please carry this card with you at all times
PP-ENB-GBR-0589. Date of preparation: August 2017



Congestive Heart Failure

If you develop symptoms suggestive of congestive heart failure or worsening of existing congestive heart failure, such as shortness of breath, swelling of ankles, persistent cough or fatigue, seek medical attention immediately.

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

Other information

Patient's name:

Doctor's name:

Doctor's phone:

My Enbrel (etanercept) Patient Alert Card

Please carry this card with you at all times

This card contains important safety information that you need to be aware of before you are given Enbrel, and during treatment with Enbrel. If you do not understand this information, please ask your doctor to explain it to you.

Show this card to any doctor involved in your treatment.

Keep this card with you for 2 months after the last Enbrel dose, since side effects may occur after your last dose of Enbrel.

See the Enbrel package leaflet for more information.

It is **important** that you and your doctor record the brand name and batch number of your medication.

Brand name:

Batch number:

Infections. Enbrel may increase your risk of getting infections, which could be serious.

You should not use Enbrel if you have an infection. If you are not sure, ask your doctor.

If you **develop symptoms** suggestive of infections, such as fever, persistent cough, weight loss, or listlessness, seek medical attention immediately.

You should be evaluated for tuberculosis (TB). Ask your doctor to record the dates and the results of your last screening for TB below.

Test:

Date:

Results:

Test:

Date:

Results:

Please ask your physician to list your other medications that may increase your risk of infection

.....