

Eydenzelt®▼

(aflibercept biosimilar)

Patient Guide

Table of Contents

Your guide to EYDENZELT for wAMD	2
Your guide to EYDENZELT for DME	10
Your guide to EYDENZELT for CRVO	18
Your guide to EYDENZELT for BRVO	26
Your guide to EYDENZELT for mCNV	34

Your guide to EYDENZELT®▼

EYDENZELT is used to treat Wet Age-Related Macular Degeneration (Wet AMD)

The guide has been produced for people who have been prescribed EYDENZELT (aflibercept solution for injection) for the treatment of wAMD. Your doctor will use the EYDENZELT 40 mg/ml solution for the recommended dose of 2mg aflibercept.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 of the SmPC for how to report adverse reactions.

The guide is available in printed and digital format, both versions allow you to access an audio version via a smart device. The audio version is a readout of the text in this patient guide. More information is available to you in the Patient Information Leaflet (PIL).

This information can be accessed by visiting <https://www.medicines.org.uk/emc> or by using the camera on a smart device to scan the relevant QR codes. This will direct the device to relevant information on the internet.

Patient Audio wAMD

QR code to Patient Audio wAMD



Patient Information Leaflet

Eydenzelt 40 mg/ml (2 mg dose) QR code to Patient Information Leaflet



Your doctor has prescribed EYDENZELT because you have been diagnosed with wet age-related macular degeneration, or Wet AMD.

What is EYDENZELT?

EYDENZELT is a type of treatment known as an anti-VEGF. Anti-VEGF is an abbreviation for anti-vascular endothelial growth factor, and this is a description of how EYDENZELT works to protect your vision. EYDENZELT blocks VEGF, and this action helps reduce the swelling in the retina and can lead to vision improvement and maintenance.

EYDENZELT is a solution (a liquid) that is injected into the eye. It is available in a 2 mg dose. Your doctor will recommend a treatment schedule for you and it is very important that you follow it.

EYDENZELT is not indicated for paediatric use unlike the original product Eylea.

What should your doctor know before you are treated with EYDENZELT?

Before your EYDENZELT treatment starts, make sure to tell your doctor and other health care providers if you:

- Have an infection in or around your eye.
- If you currently have redness in your eye or if there is any pain in your eye.
- Think you may be allergic to iodine, any pain killers or any of the ingredients in EYDENZELT.
- Have had any issues or problems with eye injections before.
- Have glaucoma or a history of high pressure in your eye.
- If you see, or have seen, flashes of light or 'floaters' in your vision.
- Are taking any medications, with or without a prescription.
- Had or will have eye surgery within 4 weeks before or after EYDENZELT treatment.
- Are pregnant, planning to become pregnant or breastfeeding. There is very little information about the safety of using EYDENZELT in pregnant women. EYDENZELT 2 mg should not be used during pregnancy, unless the benefit outweighs the risk to the foetus. Discuss this with your doctor before treatment with EYDENZELT. Women of childbearing potential should use effective contraception during their treatment with EYDENZELT. If you are treated with the EYDENZELT 2 mg dose, you should continue to use effective contraception for at least three months after the last injection. Small amounts of EYDENZELT may enter into the breastmilk. The effects of aflibercept on a breast-fed newborn/ infant is unknown. Use of EYDENZELT is not recommended during breastfeeding.

How can I get ready for my EYDENZELT appointment?

Your doctor may ask you to use eye drops for a few days before your appointment. After your treatment, your vision may be blurry so you should not drive home. Plan to have a friend or family member take you to your appointment or arrange another way to get there and home again. On the day of your appointment, do not wear any makeup.

What can I expect after the EYDENZELT treatment?

Your doctor may give you some eye tests after your EYDENZELT injection. This may include a test that measures the pressure inside your eye. After your injection, your vision will be blurry so you should not plan to drive until after your vision returns to normal.

During the next few days you may get a bloodshot eye or see moving spots in your vision. These should clear within a few days and if they don't, or if they get worse, you should contact your doctor.

Some people might feel a little bit of pain or discomfort in their eye after their injection. If this does not go away or gets worse, you should contact your doctor.

Does my treatment have side effects?

Just like any medicine, EYDENZELT has the potential to cause side effects. Not everyone who is given an EYDENZELT injection will experience a side effect.

Contact your doctor immediately if you have any signs or symptoms listed in the table below as these could be signs of a serious complication with the treatment:

Condition	Some Potential Signs or Symptoms
Infection or inflammation inside the eye	• Eye pain or increased discomfort • Worsening eye redness • Sensitivity to light • Swelling of the eyelid • Vision changes such as sudden decrease in vision or blurring of vision
Clouding of the lens (cataract)	• Blurry vision • Seeing shadows • Less vivid lines and shapes • Colour vision changes (e.g. colours look 'washed out')
Increase in pressure in the eye	• Seeing halos around lights • Eye pain • Experiencing a red eye • Nausea or vomiting • Vision changes
A detachment or tear of a layer of the retina	• Sudden flashes of light • A sudden appearance or an increase of floaters • A curtain like effect over a portion of the visual area • Vision changes

For a full list of side effects, please refer to the relevant EYDENZELT Patient Information Leaflet (PIL). You can access the PIL using the QR code at the beginning of this guide, or by asking your doctor.

If you get any side effects, which include any possible side effects not listed in this guide, talk to your doctor, pharmacist or nurse. You can also report side effects directly to the MHRA through the Yellow Card scheme, via the Yellow Card website www.mhra.gov.uk/yellowcard, the free Yellow Card app available from the Apple App Store or Google Play Store. Alternatively, you can call 0800 731 6789 for free, Monday to Friday between 9am and 5pm.

Adverse events can also be reported to Celltrion Healthcare UK by phone on +44 (0)1753 983500 or email at ukadverseevents@celltrionhc.com. By reporting side effects, you can help provide more information on the safety of this medicine.

What if I have concerns or questions?

If you have any concerns or questions, your doctor or nurse are the best people to speak to. They are very experienced and they know your individual situation so can provide you with the answers you need.

Useful contacts

The Macular Society

Helpline: 0300 3030 111

Address: The Macular Society, PO Box 1870, Andover SP10 9AD

Email: info@macularsociety.org

Website: www.macularsociety.org

RNIB (Royal National Institute of Blind People)

Helpline: 0303 123 9999

Address: Grimaldi Building, 154a Pentonville Road, London N1 9JE

Email: helpline@rnib.org.uk

Website: www.rnib.org.uk

Diabetes UK

Helpline: 0345 123 2399

Address: Wells Lawrence House, 126 Back Church Lane, London E1 1FH

Email: helpline@diabetes.org.uk

Website: www.diabetes.org.uk

Your eye care clinic is:

Contact:

Telephone:

Address:

Email:

Your guide to EYDENZELT®▼

EYDENZELT is used to treat Diabetic Macular Oedema (DME).

The guide has been produced for people who have been prescribed EYDENZELT (aflibercept solution for injection) for the treatment of DME. Your doctor will use the EYDENZELT 40 mg/ml solution for the recommended dose of 2mg aflibercept.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 of the SmPC for how to report adverse reactions.

The guide is available in printed and digital format, both versions allow you to access an audio version via a smart device. The audio version is a readout of the text in this patient guide. More information is available to you in the Patient Information Leaflet (PIL).

This information can be accessed by visiting <https://www.medicines.org.uk/emc> or by using the camera on a smart device to scan the relevant QR codes. This will direct the device to relevant information on the internet.

Patient Audio DME

QR code to Patient Audio DME



Patient Information Leaflet

Eydenzelt 40 mg/ml (2 mg dose)
QR code to Patient Information Leaflet



Your doctor has prescribed EYDENZELT because you have been diagnosed with diabetic macular oedema, or DME for short.

What is EYDENZELT?

EYDENZELT is a type of treatment known as an anti-VEGF. Anti-VEGF is an abbreviation for anti-vascular endothelial growth factor, and this is a description of how EYDENZELT works to protect your vision. EYDENZELT blocks VEGF, and this action helps reduce the swelling in the retina and can lead to vision improvement and maintenance.

EYDENZELT is a solution (a liquid) that is injected into the eye. It is available in a 2 mg dose. Your doctor will recommend a treatment schedule for you and it is very important that you follow it.

EYDENZELT is not indicated for paediatric use unlike the original product Eylea.

What should your doctor know before you are treated with EYDENZELT?

Before your EYDENZELT treatment starts, make sure to tell your doctor and other health care providers if you:

- Have an infection in or around your eye.
- If you currently have redness in your eye or if there is any pain in your eye.
- Think you may be allergic to iodine, any pain killers or any of the ingredients in EYDENZELT.
- Have had any issues or problems with eye injections before.
- Have glaucoma or a history of high pressure in your eye.
- If you see, or have seen, flashes of light or 'floaters' in your vision.
- Are taking any medications, with or without a prescription.
- Had or will have eye surgery within 4 weeks before or after EYDENZELT treatment.
- Are pregnant, planning to become pregnant or breastfeeding. There is very little information about the safety of using EYDENZELT in pregnant women. EYDENZELT 2 mg should not be used during pregnancy, unless the benefit outweighs the risk to the foetus. Discuss this with your doctor before treatment with EYDENZELT. Women of childbearing potential should use effective contraception during their treatment with EYDENZELT. If you are treated with the EYDENZELT 2 mg dose, you should continue to use effective contraception for at least three months after the last injection. Small amounts of EYDENZELT may enter into the breastmilk. The effects of aflibercept on a breast-fed newborn/ infant is unknown. Use of EYDENZELT is not recommended during breastfeeding.

How can I get ready for my EYDENZELT appointment?

Your doctor may ask you to use eye drops for a few days before your appointment. After your treatment, your vision may be blurry so you should not drive home. Plan to have a friend or family member take you to your appointment or arrange another way to get there and home again. On the day of your appointment, do not wear any makeup.

What can I expect after the EYDENZELT treatment?

Your doctor may give you some eye tests after your EYDENZELT injection. This may include a test that measures the pressure inside your eye. After your injection, your vision will be blurry so you should not plan to drive until after your vision returns to normal.

During the next few days you may get a bloodshot eye or see moving spots in your vision. These should clear within a few days and if they don't, or if they get worse, you should contact your doctor.

Some people might feel a little bit of pain or discomfort in their eye after their injection. If this does not go away or gets worse, you should contact your doctor.

Does my treatment have side effects?

Just like any medicine, EYDENZELT has the potential to cause side effects. Not everyone who is given an EYDENZELT injection will experience a side effect.

Contact your doctor immediately if you have any signs or symptoms listed in the table below as these could be signs of a serious complication with the treatment:

Condition	Some Potential Signs or Symptoms
Infection or inflammation inside the eye	• Eye pain or increased discomfort • Worsening eye redness • Sensitivity to light • Swelling of the eyelid • Vision changes such as sudden decrease in vision or blurring of vision
Clouding of the lens (cataract)	• Blurry vision • Seeing shadows • Less vivid lines and shapes • Colour vision changes (e.g. colours look 'washed out')
Increase in pressure in the eye	• Seeing halos around lights • Eye pain • Experiencing a red eye • Nausea or vomiting • Vision changes
A detachment or tear of a layer of the retina	• Sudden flashes of light • A sudden appearance or an increase of floaters • A curtain like effect over a portion of the visual area • Vision changes

For a full list of side effects, please refer to the relevant EYDENZELT Patient Information Leaflet (PIL). You can access the PIL using the QR code at the beginning of this guide, or by asking your doctor.

If you get any side effects, which include any possible side effects not listed in this guide, talk to your doctor, pharmacist or nurse. You can also report side effects directly to the MHRA through the Yellow Card scheme, via the Yellow Card website www.mhra.gov.uk/yellowcard, the free Yellow Card app available from the Apple App Store or Google Play Store. Alternatively, you can call 0800 731 6789 for free, Monday to Friday between 9am and 5pm.

Adverse events can also be reported to Celltrion Healthcare UK by phone on +44 (0)1753 983500 or email at ukadverseevents@celltrionhc.com. By reporting side effects, you can help provide more information on the safety of this medicine.

What if I have concerns or questions?

If you have any concerns or questions, your doctor or nurse are the best people to speak to. They are very experienced and they know your individual situation so can provide you with the answers you need.

Useful contacts

The Macular Society

Helpline: 0300 3030 111

Address: The Macular Society, PO Box 1870, Andover SP10 9AD

Email: info@macularsociety.org

Website: www.macularsociety.org

RNIB (Royal National Institute of Blind People)

Helpline: 0303 123 9999

Address: Grimaldi Building, 154a Pentonville Road, London N1 9JE

Email: helpline@rnib.org.uk

Website: www.rnib.org.uk

Diabetes UK

Helpline: 0345 123 2399

Address: Wells Lawrence House, 126 Back Church Lane, London E1 1FH

Email: helpline@diabetes.org.uk

Website: www.diabetes.org.uk

Your eye care clinic is:

Contact:

Telephone:

Address:

Email:

Your guide to EYDENZELT®▼

EYDENZELT is used to treat macular oedema secondary to Central Retinal Vein Occlusion (CRVO).

The guide has been produced for people who have been prescribed EYDENZELT (aflibercept solution for injection) for the treatment of CRVO. Your doctor will use the EYDENZELT 40 mg/ml solution for the recommended dose of 2mg aflibercept.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 of the SmPC for how to report adverse reactions.

The guide is available in printed and digital format, both versions allow you to access an audio version via a smart device. The audio version is a readout of the text in this patient guide. More information is available to you in the Patient Information Leaflet (PIL).

This information can be accessed by visiting <https://www.medicines.org.uk/emc> or by using the camera on a smart device to scan the relevant QR codes. This will direct the device to relevant information on the internet.

Patient Audio CRVO

QR code to Patient Audio CRVO



Patient Information Leaflet

Eydenzelt 40 mg/ml (2 mg dose)
QR code to Patient Information Leaflet



Your doctor has prescribed EYDENZELT because you have been diagnosed with macular oedema secondary to central retinal vein occlusion, or CRVO.

What is EYDENZELT?

EYDENZELT is a type of treatment known as an anti-VEGF. Anti-VEGF is an abbreviation for anti-vascular endothelial growth factor, and this is a description of how EYDENZELT works to protect your vision. EYDENZELT blocks VEGF, and this action helps reduce the swelling in the retina and can lead to vision improvement and maintenance.

EYDENZELT is a solution (a liquid) that is injected into the eye. It is available in a 2 mg dose. Your doctor will recommend a treatment schedule for you and it is very important that you follow it.

EYDENZELT is not indicated for paediatric use unlike the original product Eylea.

What should your doctor know before you are treated with EYDENZELT?

Before your EYDENZELT treatment starts, make sure to tell your doctor and other health care providers if you:

- Have an infection in or around your eye.
- If you currently have redness in your eye or if there is any pain in your eye.
- Think you may be allergic to iodine, any pain killers or any of the ingredients in EYDENZELT.
- Have had any issues or problems with eye injections before.
- Have glaucoma or a history of high pressure in your eye.
- If you see, or have seen, flashes of light or 'floaters' in your vision.
- Are taking any medications, with or without a prescription.
- Had or will have eye surgery within 4 weeks before or after EYDENZELT treatment.
- Are pregnant, planning to become pregnant or breastfeeding. There is very little information about the safety of using EYDENZELT in pregnant women. EYDENZELT 2 mg should not be used during pregnancy, unless the benefit outweighs the risk to the foetus. Discuss this with your doctor before treatment with EYDENZELT. Women of childbearing potential should use effective contraception during their treatment with EYDENZELT. If you are treated with the EYDENZELT 2 mg dose, you should continue to use effective contraception for at least three months after the last injection. Small amounts of EYDENZELT may enter into the breastmilk. The effects of aflibercept on a breast-fed newborn/ infant is unknown. Use of EYDENZELT is not recommended during breastfeeding.

How can I get ready for my EYDENZELT appointment?

Your doctor may ask you to use eye drops for a few days before your appointment. After your treatment, your vision may be blurry so you should not drive home. Plan to have a friend or family member take you to your appointment or arrange another way to get there and home again. On the day of your appointment, do not wear any makeup.

What can I expect after the EYDENZELT treatment?

Your doctor may give you some eye tests after your EYDENZELT injection. This may include a test that measures the pressure inside your eye. After your injection, your vision will be blurry so you should not plan to drive until after your vision returns to normal.

During the next few days you may get a bloodshot eye or see moving spots in your vision. These should clear within a few days and if they don't, or if they get worse, you should contact your doctor.

Some people might feel a little bit of pain or discomfort in their eye after their injection. If this does not go away or gets worse, you should contact your doctor.

Does my treatment have side effects?

Just like any medicine, EYDENZELT has the potential to cause side effects. Not everyone who is given an EYDENZELT injection will experience a side effect.

Contact your doctor immediately if you have any signs or symptoms listed in the table below as these could be signs of a serious complication with the treatment:

Condition	Some Potential Signs or Symptoms
Infection or inflammation inside the eye	• Eye pain or increased discomfort • Worsening eye redness • Sensitivity to light • Swelling of the eyelid • Vision changes such as sudden decrease in vision or blurring of vision
Clouding of the lens (cataract)	• Blurry vision • Seeing shadows • Less vivid lines and shapes • Colour vision changes (e.g. colours look 'washed out')
Increase in pressure in the eye	• Seeing halos around lights • Eye pain • Experiencing a red eye • Nausea or vomiting • Vision changes
A detachment or tear of a layer of the retina	• Sudden flashes of light • A sudden appearance or an increase of floaters • A curtain like effect over a portion of the visual area • Vision changes

For a full list of side effects, please refer to the relevant EYDENZELT Patient Information Leaflet (PIL). You can access the PIL using the QR code at the beginning of this guide, or by asking your doctor.

If you get any side effects, which include any possible side effects not listed in this guide, talk to your doctor, pharmacist or nurse. You can also report side effects directly to the MHRA through the Yellow Card scheme, via the Yellow Card website www.mhra.gov.uk/yellowcard, the free Yellow Card app available from the Apple App Store or Google Play Store. Alternatively, you can call 0800 731 6789 for free, Monday to Friday between 9am and 5pm.

Adverse events can also be reported to Celltrion Healthcare UK by phone on +44 (0)1753 983500 or email at ukadverseevents@celltrionhc.com. By reporting side effects, you can help provide more information on the safety of this medicine.

What if I have concerns or questions?

If you have any concerns or questions, your doctor or nurse are the best people to speak to. They are very experienced and they know your individual situation so can provide you with the answers you need.

Useful contacts

The Macular Society

Helpline: 0300 3030 111

Address: The Macular Society, PO Box 1870, Andover SP10 9AD

Email: info@macularsociety.org

Website: www.macularsociety.org

RNIB (Royal National Institute of Blind People)

Helpline: 0303 123 9999

Address: Grimaldi Building, 154a Pentonville Road, London N1 9JE

Email: helpline@rnib.org.uk

Website: www.rnib.org.uk

Diabetes UK

Helpline: 0345 123 2399

Address: Wells Lawrence House, 126 Back Church Lane, London E1 1FH

Email: helpline@diabetes.org.uk

Website: www.diabetes.org.uk

Your eye care clinic is:

Contact:

Telephone:

Address:

Email:

Your guide to EYDENZELT®▼

EYDENZELT is used to treat macular oedema secondary to Branch Retinal Vein Occlusion (BRVO).

The guide has been produced for people who have been prescribed EYDENZELT (aflibercept solution for injection) for the treatment of BRVO. Your doctor will use the EYDENZELT 40 mg/ml solution for the recommended dose of 2mg aflibercept.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 of the SmPC for how to report adverse reactions.

The guide is available in printed and digital format, both versions allow you to access an audio version via a smart device. The audio version is a readout of the text in this patient guide. More information is available to you in the Patient Information Leaflet (PIL).

This information can be accessed by visiting <https://www.medicines.org.uk/emc> or by using the camera on a smart device to scan the relevant QR codes. This will direct the device to relevant information on the internet.

Patient Audio BRVO

QR code to Patient Audio BRVO



Patient Information Leaflet

Eydenzelt 40 mg/ml (2 mg dose)
QR code to Patient Information Leaflet



Your doctor has prescribed EYDENZELT because you have been diagnosed with macular oedema secondary to branch retinal vein occlusion, or BRVO.

What is EYDENZELT?

EYDENZELT is a type of treatment known as an anti-VEGF. Anti-VEGF is an abbreviation for anti-vascular endothelial growth factor, and this is a description of how EYDENZELT works to protect your vision. EYDENZELT blocks VEGF, and this action helps reduce the swelling in the retina and can lead to vision improvement and maintenance.

EYDENZELT is a solution (a liquid) that is injected into the eye. It is available in a 2 mg dose. Your doctor will recommend a treatment schedule for you and it is very important that you follow it.

EYDENZELT is not indicated for paediatric use unlike the original product Eylea.

What should your doctor know before you are treated with EYDENZELT?

Before your EYDENZELT treatment starts, make sure to tell your doctor and other health care providers if you:

- Have an infection in or around your eye.
- If you currently have redness in your eye or if there is any pain in your eye.
- Think you may be allergic to iodine, any pain killers or any of the ingredients in EYDENZELT.
- Have had any issues or problems with eye injections before.
- Have glaucoma or a history of high pressure in your eye.
- If you see, or have seen, flashes of light or 'floaters' in your vision.
- Are taking any medications, with or without a prescription.
- Had or will have eye surgery within 4 weeks before or after EYDENZELT treatment.
- Are pregnant, planning to become pregnant or breastfeeding. There is very little information about the safety of using EYDENZELT in pregnant women. EYDENZELT 2 mg should not be used during pregnancy, unless the benefit outweighs the risk to the foetus. Discuss this with your doctor before treatment with EYDENZELT. Women of childbearing potential should use effective contraception during their treatment with EYDENZELT. If you are treated with the EYDENZELT 2 mg dose, you should continue to use effective contraception for at least three months after the last injection. Small amounts of EYDENZELT may enter into the breastmilk. The effects of aflibercept on a breast-fed newborn/ infant is unknown. Use of EYDENZELT is not recommended during breastfeeding.

How can I get ready for my EYDENZELT appointment?

Your doctor may ask you to use eye drops for a few days before your appointment. After your treatment, your vision may be blurry so you should not drive home. Plan to have a friend or family member take you to your appointment or arrange another way to get there and home again. On the day of your appointment, do not wear any makeup.

What can I expect after the EYDENZELT treatment?

Your doctor may give you some eye tests after your EYDENZELT injection. This may include a test that measures the pressure inside your eye. After your injection, your vision will be blurry so you should not plan to drive until after your vision returns to normal.

During the next few days you may get a bloodshot eye or see moving spots in your vision. These should clear within a few days and if they don't, or if they get worse, you should contact your doctor.

Some people might feel a little bit of pain or discomfort in their eye after their injection. If this does not go away or gets worse, you should contact your doctor.

Does my treatment have side effects?

Just like any medicine, EYDENZELT has the potential to cause side effects. Not everyone who is given an EYDENZELT injection will experience a side effect.

Contact your doctor immediately if you have any signs or symptoms listed in the table below as these could be signs of a serious complication with the treatment:

Condition	Some Potential Signs or Symptoms
Infection or inflammation inside the eye	• Eye pain or increased discomfort • Worsening eye redness • Sensitivity to light • Swelling of the eyelid • Vision changes such as sudden decrease in vision or blurring of vision
Clouding of the lens (cataract)	• Blurry vision • Seeing shadows • Less vivid lines and shapes • Colour vision changes (e.g. colours look 'washed out')
Increase in pressure in the eye	• Seeing halos around lights • Eye pain • Experiencing a red eye • Nausea or vomiting • Vision changes
A detachment or tear of a layer of the retina	• Sudden flashes of light • A sudden appearance or an increase of floaters • A curtain like effect over a portion of the visual area • Vision changes

For a full list of side effects, please refer to the relevant EYDENZELT Patient Information Leaflet (PIL). You can access the PIL using the QR code at the beginning of this guide, or by asking your doctor.

If you get any side effects, which include any possible side effects not listed in this guide, talk to your doctor, pharmacist or nurse. You can also report side effects directly to the MHRA through the Yellow Card scheme, via the Yellow Card website www.mhra.gov.uk/yellowcard, the free Yellow Card app available from the Apple App Store or Google Play Store. Alternatively, you can call 0800 731 6789 for free, Monday to Friday between 9am and 5pm.

Adverse events can also be reported to Celltrion Healthcare UK by phone on +44 (0)1753 983500 or email at ukadverseevents@celltrionhc.com. By reporting side effects, you can help provide more information on the safety of this medicine.

What if I have concerns or questions?

If you have any concerns or questions, your doctor or nurse are the best people to speak to. They are very experienced and they know your individual situation so can provide you with the answers you need.

Useful contacts

The Macular Society

Helpline: 0300 3030 111

Address: The Macular Society, PO Box 1870, Andover SP10 9AD

Email: info@macularsociety.org

Website: www.macularsociety.org

RNIB (Royal National Institute of Blind People)

Helpline: 0303 123 9999

Address: Grimaldi Building, 154a Pentonville Road, London N1 9JE

Email: helpline@rnib.org.uk

Website: www.rnib.org.uk

Diabetes UK

Helpline: 0345 123 2399

Address: Wells Lawrence House, 126 Back Church Lane, London E1 1FH

Email: helpline@diabetes.org.uk

Website: www.diabetes.org.uk

Your eye care clinic is:

Contact:

Telephone:

Address:

Email:

Your guide to EYDENZELT®▼

EYDENZELT is used to treat myopic Choroidal Neovascularization (mCNV).

The guide has been produced for people who have been prescribed EYDENZELT (aflibercept solution for injection) for the treatment of mCNV. Your doctor will use the EYDENZELT 40 mg/ml solution for the recommended dose of 2mg aflibercept.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 of the SmPC for how to report adverse reactions.

The guide is available in printed and digital format, both versions allow you to access an audio version via a smart device. The audio version is a readout of the text in this patient guide. More information is available to you in the Patient Information Leaflet (PIL).

This information can be accessed by visiting <https://www.medicines.org.uk/emc> or by using the camera on a smart device to scan the relevant QR codes. This will direct the device to relevant information on the internet.

Patient Audio mCNV

QR code to Patient Audio mCNV



Patient Information Leaflet

Eydenzelt 40 mg/ml (2 mg dose)
QR code to Patient Information Leaflet



Your doctor has prescribed EYDENZELT because you have been diagnosed with myopic choroidal neovascularization, or mCNV.

What is EYDENZELT?

EYDENZELT is a type of treatment known as an anti-VEGF. Anti-VEGF is an abbreviation for anti-vascular endothelial growth factor, and this is a description of how EYDENZELT works to protect your vision. EYDENZELT blocks VEGF, and this action helps reduce the swelling in the retina and can lead to vision improvement and maintenance.

EYDENZELT is a solution (a liquid) that is injected into the eye. It is available in a 2 mg dose. Your doctor will recommend a treatment schedule for you and it is very important that you follow it.

EYDENZELT is not indicated for paediatric use unlike the original product Eylea.

What should your doctor know before you are treated with EYDENZELT?

Before your EYDENZELT treatment starts, make sure to tell your doctor and other health care providers if you:

- Have an infection in or around your eye.
- If you currently have redness in your eye or if there is any pain in your eye.
- Think you may be allergic to iodine, any pain killers or any of the ingredients in EYDENZELT.
- Have had any issues or problems with eye injections before.
- Have glaucoma or a history of high pressure in your eye.
- If you see, or have seen, flashes of light or 'floaters' in your vision.
- Are taking any medications, with or without a prescription.
- Had or will have eye surgery within 4 weeks before or after EYDENZELT treatment.
- Are pregnant, planning to become pregnant or breastfeeding. There is very little information about the safety of using EYDENZELT in pregnant women. EYDENZELT 2 mg should not be used during pregnancy, unless the benefit outweighs the risk to the foetus. Discuss this with your doctor before treatment with EYDENZELT. Women of childbearing potential should use effective contraception during their treatment with EYDENZELT. If you are treated with the EYDENZELT 2 mg dose, you should continue to use effective contraception for at least three months after the last injection. Small amounts of EYDENZELT may enter into the breastmilk. The effects of aflibercept on a breast-fed newborn/ infant is unknown. Use of EYDENZELT is not recommended during breastfeeding.

How can I get ready for my EYDENZELT appointment?

Your doctor may ask you to use eye drops for a few days before your appointment. After your treatment, your vision may be blurry so you should not drive home. Plan to have a friend or family member take you to your appointment or arrange another way to get there and home again. On the day of your appointment, do not wear any makeup.

What can I expect after the EYDENZELT treatment?

Your doctor may give you some eye tests after your EYDENZELT injection. This may include a test that measures the pressure inside your eye. After your injection, your vision will be blurry so you should not plan to drive until after your vision returns to normal.

During the next few days you may get a bloodshot eye or see moving spots in your vision. These should clear within a few days and if they don't, or if they get worse, you should contact your doctor.

Some people might feel a little bit of pain or discomfort in their eye after their injection. If this does not go away or gets worse, you should contact your doctor.

Does my treatment have side effects?

Just like any medicine, EYDENZELT has the potential to cause side effects. Not everyone who is given an EYDENZELT injection will experience a side effect.

Contact your doctor immediately if you have any signs or symptoms listed in the table below as these could be signs of a serious complication with the treatment:

Condition	Some Potential Signs or Symptoms
Infection or inflammation inside the eye	• Eye pain or increased discomfort • Worsening eye redness • Sensitivity to light • Swelling of the eyelid • Vision changes such as sudden decrease in vision or blurring of vision
Clouding of the lens (cataract)	• Blurry vision • Seeing shadows • Less vivid lines and shapes • Colour vision changes (e.g. colours look 'washed out')
Increase in pressure in the eye	• Seeing halos around lights • Eye pain • Experiencing a red eye • Nausea or vomiting • Vision changes
A detachment or tear of a layer of the retina	• Sudden flashes of light • A sudden appearance or an increase of floaters • A curtain like effect over a portion of the visual area • Vision changes

For a full list of side effects, please refer to the relevant EYDENZELT Patient Information Leaflet (PIL). You can access the PIL using the QR code at the beginning of this guide, or by asking your doctor.

If you get any side effects, which include any possible side effects not listed in this guide, talk to your doctor, pharmacist or nurse. You can also report side effects directly to the MHRA through the Yellow Card scheme, via the Yellow Card website www.mhra.gov.uk/yellowcard, the free Yellow Card app available from the Apple App Store or Google Play Store. Alternatively, you can call 0800 731 6789 for free, Monday to Friday between 9am and 5pm.

Adverse events can also be reported to Celltrion Healthcare UK by phone on +44 (0)1753 983500 or email at ukadverseevents@celltrionhc.com. By reporting side effects, you can help provide more information on the safety of this medicine.

What if I have concerns or questions?

If you have any concerns or questions, your doctor or nurse are the best people to speak to. They are very experienced and they know your individual situation so can provide you with the answers you need.

Useful contacts

The Macular Society

Helpline: 0300 3030 111

Address: The Macular Society, PO Box 1870, Andover SP10 9AD

Email: info@macularsociety.org

Website: www.macularsociety.org

Diabetes UK

Helpline: 0345 123 2399

Address: Wells Lawrence House, 126 Back Church Lane, London E1 1FH

Email: helpline@diabetes.org.uk

Website: www.diabetes.org.uk

RNIB (Royal National Institute of Blind People)

Helpline: 0303 123 9999

Address: Grimaldi Building, 154a Pentonville Road, London N1 9JE

Email: helpline@rnib.org.uk

Website: www.rnib.org.uk

Your eye care clinic is:

Contact:

Telephone:

Address:

Email: