

**Vgenfli<sup>▼</sup> 40 mg/mL**  
solution for injection  
**aflibercept**

# **Vgenfli<sup>▼</sup>** **PATIENT GUIDE**

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 of the SmPC for details on how to report adverse reactions.

TABLE OF CONTENTS	PAGE
<b>YOUR GUIDE TO VGENFLI FOR:</b>	
● <b>WET AGE-RELATED MACULAR DEGENERATION (WAMD) .....</b>	<b>1</b>
● <b>DIABETIC MACULAR OEDEMA (DMO) .....</b>	<b>7</b>
● <b>CENTRAL RETINAL VEIN OCCLUSION (CRVO) .....</b>	<b>13</b>
● <b>BRANCH RETINAL VEIN OCCLUSION (BRVO) .....</b>	<b>19</b>
● <b>MYOPIC CHOROIDAL NEOVASCULARISATION (MCNV) .....</b>	<b>25</b>

## **YOUR GUIDE TO Vgenfli<sup>▼</sup>**

### **Vgenfli is used to treat Wet Age-Related Macular Degeneration (Wet AMD)**

The guide has been produced for people who have been prescribed Vgenfli (aflibercept solution for injection) for the treatment of wAMD. The Vgenfli 2 mg dose has been studied in wAMD.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 of the SmPC for details on how to report adverse reactions.

The guide is available in printed and digital format, both versions allow you to access an audio version via a smart device. The audio version is a readout of the text in this patient guide. More information is available to you in the Patient Information Leaflet (PIL).

**This information can be accessed by visiting [www.medicines.org.uk/emc](http://www.medicines.org.uk/emc) or by using the camera on a smart device to scan the relevant QR codes. This will direct the device to relevant information on the internet.**

#### **Vgenfli 40 mg/mL (2 mg dose) (pre-filled syringe)**

QR code to  
**Patient Information Leaflet**



QR code to  
**Patient Audio wAMD**



# Your doctor has prescribed Vgenfli because you have been diagnosed with wet age-related macular degeneration, or Wet AMD.

## What is Vgenfli?

Vgenfli is a type of treatment known as an anti-VEGF. Anti-VEGF is an abbreviation for anti-vascular endothelial growth factor, and this is a description of how Vgenfli works to protect your vision. Vgenfli blocks VEGF, and this action helps reduce the swelling in the retina and can lead to vision improvement and maintenance.

Vgenfli is a solution (a liquid) that is injected into the eye. It is available in a 2 mg dose. Your doctor will recommend a treatment schedule for you and it is very important that you follow it.

## What should your doctor know before you are treated with Vgenfli?

Before your Vgenfli treatment starts, make sure to tell your doctor and other health care providers if you:

- Have an infection in or around your eye.
- Currently have redness in your eye or if there is any pain in your eye.
- Think you may be allergic to iodine, any pain killers or any of the ingredients in Vgenfli.
- Have had any issues or problems with eye injections before.
- Have glaucoma or a history of high pressure in your eye.
- See, or have seen, flashes of light or 'floaters' in your vision.
- Are taking any medications, with or without a prescription.
- Had or will have eye surgery within 4 weeks before or after Vgenfli treatment.
- Are pregnant, planning to become pregnant or breastfeeding. There is very little information about the safety of using Vgenfli in pregnant women. Vgenfli 2 mg should not be used during pregnancy, unless the benefit outweighs the risk to the foetus. Discuss this with your doctor before treatment with Vgenfli.

Women of childbearing potential should use effective contraception during their treatment with Vgenfli.

If you are treated with the Vgenfli 2 mg dose, you should continue to use effective contraception for at least three months after the last injection. Small amounts of Vgenfli may enter into the breastmilk. The effects of afibercept on a breast-fed newborn/infant is unknown. Use of Vgenfli is not recommended during breastfeeding.

## How can I get ready for my Vgenfli appointment?

Your doctor may ask you to use eye drops for a few days before your appointment. After your treatment, your vision may be blurry so you should not drive home. Plan to have a friend or family member take you to your appointment or arrange another way to get there and home again. On the day of your appointment, do not wear any makeup.

## What can I expect after the Vgenfli treatment?

Your doctor may give you some eye tests after your Vgenfli injection. This may include a test that measures the pressure inside your eye. After your injection, your vision will be blurry so you should not plan to drive until after your vision returns to normal.

If you use contact lenses, your doctor or nurse can advise you when you can start using them again.

During the next few days you may get a bloodshot eye or see moving spots in your vision. These should clear within a few days and if they don't, or if they get worse, you should contact your doctor.

Some people might feel a little bit of pain or discomfort in their eye after their injection. If this does not go away or gets worse, you should contact your doctor.

## Does Vgenfli have side effects?

Just like any medicine, Vgenfli has the potential to cause side effects. Not everyone who is given a Vgenfli injection will experience a side effect.

**Contact your doctor immediately if you have any signs or symptoms listed in the table below as these could be signs of a serious complication with the treatment:**

Condition	Some Potential Signs or Symptoms
<b>Infection or inflammation inside the eye</b>	<ul style="list-style-type: none"><li>• Eye pain or increased discomfort</li><li>• Worsening eye redness</li><li>• Sensitivity to light</li><li>• Swelling of the eyelid</li><li>• Vision changes such as sudden decrease in vision or blurring of vision</li></ul>
<b>Clouding of the lens (cataract)</b>	<ul style="list-style-type: none"><li>• Blurry vision</li><li>• Seeing shadows</li><li>• Less vivid lines and shapes</li><li>• Colour vision changes (e.g. colours looked 'washed out')</li></ul>
<b>Increase in pressure in the eye</b>	<ul style="list-style-type: none"><li>• Seeing halos around lights</li><li>• Eye pain</li><li>• Experiencing a red eye</li><li>• Nausea or vomiting</li><li>• Vision changes</li></ul>
<b>A detachment or tear of a layer of the retina</b>	<ul style="list-style-type: none"><li>• Sudden flashes of light</li><li>• A sudden appearance or an increase of floaters</li><li>• A curtain like effect over a portion of the visual area</li><li>• Vision changes</li></ul>

For a full list of side effects, please refer to the relevant Vgenfli Patient Information Leaflet (PIL). You can access the PIL using the QR code at the beginning of this guide, or by asking your doctor. The PIL is also available on [www.medicines.org.uk/emc](http://www.medicines.org.uk/emc)

**If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.**

**Adverse events can also be reported to the representative of the marketing authorization holder: Zakłady Farmaceutyczne POLPHARMA S.A.:  
MDBiologics UK Safety Department by phone on 0800 861 1298 or by e-mail at [drugsafety@mdbiologics.co.uk](mailto:drugsafety@mdbiologics.co.uk). By reporting side effects, you can help provide more information on the safety of this medicine.**

## What if I have concerns or questions?

If you have any concerns or questions, your doctor or nurse are the best people to speak to. They are very experienced and they know your individual situation so can provide you with the answers you need.

### Your eye care clinic is:

<b>Contact:</b>	
<b>Telephone:</b>	
<b>Address:</b>	
<b>Email:</b>	
<b>Clinic Stamp:</b>	



# **YOUR GUIDE TO Vgenfli<sup>▼</sup>**

**Vgenfli is used to treat Diabetic Macular Oedema (DMO).**

The guide has been produced for people who have been prescribed Vgenfli (aflibercept solution for injection) for the treatment of DMO. The Vgenfli 2 mg dose has been studied in DMO.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 of the SmPC for details on how to report adverse reactions.

The guide is available in printed and digital format, both versions allow you to access an audio version via a smart device. The audio version is a readout of the text in this patient guide. More information is available to you in the Patient Information Leaflet (PIL).

**This information can be accessed by visiting [www.medicines.org.uk/emc](http://www.medicines.org.uk/emc) or by using the camera on a smart device to scan the relevant QR codes. This will direct the device to relevant information on the internet.**

---

## **Vgenfli 40 mg/mL (2 mg dose) (pre-filled syringe)**

---

QR code to  
**Patient Information Leaflet**



QR code to  
**Patient Audio DMO**



# Your doctor has prescribed Vgenfli because you have been diagnosed with diabetic macular oedema, or DMO for short.

## What is Vgenfli?

Vgenfli is a type of treatment known as an anti-VEGF. Anti-VEGF is an abbreviation for anti-vascular endothelial growth factor, and this is a description of how Vgenfli works to protect your vision. Vgenfli blocks VEGF, and this action helps reduce the swelling in the retina and can lead to vision improvement and maintenance.

Vgenfli is a solution (a liquid) that is injected into the eye. It is available in a 2 mg dose. Your doctor will recommend a treatment schedule for you and it is very important that you follow it.

## What should your doctor know before you are treated with Vgenfli?

Before your Vgenfli treatment starts, make sure to tell your doctor and other health care providers if you:

- Have an infection in or around your eye.
- Currently have redness in your eye or if there is any pain in your eye.
- Think you may be allergic to iodine, any pain killers or any of the ingredients in Vgenfli.
- Have had any issues or problems with eye injections before.
- Have glaucoma or a history of high pressure in your eye.
- See, or have seen, flashes of light or 'floaters' in your vision.
- Are taking any medications, with or without a prescription.
- Had or will have eye surgery within 4 weeks before or after Vgenfli treatment.
- Are pregnant, planning to become pregnant or breastfeeding. There is very little information about the safety of using Vgenfli in pregnant women. Vgenfli 2 mg should not be used during pregnancy, unless the benefit outweighs the risk to the foetus. Discuss this with your doctor before treatment with Vgenfli.

Women of childbearing potential should use effective contraception during their treatment with Vgenfli.

If you are treated with the Vgenfli 2 mg dose, you should continue to use effective contraception for at least three months after the last injection. Small amounts of Vgenfli may enter into the breastmilk. The effects of afibercept on a breast-fed newborn/infant is unknown. Use of Vgenfli is not recommended during breastfeeding.

## How can I get ready for my Vgenfli appointment?

Your doctor may ask you to use eye drops for a few days before your appointment. After your treatment, your vision may be blurry so you should not drive home. Plan to have a friend or family member take you to your appointment or arrange another way to get there and home again. On the day of your appointment, do not wear any makeup.

## What can I expect after the Vgenfli treatment?

Your doctor may give you some eye tests after your Vgenfli injection. This may include a test that measures the pressure inside your eye. After your injection, your vision will be blurry so you should not plan to drive until after your vision returns to normal.

If you use contact lenses, your doctor or nurse can advise you when you can start using them again.

During the next few days you may get a bloodshot eye or see moving spots in your vision. These should clear within a few days and if they don't, or if they get worse, you should contact your doctor.

Some people might feel a little bit of pain or discomfort in their eye after their injection. If this does not go away or gets worse, you should contact your doctor.

## Does Vgenfli have side effects?

Just like any medicine, Vgenfli has the potential to cause side effects. Not everyone who is given a Vgenfli injection will experience a side effect.

**Contact your doctor immediately if you have any signs or symptoms listed in the table below as these could be signs of a serious complication with the treatment:**

Condition	Some Potential Signs or Symptoms
<b>Infection or inflammation inside the eye</b>	<ul style="list-style-type: none"><li>• Eye pain or increased discomfort</li><li>• Worsening eye redness</li><li>• Sensitivity to light</li><li>• Swelling of the eyelid</li><li>• Vision changes such as sudden decrease in vision or blurring of vision</li></ul>
<b>Clouding of the lens (cataract)</b>	<ul style="list-style-type: none"><li>• Blurry vision</li><li>• Seeing shadows</li><li>• Less vivid lines and shapes</li><li>• Colour vision changes (e.g. colours looked 'washed out')</li></ul>
<b>Increase in pressure in the eye</b>	<ul style="list-style-type: none"><li>• Seeing halos around lights</li><li>• Eye pain</li><li>• Experiencing a red eye</li><li>• Nausea or vomiting</li><li>• Vision changes</li></ul>
<b>A detachment or tear of a layer of the retina</b>	<ul style="list-style-type: none"><li>• Sudden flashes of light</li><li>• A sudden appearance or an increase of floaters</li><li>• A curtain like effect over a portion of the visual area</li><li>• Vision changes</li></ul>

For a full list of side effects, please refer to the relevant Vgenfli Patient Information Leaflet (PIL). You can access the PIL using the QR code at the beginning of this guide, or by asking your doctor. The PIL is also available on [www.medicines.org.uk/emc](http://www.medicines.org.uk/emc)

**If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.**

**Adverse events can also be reported to the representative of the marketing authorization holder: Zakłady Farmaceutyczne POLPHARMA S.A.:  
MDBiologics UK Safety Department by phone on 0800 861 1298 or by e-mail at [drugsafety@mdbiologics.co.uk](mailto:drugsafety@mdbiologics.co.uk). By reporting side effects, you can help provide more information on the safety of this medicine.**

## What if I have concerns or questions?

If you have any concerns or questions, your doctor or nurse are the best people to speak to. They are very experienced and they know your individual situation so can provide you with the answers you need.

### Your eye care clinic is:

<b>Contact:</b>	
<b>Telephone:</b>	
<b>Address:</b>	
<b>Email:</b>	
<b>Clinic Stamp:</b>	



## **YOUR GUIDE TO Vgenfli<sup>▼</sup>**

**Vgenfli is used to treat macular oedema secondary to Central Retinal Vein Occlusion (CRVO).**

The guide has been produced for people who have been prescribed the Vgenfli 2 mg dose (aflibercept solution for injection) for the treatment of CRVO.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 of the SmPC for details on how to report adverse reactions.

The guide is available in printed and digital format, both versions allow you to access an audio version via a smart device. The audio version is a readout of the text in this patient guide. More information is available to you in the Patient Information Leaflet (PIL).

**This information can be accessed by visiting [www.medicines.org.uk/emc](http://www.medicines.org.uk/emc) or by using the camera on a smart device to scan the relevant QR codes. This will direct the device to relevant information on the internet.**

---

### **Vgenfli 40 mg/mL (2 mg dose) (pre-filled syringe)**

---

QR code to  
**Patient Information Leaflet**



QR code to  
**Patient Audio CRVO**



# Your doctor has prescribed Vgenfli because you have been diagnosed with macular oedema secondary to central retinal vein occlusion, or CRVO.

## What is Vgenfli?

Vgenfli is a type of treatment known as an anti-VEGF. Anti-VEGF is an abbreviation for anti-vascular endothelial growth factor, and this is a description of how Vgenfli works to protect your vision. Vgenfli blocks VEGF, and this action helps reduce the swelling in the retina and can lead to vision improvement and maintenance.

Vgenfli is a solution (a liquid) that is injected into the eye. For the treatment of CRVO, Vgenfli is available in a 2 mg dose. Your doctor will also recommend a treatment schedule for you and it is very important that you follow it.

## What should your doctor know before you are treated with Vgenfli?

Before your Vgenfli treatment starts, make sure to tell your doctor and other health care providers if you:

- Have an infection in or around your eye.
- Currently have redness in your eye or if there is any pain in your eye.
- Think you may be allergic to iodine, any pain killers or any of the ingredients in Vgenfli.
- Have had any issues or problems with eye injections before.
- Have glaucoma or a history of high pressure in your eye.
- See, or have seen, flashes of light or 'floaters' in your vision.
- Are taking any medications, with or without a prescription.
- Had or will have eye surgery within 4 weeks before or after Vgenfli treatment.
- Are pregnant, planning to become pregnant or breastfeeding. There is very little information about the safety of using Vgenfli in pregnant women. Vgenfli 2 mg should not be used during pregnancy, unless the benefit outweighs the risk to the foetus. Discuss this with your doctor before treatment with Vgenfli.

Women of childbearing potential should use effective contraception during their treatment and for at least three months after the last injection of Vgenfli. Small amounts of Vgenfli may enter into the breastmilk. The effects of afibercept on a breast-fed newborn/infant is unknown. Use of Vgenfli is not recommended during breastfeeding.

## How can I get ready for my Vgenfli appointment?

Your doctor may ask you to use eye drops for a few days before your appointment. After your treatment, your vision may be blurry so you should not drive home. Plan to have a friend or family member take you to your appointment or arrange another way to get there and home again. On the day of your appointment, do not wear any makeup.

## What can I expect after the Vgenfli treatment?

Your doctor may give you some eye tests after your Vgenfli injection. This may include a test that measures the pressure inside your eye. After your injection, your vision will be blurry so you should not plan to drive until after your vision returns to normal.

If you use contact lenses, your doctor or nurse can advise you when you can start using them again.

During the next few days you may get a bloodshot eye or see moving spots in your vision. These should clear within a few days and if they don't, or if they get worse, you should contact your doctor.

Some people might feel a little bit of pain or discomfort in their eye after their injection. If this does not go away or gets worse, you should contact your doctor.

## Does Vgenfli have side effects?

Just like any medicine, Vgenfli has the potential to cause side effects. Not everyone who is given a Vgenfli injection will experience a side effect.

**Contact your doctor immediately if you have any signs or symptoms listed in the table below as these could be signs of a serious complication with the treatment:**

Condition	Some Potential Signs or Symptoms
<b>Infection or inflammation inside the eye</b>	<ul style="list-style-type: none"><li>• Eye pain or increased discomfort</li><li>• Worsening eye redness</li><li>• Sensitivity to light</li><li>• Swelling of the eyelid</li><li>• Vision changes such as sudden decrease in vision or blurring of vision</li></ul>
<b>Clouding of the lens (cataract)</b>	<ul style="list-style-type: none"><li>• Blurry vision</li><li>• Seeing shadows</li><li>• Less vivid lines and shapes</li><li>• Colour vision changes (e.g. colours looked 'washed out')</li></ul>
<b>Increase in pressure in the eye</b>	<ul style="list-style-type: none"><li>• Seeing halos around lights</li><li>• Eye pain</li><li>• Experiencing a red eye</li><li>• Nausea or vomiting</li><li>• Vision changes</li></ul>
<b>A detachment or tear of a layer of the retina</b>	<ul style="list-style-type: none"><li>• Sudden flashes of light</li><li>• A sudden appearance or an increase of floaters</li><li>• A curtain like effect over a portion of the visual area</li><li>• Vision changes</li></ul>

For a full list of side effects, please refer to the relevant Vgenfli Patient Information Leaflet (PIL). You can access the PIL using the QR code at the beginning of this guide, or by asking your doctor. The PIL is also available on [www.medicines.org.uk/emc](http://www.medicines.org.uk/emc)

**If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.**

**Adverse events can also be reported to the representative of the marketing authorization holder: Zakłady Farmaceutyczne POLPHARMA S.A.:  
MDBiologics UK Safety Department by phone on 0800 861 1298 or by e-mail at [drugsafety@mdbiologics.co.uk](mailto:drugsafety@mdbiologics.co.uk). By reporting side effects, you can help provide more information on the safety of this medicine.**

## What if I have concerns or questions?

If you have any concerns or questions, your doctor or nurse are the best people to speak to. They are very experienced and they know your individual situation so can provide you with the answers you need.

### Your eye care clinic is:

**Contact:**

**Telephone:**

**Address:**

**Email:**

**Clinic Stamp:**



## **YOUR GUIDE TO Vgenfli<sup>▼</sup>**

**Vgenfli is used to treat macular oedema secondary to Branch Retinal Vein Occlusion (BRVO)**

The guide has been produced for people who have been prescribed the Vgenfli 2 mg dose (aflibercept solution for injection) for the treatment of BRVO.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 of the SmPC for details on how to report adverse reactions.

The guide is available in printed and digital format, both versions allow you to access an audio version via a smart device. The audio version is a readout of the text in this patient guide. More information is available to you in the Patient Information Leaflet (PIL).

**This information can be accessed by visiting [www.medicines.org.uk/emc](http://www.medicines.org.uk/emc) or by using the camera on a smart device to scan the relevant QR codes. This will direct the device to relevant information on the internet.**

---

### **Vgenfli 40 mg/mL (2 mg dose) (pre-filled syringe)**

---

QR code to  
**Patient Information Leaflet**



QR code to  
**Patient Audio BRVO**



# Your doctor has prescribed Vgenfli because you have been diagnosed with macular oedema secondary to branch retinal vein occlusion, or BRVO.

## What is Vgenfli?

Vgenfli is a type of treatment known as an anti-VEGF. Anti-VEGF is an abbreviation for anti-vascular endothelial growth factor, and this is a description of how Vgenfli works to protect your vision. Vgenfli blocks VEGF, and this action helps reduce the swelling in the retina and can lead to vision improvement and maintenance.

Vgenfli is a solution (a liquid) that is injected into the eye. For the treatment of BRVO, Vgenfli is available in a 2 mg dose. Your doctor will also recommend a treatment schedule for you and it is very important that you follow it.

## What should your doctor know before you are treated with Vgenfli?

Before your Vgenfli treatment starts, make sure to tell your doctor and other health care providers if you:

- Have an infection in or around your eye.
- Currently have redness in your eye or if there is any pain in your eye.
- Think you may be allergic to iodine, any pain killers or any of the ingredients in Vgenfli.
- Have had any issues or problems with eye injections before.
- Have glaucoma or a history of high pressure in your eye.
- See, or have seen, flashes of light or 'floaters' in your vision.
- Are taking any medications, with or without a prescription.
- Had or will have eye surgery within 4 weeks before or after Vgenfli treatment.
- Are pregnant, planning to become pregnant or breastfeeding. There is very little information about the safety of using Vgenfli in pregnant women. Vgenfli 2 mg should not be used during pregnancy, unless the benefit outweighs the risk to the foetus. Discuss this with your doctor before treatment with Vgenfli.

Women of childbearing potential should use effective contraception during their treatment and for at least three months after the last injection of Vgenfli. Small amounts of Vgenfli may enter into the breastmilk. The effects of afibercept on a breast-fed newborn/infant is unknown. Use of Vgenfli is not recommended during breastfeeding.

## How can I get ready for my Vgenfli appointment?

Your doctor may ask you to use eye drops for a few days before your appointment. After your treatment, your vision may be blurry so you should not drive home. Plan to have a friend or family member take you to your appointment or arrange another way to get there and home again. On the day of your appointment, do not wear any makeup.

## What can I expect after the Vgenfli treatment?

Your doctor may give you some eye tests after your Vgenfli injection. This may include a test that measures the pressure inside your eye. After your injection, your vision will be blurry so you should not plan to drive until after your vision returns to normal.

If you use contact lenses, your doctor or nurse can advise you when you can start using them again.

During the next few days you may get a bloodshot eye or see moving spots in your vision. These should clear within a few days and if they don't, or if they get worse, you should contact your doctor.

Some people might feel a little bit of pain or discomfort in their eye after their injection. If this does not go away or gets worse, you should contact your doctor.

## Does Vgenfli have side effects?

Just like any medicine, Vgenfli has the potential to cause side effects. Not everyone who is given a Vgenfli injection will experience a side effect.

**Contact your doctor immediately if you have any signs or symptoms listed in the table below as these could be signs of a serious complication with the treatment:**

Condition	Some Potential Signs or Symptoms
<b>Infection or inflammation inside the eye</b>	<ul style="list-style-type: none"><li>• Eye pain or increased discomfort</li><li>• Worsening eye redness</li><li>• Sensitivity to light</li><li>• Swelling of the eyelid</li><li>• Vision changes such as sudden decrease in vision or blurring of vision</li></ul>
<b>Clouding of the lens (cataract)</b>	<ul style="list-style-type: none"><li>• Blurry vision</li><li>• Seeing shadows</li><li>• Less vivid lines and shapes</li><li>• Colour vision changes (e.g. colours looked 'washed out')</li></ul>
<b>Increase in pressure in the eye</b>	<ul style="list-style-type: none"><li>• Seeing halos around lights</li><li>• Eye pain</li><li>• Experiencing a red eye</li><li>• Nausea or vomiting</li><li>• Vision changes</li></ul>
<b>A detachment or tear of a layer of the retina</b>	<ul style="list-style-type: none"><li>• Sudden flashes of light</li><li>• A sudden appearance or an increase of floaters</li><li>• A curtain like effect over a portion of the visual area</li><li>• Vision changes</li></ul>

For a full list of side effects, please refer to the relevant Vgenfli Patient Information Leaflet (PIL). You can access the PIL using the QR code at the beginning of this guide, or by asking your doctor. The PIL is also available on [www.medicines.org.uk/emc](http://www.medicines.org.uk/emc)

**If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.**

**Adverse events can also be reported to the representative of the marketing authorization holder: Zakłady Farmaceutyczne POLPHARMA S.A.:  
MDBiologics UK Safety Department by phone on 0800 861 1298 or by e-mail at [drugsafety@mdbiologics.co.uk](mailto:drugsafety@mdbiologics.co.uk). By reporting side effects, you can help provide more information on the safety of this medicine.**

## What if I have concerns or questions?

If you have any concerns or questions, your doctor or nurse are the best people to speak to. They are very experienced and they know your individual situation so can provide you with the answers you need.

### Your eye care clinic is:

**Contact:**

**Telephone:**

**Address:**

**Email:**

**Clinic Stamp:**



# **YOUR GUIDE TO Vgenfli<sup>▼</sup>**

**Vgenfli is used to treat myopic Choroidal Neovascularisation (mCNV).**

The guide has been produced for people who have been prescribed the Vgenfli 2 mg dose (aflibercept solution for injection) for the treatment of mCNV.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 of the SmPC for details on how to report adverse reactions.

The guide is available in printed and digital format, both versions allow you to access an audio version via a smart device. The audio version is a readout of the text in this patient guide. More information is available to you in the Patient Information Leaflet (PIL).

**This information can be accessed by visiting [www.medicines.org.uk/emc](http://www.medicines.org.uk/emc) or by using the camera on a smart device to scan the relevant QR codes. This will direct the device to relevant information on the internet.**

---

## **Vgenfli 40 mg/mL (2 mg dose) (pre-filled syringe)**

---

QR code to  
**Patient Information Leaflet**



QR code to  
**Patient Audio Myopic CNV**



# Your doctor has prescribed Vgenfli because you have been diagnosed with myopic choroidal neovascularisation, or mCNV.

## What is Vgenfli?

Vgenfli is a type of treatment known as an anti-VEGF. Anti-VEGF is an abbreviation for anti-vascular endothelial growth factor, and this is a description of how Vgenfli works to protect your vision. Vgenfli blocks VEGF, and this action helps reduce the swelling in the retina and can lead to vision improvement and maintenance.

Vgenfli is a solution (a liquid) that is injected into the eye. For the treatment of mCNV, Vgenfli is available in a 2 mg dose. Your doctor will also recommend a treatment schedule for you and it is very important that you follow it.

## What should your doctor know before you are treated with Vgenfli?

Before your Vgenfli treatment starts, make sure to tell your doctor and other health care providers if you:

- Have an infection in or around your eye.
- Currently have redness in your eye or if there is any pain in your eye.
- Think you may be allergic to iodine, any pain killers or any of the ingredients in Vgenfli.
- Have had any issues or problems with eye injections before.
- Have glaucoma or a history of high pressure in your eye.
- See, or have seen, flashes of light or 'floaters' in your vision.
- Are taking any medications, with or without a prescription.
- Had or will have eye surgery within 4 weeks before or after Vgenfli treatment.
- Are pregnant, planning to become pregnant or breastfeeding. There is very little information about the safety of using Vgenfli in pregnant women. Vgenfli 2 mg should not be used during pregnancy, unless the benefit outweighs the risk to the foetus. Discuss this with your doctor before treatment with Vgenfli.

Women of childbearing potential should use effective contraception during their treatment and for at least three months after the last injection of Vgenfli. Small amounts of Vgenfli may enter into the breastmilk. The effects of afibercept on a breast-fed newborn/infant is unknown. Use of Vgenfli is not recommended during breastfeeding.

## How can I get ready for my Vgenfli appointment?

Your doctor may ask you to use eye drops for a few days before your appointment. After your treatment, your vision may be blurry so you should not drive home. Plan to have a friend or family member take you to your appointment or arrange another way to get there and home again. On the day of your appointment, do not wear any makeup.

## What can I expect after the Vgenfli treatment?

Your doctor may give you some eye tests after your Vgenfli injection. This may include a test that measures the pressure inside your eye. After your injection, your vision will be blurry so you should not plan to drive until after your vision returns to normal.

If you use contact lenses, your doctor or nurse can advise you when you can start using them again.

During the next few days you may get a bloodshot eye or see moving spots in your vision. These should clear within a few days and if they don't, or if they get worse, you should contact your doctor.

Some people might feel a little bit of pain or discomfort in their eye after their injection. If this does not go away or gets worse, you should contact your doctor.

## Does Vgenfli have side effects?

Just like any medicine, Vgenfli has the potential to cause side effects. Not everyone who is given a Vgenfli injection will experience a side effect.

**Contact your doctor immediately if you have any signs or symptoms listed in the table below as these could be signs of a serious complication with the treatment:**

Condition	Some Potential Signs or Symptoms
<b>Infection or inflammation inside the eye</b>	<ul style="list-style-type: none"><li>• Eye pain or increased discomfort</li><li>• Worsening eye redness</li><li>• Sensitivity to light</li><li>• Swelling of the eyelid</li><li>• Vision changes such as sudden decrease in vision or blurring of vision</li></ul>
<b>Clouding of the lens (cataract)</b>	<ul style="list-style-type: none"><li>• Blurry vision</li><li>• Seeing shadows</li><li>• Less vivid lines and shapes</li><li>• Colour vision changes (e.g. colours looked 'washed out')</li></ul>
<b>Increase in pressure in the eye</b>	<ul style="list-style-type: none"><li>• Seeing halos around lights</li><li>• Eye pain</li><li>• Experiencing a red eye</li><li>• Nausea or vomiting</li><li>• Vision changes</li></ul>
<b>A detachment or tear of a layer of the retina</b>	<ul style="list-style-type: none"><li>• Sudden flashes of light</li><li>• A sudden appearance or an increase of floaters</li><li>• A curtain like effect over a portion of the visual area</li><li>• Vision changes</li></ul>

For a full list of side effects, please refer to the relevant Vgenfli Patient Information Leaflet (PIL). You can access the PIL using the QR code at the beginning of this guide, or by asking your doctor. The PIL is also available on [www.medicines.org.uk/emc](http://www.medicines.org.uk/emc)

**If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.**

**Adverse events can also be reported to the representative of the marketing authorization holder: Zakłady Farmaceutyczne POLPHARMA S.A.:  
MDBiologics UK Safety Department by phone on 0800 861 1298 or by e-mail at [drugsafety@mdbiologics.co.uk](mailto:drugsafety@mdbiologics.co.uk). By reporting side effects, you can help provide more information on the safety of this medicine.**

## What if I have concerns or questions?

If you have any concerns or questions, your doctor or nurse are the best people to speak to. They are very experienced and they know your individual situation so can provide you with the answers you need.

### Your eye care clinic is:

<b>Contact:</b>	
<b>Telephone:</b>	
<b>Address:</b>	
<b>Email:</b>	
<b>Clinic Stamp:</b>	

For more information about educational materials for Vgenfli, visit:  
[www.medicines.org.uk/emc](http://www.medicines.org.uk/emc)



Zakłady Farmaceutyczne Polpharma S.A.  
Bobrowiecka St. 6, 00-728 Warsaw, POLAND  
Date of version: July 2025