

This card has been developed by Pfizer Ltd and is intended only for patients who have been prescribed HYMPAVZI (marstacimab) in the UK.

HYMPAVZI[®] ▼ (marstacimab) Patient Card

Important safety information for patients receiving treatment with HYMPAVZI. This information does not replace the Patient Information Leaflet that comes with your medicine. Please read it carefully.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. For more information, see the section on reporting side effects.



Patient Information

Always carry this card with you and show it to healthcare professionals (doctor or nurse) whenever you have a medical appointment or emergency.

What is HYMPAVZI used for?

HYMPAVZI is a medicine used to prevent or reduce bleeding in patients 12 years of age and older, weighing at least 35 kg, with:

- severe haemophilia A who have not developed factor VIII inhibitors, or
- severe haemophilia B who have not developed factor IX inhibitors.

Blood clots (thromboembolic events)

HYMPAVZI increases how easily your blood clots and may cause blood clots in the blood vessels (so called thromboembolic events). Blood clots may be life threatening. Tell your doctor if you have a history of blood clots or if you have a condition which increases your risk of blood clots.

Stop using HYMPAVZI, and talk to a doctor immediately if you notice any symptoms of a possible blood clot - see further details on the reverse.

Reporting of Side Effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet.

You can also report side effects directly via the Yellow Card Scheme at <https://yellowcard.mhra.gov.uk> and the free Yellow Card app, available from the Google Play Store or Apple App Store. Alternatively you can call 0800 731 6789 for free, Monday to Friday between 9am and 5pm.

By reporting side effects, you can help provide more information on the safety of this medicine.



Personal Details

Name of patient:

Date of starting HYMPAVZI & batch number:

Doctor's name & contact details:

Emergency contact & contact details:

Blood clots (thromboembolic events)

Stop using HYMPAVZI, and talk to a doctor immediately if you notice any symptoms of a possible blood clot including the following:

- swelling or pain in arms or legs
- redness or discolouration in your arms or legs
- shortness of breath
- pain in chest or upper back
- fast heart rate
- coughing up blood
- feeling faint
- headache
- numbness in your face
- eye pain or swelling
- trouble seeing