

Pregnancy: IV iron should not be used during pregnancy unless clearly necessary. If you are pregnant or think you could be pregnant, it is important to discuss this with your doctor.

You should contact your Doctor or Nurse immediately if:

- you have any signs or symptoms of an allergic reaction during or shortly after treatment with IV Iron

For example: hives or rash, itching, dizziness, light-headedness, swelling of the lips, tongue, throat or body, difficulty breathing, shortness of breath or wheezing.

Your doctor and nursing staff are aware of these possible side effects and you will be monitored during and after you receive IV iron.

Reporting of side effects

Please report side effects via the Yellow Card Scheme at <https://yellowcard.mhra.gov.uk/> or search for MHRA Yellow Card in the Google Play or Apple App Store.

Alternatively, you can call 0800 731 6789 (freephone), Monday to Friday, 9am–5pm.

When reporting, please ensure you include the name of the specific product administered.

By reporting side effects, you can help provide more information on the safety of this medicine.

Intravenous (IV) Iron

**Important Information for Patients
about the Possible Risk of Serious
Allergic Reactions with IV iron
treatment (iron given by injection
or infusion into a vein)**

ferric carboxymaltose, iron dextran, ferric
derisomaltose, iron sucrose

**Please read this leaflet carefully and
discuss any questions you may have
with your Doctor.**

**You should also read the patient
information leaflet that comes with
the particular IV iron that you have
been prescribed. This contains
important additional information
regarding the medicine.**

IV Iron is used to treat iron deficiency when iron preparations taken by mouth have not worked or cannot be used. The aim of treatment is to replenish the body's iron stores

IV Iron can cause allergic reactions. In a small number of patients these allergic reactions can become severe or life-threatening (known as anaphylactic reactions) and can cause problems with your heart and blood pressure and/or cause you to faint or lose consciousness.

You should not be prescribed or given an IV Iron if:

- you are allergic (hypersensitive) to the product or any of the other ingredients of this medicine
- you have experienced serious allergic (hypersensitive) reactions to other IV iron treatments in the past*
- you have iron overload (too much iron in your body)
- your anaemia is not caused by iron deficiency

You should tell your doctor before they prescribe / administer an IV Iron if you have any of these allergies or conditions.

*** It is important to know that a reaction can still happen even if you have not had any problems in the past with IV iron.**

You may have an increased risk of having an allergic reaction if you have:

- known allergies including drug allergies
- a history of severe asthma, eczema or other allergies (for example dust, pollen, pet dander) or
- immune or inflammatory conditions (e.g. rheumatoid arthritis, lupus erythematosus)

You should tell your doctor before they prescribe or give you IV Iron if you have any of these allergies or conditions.

Your Doctor will decide whether IV iron is suitable for you.