

Front

You should immediately contact your doctor if you think you are having a hypersensitivity reaction to Abacavir/Lamivudine. Write your doctor's details below:

Doctor:

Tel:

If your doctor is not available, you must urgently seek alternative medical advice (e.g. the emergency unit of the nearest hospital).

For general Abacavir/Lamivudine information enquiries, contact Mylan, Tel: + 44 1707 853000.

IMPORTANT - ALERT CARD

Abacavir/Lamivudine tablets
Carry this card with you at all times

Back

Since Abacavir/Lamivudine contains abacavir some patients taking Abacavir/Lamivudine may develop a hypersensitivity reaction (serious allergic reaction) which **can be life-threatening** if treatment with Abacavir/Lamivudine is continued.

CONTACT YOUR DOCTOR IMMEDIATELY for advice on whether you should stop taking Abacavir/Lamivudine if:

- 1) you get a skin rash OR**
- 2) you get one or more symptoms from at least TWO of the following groups**

- fever
- shortness of breath, sore throat or cough
- nausea or vomiting or diarrhoea or abdominal pain
- severe tiredness or achiness or generally feeling ill.

If you have discontinued Abacavir/Lamivudine due to this reaction, **YOU MUST NEVER TAKE** Abacavir/Lamivudine, or any medicine containing abacavir (e.g. Ziagen, Trizivir, Triumeq) again as **within hours** you may experience a life-threatening lowering of your blood pressure or death.

(see reverse of card)

792372

 Mylan