

This card contains important safety information that you need to be aware of before you are given Benepali and during treatment with Benepali.

If you do not understand this information, please ask your doctor to explain it to you. This medicine is for adults aged 18 and over.

- Show this card to any doctor involved in your treatment.

Use in children and adolescents (less than 18 years old)

Benepali is not indicated for use in children and adolescents. Please ask your doctor if you have any questions about this.

Infections

Benepali may increase your risk of getting infections, which could be serious.

- You should not use Benepali if you have an infection. If you are not sure, ask your doctor.
- If you develop symptoms suggestive of infections, such as fever, persistent cough, weight loss, or listlessness, seek medical attention immediately.
- You should be evaluated for tuberculosis (TB). Ask your doctor to record the dates and the results of your last screening for TB below:

Test:
Date:
Results:


Test:
Date:
Results:



STB-144372-01



48005-01

<p>Other information (please complete)</p> <p>Patient's name: -----</p> <p>Doctor's name: -----</p> <p>Doctor's phone: -----</p> <ul style="list-style-type: none">• Keep this card with you for 2 months after the last Benepali dose, since side effects may occur after your last dose of Benepali.	<ul style="list-style-type: none">• Please ask your physician to list your other medicines that may increase your risk of infection. _____ _____ <p>Congestive heart failure</p> <ul style="list-style-type: none">• If you have congestive heart failure, and you think your symptoms (e.g., shortness of breath or swelling of the feet) are becoming worse, seek medical attention immediately.	<p>Patient Alert Card</p> <p> Benepali[®] 50 mg</p> <p>Etanercept</p>
---	---	---