

Acitretin : Pregnancy Prevention Programme

You have been given this card because you have been prescribed Acitretin

What you must do

Please read this card and the package information leaflet thoroughly before starting your treatment with acitretin.

- You must not become pregnant while taking acitretin, or for 3 years after stopping treatment.
- You must use effective contraception before, during and for 3 years after stopping treatment with acitretin.
- You must attend regular follow-up visits and have regular pregnancy testing.

Reminder for Men and Women

***Do not share this medication with anybody and return any unused capsules back to the pharmacy.
You should not donate blood during treatment with this medicine and for three years after stopping treatment.***

Acitretin [▼]: Pregnancy Prevention Programme

You have been given this card because you have been prescribed Acitretin

Important information to know

- Acitretin must not be taken during pregnancy.
- Acitretin can seriously harm an unborn baby if a pregnant women takes it.
- If you become pregnant or think you might be pregnant, stop taking acitretin immediately and contact your doctor.

If you have any questions or concerns about taking acitretin, talk to your doctor or pharmacist.

[▼] This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See www.mhra.gov.uk/yellowcard for how to report side effects or search for MHRA Yellow Card in the Google Play or Apple App Store.

Keep this card safe so you always know what to do
UK/MED/18/0197a

Date of Preparation: May 2019