

My name: \_\_\_\_\_

My contact number: \_\_\_\_\_

Emergency contact: \_\_\_\_\_

Emergency contact number: \_\_\_\_\_

Name of Haematologist/  
Oncologist/Oncology Nurse: \_\_\_\_\_

Contact number: \_\_\_\_\_

After-hours contact number: \_\_\_\_\_

Name of my Hospital: \_\_\_\_\_

My Hospital contact number: \_\_\_\_\_

PATIENT ALERT CARD


GILTERITINIB  
ASTELLAS


- Carry this card with you **at all times**, especially when you travel or when you see another doctor.
- Please ensure you show this card to any doctor, pharmacist or nurse for any medical treatment or at any visits to the hospital or clinic.
- Please contact your doctor **immediately**, if you develop any side effects, in particular those listed on this card.

171090



4008014

4008014/ 171090
GILTERITINIB ASTELLAS
CRD 40
United Kingdom GB
106 X 115 mm FD
CRD_002
Project 91376
Eq. to: N/A
Proof_01
09 Jul 2025 19:52
 Perigord life science artwork solutions

Black	PMS 7680 C
	Varnish free

## IMPORTANT SAFETY INFORMATION FOR PATIENTS

**Gilteritinib Astellas may cause serious side effects, including differentiation syndrome.**

Differentiation syndrome is a condition that affects your blood cells and may be life-threatening or lead to death if not treated in a timely manner.

Talk to your doctor, pharmacist or nurse **immediately** if you have any of the following symptoms:

- Fever
- Trouble Breathing
- Rash
- Dizziness or lightheadedness
- Rapid weight gain
- Swelling of your arms or legs

Differentiation syndrome can happen any time during the first 3 months of treatment from as early as 1 day after starting treatment. Getting medical treatment early may stop the problem from becoming more serious.

Your doctor will monitor you, may pause your treatment and/or may give you a medicine to treat your condition.

If you have any further questions about your treatment, please contact your doctor.

## IMPORTANT INFORMATION FOR HEALTHCARE PROVIDERS

- This patient is being treated with Gilteritinib Astellas, which can cause differentiation syndrome.
- Symptoms include fever, dyspnoea, pleural effusion, pericardial effusion, pulmonary oedema, hypotension, rapid weight gain, peripheral oedema, rash, and renal dysfunction.
- If differentiation syndrome is suspected, corticosteroid therapy should be initiated along with hemodynamic monitoring until symptom resolution.
- If severe signs and/or symptoms persist for more than 48 hours after initiation of corticosteroids, Gilteritinib Astellas should be interrupted until signs and symptoms are no longer severe.

Please contact the patient's Haematologist/Oncologist for more information and consult the Product Information for gilteritinib available at [www.medicines.org.uk](http://www.medicines.org.uk).