

Show this card to any healthcare professional involved in your medical care – for example, your dentist or an emergency doctor.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See <https://yellowcard.mhra.gov.uk> for how to report side effects or via the Yellow Card app from the Google Play or Apple app Store.

Your name:

.....
Consultant's name – who prescribed RINVOQ®:

.....
Consultant's phone number:

.....
The date you started RINVOQ®:

www.gov.uk/mhra

UK-UPAD-190110/October 2019



abbvie

Patient Alert Card

Keep this card with you all the time

Safety Information about RINVOQ® ▼ (upadacitinib) for Patients

- This card contains important safety information you should be aware of – before and during treatment with RINVOQ®.
- Read the patient information leaflet for more information.

Risk of infections

RINVOQ® may make an existing infection worse or increase the chance of you getting a new infection – for example tuberculosis (TB), shingles or viral hepatitis.

Tell your doctor straight away if you notice signs of infection, such as:

- Fever, sweating, chills, weight loss, or a cough that won't go away – these may be signs of TB
- Painful skin rash with blisters – this may be a sign of shingles
- Feeling tired or short of breath – this may be a sign of pneumonia.

Vaccines – used to help prevent infections

Live vaccines (for example the influenza vaccine nasal spray) should not be given during RINVOQ® treatment, or just before starting RINVOQ® treatment.

Before being given any vaccines, talk to your doctor – they will know which vaccines you should **not** be given before or during treatment with RINVOQ®.

Cholesterol

Your doctor will check your cholesterol levels while you are taking RINVOQ® – this is to decide if you need to start a medicine to lower your cholesterol levels.

Contraception, pregnancy, and breast-feeding

RINVOQ® must not be taken during pregnancy.

- Use effective contraception while taking RINVOQ® – and for 4 weeks after your last dose. Talk to your doctor about effective contraception.
- Tell your doctor straight away if you wish to become pregnant, or if you become pregnant.
- Do not breast-feed while using RINVOQ®.

Blood clots in veins (DVT) or lungs (PE)

Tell your doctor straight away if you get signs of DVT or PE, such as a painful swollen leg or chest pain.