

Patient Card

Keep this card with you (or caregiver) all the time

Safety Information about RINVOQ[®] ▼ (upadacitinib) for patients

- This card contains important safety information you should be aware of – before and during treatment with RINVOQ[®].
- Read the patient information leaflet for more information.

Your name:

.....
Consultant's name – who prescribed RINVOQ®:

.....
Consultant's phone number:

.....
The date you started RINVOQ®:



.....
www.gov.uk/mhra

Version 3.0. UK-UPAD-220290/April 2023

GLUE TO FRONT COVER

Risk of infections

RINVOQ® may make an existing infection worse or increase the chance of you getting a new infection – for example tuberculosis (TB), shingles or viral hepatitis.

Tell your doctor straight away if you notice signs of infection, such as:

- Fever, sweating, chills, weight loss, or a cough that won't go away – these may be signs of TB.
- Painful skin rash with blisters – this may be a sign of shingles.
- Feeling tired or short of breath – this may be a sign of pneumonia.

Vaccines used to help prevent infections

Live vaccines (for example influenza vaccine by nasal spray, varicella, measles/mumps/rubella) should not be given during RINVOQ® treatment, or just before starting RINVOQ® treatment.

Before being given any vaccines, talk to your doctor – they will know which vaccines you should not be given before or during treatment with RINVOQ®.

Risk of a hole in your bowel

RINVOQ may increase your risk of a hole in your bowel especially if you have Crohn's disease. Tell your doctor straight away if you have unexplained or unexpected stomach pain.

Risk of heart disease

Treatment with RINVOQ® was associated with increases in cholesterol (blood fat). Your doctor will check your cholesterol levels while you are taking RINVOQ®.

Tell your doctor straight away if you notice symptoms such as chest pain or tightness since this may be a symptom of heart disease.

GLUE TO BACK COVER

Contraception, pregnancy, and breast-feeding

RINVOQ® must not be taken during pregnancy.

- Use effective contraception while taking RINVOQ® – and for 4 weeks after your last dose. Talk to your doctor about effective contraception.
- Tell your doctor straight away if you wish to become pregnant, or if you become pregnant.
- Do not breast-feed while using RINVOQ®.

Risk of blood clots in veins or lungs

Blood clots in veins or lungs have been observed with RINVOQ®. Tell your doctor straight away if you get signs of blood clots in veins or lungs, such as a painful swollen leg, shortness of breath, or chest pain.

Risk of cancers

RINVOQ® may potentially increase your risk of developing cancers particularly skin cancer.

Let your doctor know if you notice any change in the appearance of an area on your skin, or notice any new growth on your skin.

Show this card to any healthcare professional involved in your medical care – for example, your dentist or an emergency doctor.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See <https://yellowcard.mhra.gov.uk> for how to report side effects or via the Yellow Card app from the Google Play or Apple app Store.