

My information

Name:

Date of birth

DD / MM / YYYY

Phone number:

Emergency contact (name):

Emergency contact
(phone number):

NHS number:

My treatment details

Please complete this section or ask your doctor to do it.

Isatuximab recommended dose of 10 mg/kg and dosing schedule:

Cycle 1 (28 days): Days 1, 8, 15 and 22 (weekly)

Cycle 2 (28 days) and beyond: Days 1 and 15 (every 2 weeks)

Start date: DD/MM/YYYY

End date: DD/MM/YYYY

It is important to record the end date of your treatment because isatuximab may interfere with the indirect antiglobulin test (indirect Coombs test) for approximately 6 months after the last infusion.

My blood results

Before starting isatuximab, the results of my blood test collected on DD / MM / YYYY were:

Blood type:

A B AB O Rh+ Rh-

The result of my indirect antiglobulin test (indirect Coombs test) was:

Negative Positive for the following antibodies:

Other

Contact details of institution
where blood test was performed:

Phone number

My haematologist's information

In case of emergency, or if you find this card, please contact my haematologist using the details below.

Haematologist's name:

Haematologist's phone number:

Name of hospital:

MAT-GB-2000062(v2.0)

Date of prep: June 2021

MHRA approval: June 2021

SARCLISA[®]▼ (isatuximab)

Patient card

For patients receiving isatuximab

- Provide this card to healthcare providers before blood transfusion
- Keep this card with you at all times and until 6 months after the last dose of isatuximab
- If you notice any side effects, talk to your doctor or pharmacist
- ▼ This medicine is subject to additional monitoring. This will allow for the identification of new safety information. You can help by reporting any side effects you may get. It is easiest and quickest to report side effects online via the Yellow Cards website - www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. Alternatively, pre-paid Yellow Cards for reporting are available by writing to FREEPOST YELLOW CARD (no other address details necessary) or by emailing yellowcard@mhra.gov.uk. Side effects should also be reported to SANOFI on Tel: 0800 090 2314 or via email: UK-drugsafety@sanofi.com. By reporting side effects, you can help to provide more information on the safety of this medicine
- For further information about isatuximab, you can consult the patient information leaflet (PIL).

Warning message for healthcare professionals treating the patient at any time, including in emergency situations

- Please note this patient is receiving treatment with isatuximab for the treatment of multiple myeloma
- This patient card contains important safety information that you need to be aware of before, during, and after treatment with isatuximab
- Isatuximab can bind to CD38 on red blood cells (RBCs) and is associated with a risk of interference with blood typing (positive indirect Coombs test), which may persist for approximately 6 months after the last isatuximab infusion
- To avoid potential problems with RBC transfusion, you should perform blood type and screen tests prior to the first infusion of isatuximab. Phenotyping may be considered as per local practice
- If treatment with isatuximab has already started and in the event of a planned transfusion, you should notify the blood bank that the patient is receiving isatuximab and alert them to the risk of interference with the indirect antiglobulin test (indirect Coombs test)
- For additional information about isatuximab, please refer to the Summary of Product Characteristics (SmPC).