

Information for pharmacists dispensing Librachrom 2% w/v eye drops solution PL35533/0132

This product uses a different type of eye-dropper to those supplied with other non-prescription 'Pharmacy' category eye drops.

It is important to draw patients' attention to Step 6 in the administration process summarised below, as patients may not be familiar with this. In this step, after use of the dropper and prior to recapping, the bottle should be shaken once in a downwards direction to remove any residual liquid on the tip.

This step is important to minimise leaching of antimicrobial silver ions into any residual drop prior to next use and should be emphasised to patients prior to dispensing.

Please ensure the patient has read and fully understood the information on the leaflet contained within the packaging and address any questions they have.

If the patient develops any eye irritation or if the itching, redness or swelling gets worse, advise them to stop using this medicine and contact their doctor immediately.


Patient instructions for use

Before instillation of the eye drops:

- Wash your hands before opening the bottle.
- Do not use this medicine if you notice that the tamper-proof seal on the bottle neck is broken before you first use it.
- When using for the first time, before delivering a drop to the eye, you should first practice using the dropper bottle by squeezing it slowly to deliver one drop into the air, away from the eye.
- When you are confident that you can deliver one drop at a time, choose the position that you find most comfortable for the instillation of the drops (you can sit down, lie on your back, or stand in front of a mirror).

Instillation:

1. Hold the bottle directly below the cap and turn the cap to open the bottle. Do not touch anything with the tip of the bottle to avoid contamination of the solution.
4. Blink a few times so that the drop spreads over the eye.



2. Tilt your head backwards and hold the bottle above your eye.



3. Pull the lower eyelid down and look up. Squeeze the bottle gently in the middle and let a drop fall into your eye. **Please note that there might be a few seconds delay between squeezing and the drop coming out. Do not squeeze too hard.** If a drop misses your eye, try again. If you are not sure how to administer your medicine, ask your pharmacist.



5. Repeat the instructions 2. – 4. to deliver a drop into the other eye if needed.

6. After use and prior to recapping, the bottle should be shaken once in a downwards direction, without touching the dropper tip, in order to remove any residual liquid on the tip. This is necessary in order to ensure delivery of subsequent drops.



7. At the end of the 28-day in-use shelf life of the medicine, there will be some eye drops left in the bottle. Do not use the eye drops for longer than 28 days after first opening the bottle.

Reporting of Side Effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the product's package leaflet. You can also report side effects directly via the Yellow Card Scheme (Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store). By reporting side effects you can help provide more information on the safety of this medicine.

Last revised: 09/2020

1010489 - LP1.5

ASPIRE[®]
P H A R M A

Aspire Pharma Limited	
Product name	Librachrom
Size:	10ml
PL/PA no:	35533.0132
Type:	Leave Piece
Artwork dimensions (mm):	148mm x 210
Reason for request:	Added product code
Version no:	1.5
Date of revision:	15.9.20
Font(s):	Strada Pro
A/W software:	Indesign CC

