FRONT SIDE

Treatment start date and contact details

Date of most recent infusion:_____

Date of first infusion:_____

Patient's Name:_____

Doctor's Name:_____

Doctor's contact details:_____

Make sure you have a list of all your medicines when you see a health care professional.

Please talk to your doctor or nurse if you have any questions about the information in this card.

12345678 SZ00000AC000

Rixathon Alert Card for patients with non-oncology diseases

Why have I been given this card?

This medicine may make you more likely to get infections. This card tells you:

- What you need to know before having Rixathon
- What the signs of an infection are
- What to do if you think you might be getting an infection.

It also includes your name and doctor's name and phone number on the back.

What should I do with this card?

- Keep this card with you all the time – such as in your wallet or purse.
- Show this card to any doctor, nurse or dentist you see – not just the specialist who prescribes your Rixathon.

12345678 SANDOZ

4

This side will be visible when folded

1

This side will be visible when folded

BACK SIDE

Keep this card with you for 2 years after your last dose of Rixathon. This is because side effects can develop several months after you have had treatment.

When should I not have Rixathon?

Do not have Rixathon if you have an active infection or a serious problem with your immune system.

Tell your doctor or nurse if you are taking or have previously taken medicines which may affect your immune system this includes chemo-therapy.

What are the signs of getting an infection?

Look out for the following possible signs of infection:

- Fever or persistent cough
- Weight loss
- Pain without injuring yourself
- Feeling generally unwell or listless.

2

If you get any of these, tell a doctor or nurse straight away.
You should also tell them about your Rixathon treatment.

What else do I need to know?

Rarely Rixathon can cause a serious brain infection, called "Progressive Multifocal Leukoencephalopathy" or PML. This can be fatal.

- Signs of PML include:
  - Confusion, memory loss or problems thinking
  - Loss of balance or a change in the way you walk or talk
  - Decreased strength or weakness on one side of your body
  - Blurred vision or loss of vision.

If you get any of these, tell a doctor or nurse straight away. You should also tell them about your Rixathon treatment.

Where can I get more information?

See the Rixathon package leaflet for more information.

3

Artwork Proof Box:

Variation:	Technical changes to the PAC
Proof no:	004.0
Date prepared:	02/04/2026
Font size:	7.5 pt
Fonts:	FuturaEF EL
Dimensions:	110x85 mm
Technical data:	
SKUs:	00000000
SZ codes:	SZ00000AC000
Item codes:	00000000
Pharma codes:	00000000

Technical Colours:	
	Legend:
	Cutting:
	TextArea
	Guides:
Colours:	
	PANTONE 541 C
	Black

SANDOZ

Signature Panel: