	Front	
Important Safety Alerts for Patients taking Bosentan Tablets (bosentan)  This card contains important information about Bosentan Tablets. Please read this card carefully before starting your treatment with Bosentan Tablets.  Your name:  Prescribing doctor:  If you have questions about Bosentan Tablets ask your doctor.  Cipla Europe NV 21065065	Contraception Do you currently use or take contraceptives? Yes No If Yes, write the names of these here: Take this card to your doctor or your gynaecologist at your next visit and he/she will be able to advise you on whether you need to use additional or alternative contraceptive methods.	If you are a woman of child bearing age read this page carefully Pregnancy Bosentan Tablets may harm the development of the foetus. Therefore, you must not take Bosentan Tablets if you are pregnant and you must also not become pregnant while taking Bosentan Tablets.  Moreover, if you are suffering from pulmonary hypertension disease, the occurrence of a pregnancy can severely deteriorate the symptoms of your disease. If you suspect you may be pregnant, tell your doctor or gynaecologist.  Contraception Birth control based on hormones — such as oral contraceptives or birth control pills,
	Back	
ormone injections, implants, or birth control skin patches don't reliably prevent regnancy in women who are taking Bosentan Tablets. You need to use a barrier orm of birth control – like a condom, diaphragm or vaginal sponge – in addition to ny of these kinds of hormonal birth control. Be sure to discuss any questions you nay have with your doctor or your gynaecologist – complete the details on the back of this card and take it to your doctor or gynaecologist at your next visit. You should have a pregnancy test before initiation of Bosentan Tablets and every nonth during the treatment even if you think that you are not pregnant.	Blood Test for Liver Function Some patients taking Bosentan Tablets were found to have abnormal liver function tests. During treatment with Bosentan Tablets, your doctor will arrange for regular blood tests to check for changes in your liver function. Remember to have your liver blood test every month. After an increase in dose, an additional test will be done after 2 weeks.	Date of first monthly test: