

Patient Alert Card

Please keep this card with you at all times

This non-promotional material is intended for patients prescribed **Tolvaptan Teva**

TOL-GB-NP-00013
Version 2

Date of Preparation: December 2023
Date of MHRA Approval: January 2024

Patient's name:

Date Tolvaptan Teva first prescribed:

Doctor's name:

Treatment Centre name:

Treatment Centre contact number:

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse.

This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard.

By reporting side effects, you can help provide more information on the safety of this medicine. Adverse events may also be reported to Teva UK Limited via email to medinfo@tevauk.com or via phone on +44 (0) 207 540 7117.

Important safety information for patients

Tolvaptan Teva can cause your liver not to work properly.

Your doctor will arrange blood tests for liver function testing:

- Before starting treatment with Tolvaptan Teva
- Monthly for the first 18 months of treatment
- Then every 3 months thereafter

Therapy should be interrupted or discontinued if significant increase of liver enzymes and/or clinical symptoms of liver injury persist.

Inform your doctor immediately if you experience any of these signs that could indicate potential liver problems:

- Tiredness
- Loss of appetite
- Pain in the abdomen
- Dark urine
- Yellowing of skin or eyes (jaundice)
- Nausea
- Vomiting
- Fever
- Itching of your skin
- Flu-like syndrome (joint and muscle pain with fever)

Tolvaptan Teva can cause severe dehydration.

Tolvaptan Teva can increase your urine production which may result in excessive water loss and dehydration.

Drink plenty of fluids to avoid dehydration and consult your doctor if you are unable to drink or if you experience any of these symptoms.

Symptoms of dehydration can include:

- Increased thirst
- Dark yellow and strong-smelling urine
- Feeling tired
- Feeling dizzy or lightheaded
- Decreased urination
- Dry mouth, lips, eyes or skin

If dehydration is left untreated, it can become severe.

Severe dehydration is a medical emergency and requires immediate medical attention.

Symptoms of severe dehydration can include:

- Unusual tiredness
- Weak/rapid pulse
- Confusion
- Dizziness
- Not urinated all day
- Fits (seizures)

If you experience any of these symptoms, contact your doctor, call 999 or go to A&E immediately to seek medical advice.

For more information regarding Tolvaptan Teva, please contact Teva UK Limited Medical Information on Tel: 0207 540 7117 or medinfo@tevauk.com