

In case of an adverse reaction or pregnancy, your doctor may recommend an accelerated elimination procedure in order to speed up the elimination of AUBAGIO from your body.

This procedure consists of administration of 8 g cholestyramine three times daily (or 4 g if 8 g is not well tolerated), or 50 g activated charcoal twice daily for 11 days. The efficacy of this procedure is to be confirmed with blood tests prescribed by your doctor.

Please read the AUBAGIO package leaflet for more information.

For further information or to report an adverse event, please contact your doctor.

Please show this card to any doctor or healthcare professional involved in your medical care (e.g. in case of unexpected problems that may involve new doctors).

AUBAGIO is for treating adult patients with relapsing-remitting multiple sclerosis. This drug may affect your liver function. It may also affect blood cell counts and the immune system, and may increase the risk of infections/serious infections. Do not take AUBAGIO if you are, think you may be, or plan to become pregnant.

Once-daily 
AUBAGIO 
(teriflunomide) 14 mg tablets

SANOFI GENZYME 

PATIENT CARD

This Patient Card contains important safety information you need to be aware of when receiving treatment with AUBAGIO.

▼This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get to:
www.mhra.gov.uk/yellowcard

If you have any of the following symptoms, please contact your doctor immediately:

- Unexplained nausea
- Vomiting
- Stomach pain
- Your skin or the whites of your eyes turn yellow
- Dark urine
- Any signs or symptoms of an infection.

If you are a woman of childbearing potential, you should not be pregnant before starting AUBAGIO.

Your doctor may ask for a precautionary pregnancy test before issuing your AUBAGIO prescription. In addition, you should use an effective contraceptive method, agreed upon with your doctor, while you are on therapy.

AUBAGIO is not expected to adversely impact efficacy of oral hormonal contraceptives. You should inform your doctor before changing or stopping contraceptive measures during treatment with AUBAGIO.

If there is any possibility you may be pregnant, please contact your doctor immediately.

You should also discuss with your doctor if you plan to or are breastfeeding.

Please ensure you perform any follow up blood tests and blood pressure check-ups, as arranged by your doctor. Please also make sure you have a list of all your other medicines and medical conditions with you at every visit to a healthcare professional.

Patient's name:

.....

Date AUBAGIO first prescribed: __ __ / __ __ / __ __ __ __

.....

Centre name:

.....

Treating doctor's name:

.....

Treating doctor's phone number:

.....