ī	ATIENT ALERT CARD	
	bigatran etexilate hard capsules	E
•	This card should be with you / the caregiver at all times Make sure to use the latest version	
l		
ì		
1		
		E
' -		
1		
· I		
1		
1		i
1		
1		
T.		
1		
1		
1		
Ī		
1		ı
1		
1		
1		
1		
	STADA	

<u>-</u>	
Your	Patient / Caregiver of a paediatric patient, / your child's doctor has initiated treatment with
Dabig	gatran etexilate. In order to use this medicine safely, se consider the important information in the package
As th abou you /	is patient alert card contains important information tyour / your child's treatment, this card should be with your child at all times to inform healthcare professional tyour / your child's intake of this medicine.
	gatran etexilate Information for Patients / Caregiver nediatric patients
• Da	t your / your child's treatment abigatran etexilate thins the blood. It is used to treat sisting blood clots or to prevent the formation of ungerous blood clots.
Da of	ollow your / your child's doctor's instructions when takinabigatran etexilate. Never skip a dose or stop the intake this medicine without talking to your / your child's actor.
yo • Inf Da pr	form your / your child's doctor about all medicines you / ur child are / is currently taking. form your / your child's doctor about the intake of abigatran etexilate before any surgery / invasive ocedure.
foo of	abigatran etexilate capsules can be taken with or withou od. The capsule should be swallowed whole with a glas water. The capsule must not be broken or chewed and e pellets must not be emptied from the capsule.
Takin Spea child swell palen of gu flow pink blood • In se • Do	n to seek medical advice go Dabigatran etexilate may increase the risk of bleeding k to your / your child's doctor immediately if you / your experience(s) signs and symptoms of bleeding such as ling, discomfort, unusual pain or headache, dizziness, ness, weakness, unusual bruising, nosebleeds, bleeding ms, unusual long bleeding cuts, abnormal menstrual or vaginal bleeding, blood in the urine which may be or brown, red/black stools, coughing up blood, vomiting d or coffee ground like material. case of fall or injury, especially if the head is hit, urgent tek medical advice.
ex	lking to your / your child's doctor, if you / your child sperience(s) heartburn, nausea, vomiting, stomach scomfort, bloating or upper abdominal pain.
	gatran etexilate Information for Healthcare essionals
thi Da su In	abigatran etexilate is an oral anticoagulant (direct rombin inhibitor). abigatran etexilate may need to be stopped in advance or a stopped in a
ad re	specific reversal agent (idarucizumab) is available for fult patients. The efficacy and safety of the specific versal agent idarucizumab have not been established in dediatric patients.
effec of Pro	etails and more advice to antagonise the anticoagulant t of Dabigatran etexilate please refer to the Summary oduct Characteristics of Dabigatran etexilate and cizumab.
ad	abigatran etexilate is mainly eliminated by the kidneys; lequate diuresis must be maintained. Dabigatran exilate is dialyzable.
	se complete this section or ask your / your child's or to do it.
Patie	ent Information
Nam	e of the patient
D-4-	of birth
Date	of birth
Indic	eation for anticoagulation
Dose	e of Dabigatran etexilate
· 	
' 	