Please remember:

- BLINCYTO may cause neurological events including:
 - » Shaking (tremor)
- » Confusion
- » Disturbances of brain function (encephalopathy)
- Difficulty in communicating (aphasia)
- » Seizure (convulsion)
- Please call your doctor or nurse immediately if you experience any of these symptoms
- Please travel home safely and do not drive or operate moving vehicles/heavy machinery, or engage in hazardous activities whilst receiving BLINCYTO
- If there are problems or unexpected stopping of the infusion pump, call your doctor or nurse immediately. Please don't try to fix any issues yourself.

 Version 3.0 Approved: January 2022

Patient Card



Please show this card to all emergency and healthcare providers

Information about BLINCYTO®▼(blinatumomab)

My name is	Before providing any treatment, please call my prescribing physician at the number below. any medical evaluations are undertaken, please provide copies of all medical records, including ar treatments and/or test results, to the doctor(s) named below.				
In Case of Emergency (ICE) contact		Name	Hospital	City	Phone Number
	Haematologist				
	Oncologist				
	Haematology Nurse				
I am being treated with BLINCYTO, a treatment for B-precursor acute lymphoblastic leukaemia, which can lower my immune system.	▼This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App				

Store. By reporting side effects you can help provide more information on the safety of this medicine.

I started treatment on .

Version 3.0 Approved: January 2022